


HERBICIDE ROTATION - 2014

Resistance to a pesticide group will reduce the effectiveness of pesticides in that group over time. Rotation of herbicides and other pesticides is an important measure to delay the onset of resistance to any one pesticide group or mode-of-action.

By rotating herbicides, the risk of developing widespread resistance over a field drops. The illustration below gives a relative ranking of risk of resistance developing from repeated use. The top of the triangle indicates groups that may develop resistance quickly and those at the bottom of the triangle have demonstrated a low risk of resistance developing over long-term use. Be aware that low risk does not mean no risk, since weeds have developed resistance to herbicides in these groups as well.

Herbicide Resistance Development Risk based on Number of Applications

Classification of herbicide groups numbers by risk of selection for weed resistance ('Other': all other herbicide groups that pose a low or moderate risk)
Adapted from Beckie, H. J., 2006 *Herbicide Resistance Weeds: Management Tactics & Practices*; Weed Technology Vol. 20 Issue 3 (July-September) pp. 793-814



Rotation of Herbicides

OTHER GROUPS

Linuron (7), Liberty (10), Amitrol (11), Battalion* (15), Dual II Magnum (15), Frontier Max (15), Kerb (15), Primextra II Magnum* (15), Distind* (19), Casaron (20), Gramoxone (22), Diquat (22), Infinity* (27), Tundra (27)*, Velocity m3(27)*

GROUP 14

(Grass & Broadleaf Control)
Aim, Authority, Authority Charge, Blazer, CleanStart*, Flumioxazin, Heat, Reflex

GROUP 9

(Grass & Broadleaf Control)
CleanStart*, Eclipse*, Glyphosate, glyphosate+dicamba*, Florasulam+glyphosate*

GROUP 8

(Grass & Broadleaf Control)
Avadex, Avenge, Eptam, Fortress*

GROUP 1 (Grass Control)

Axial, clethodim, Clodinafop, Equinox, Fenoxaprop, Poast Ultra, quizalofop, tralkoxydim, Traxos, Tundra*

GROUP 2 (Grass & Broadleaf Control)

Accent, Adrenalin*, Ally, Altitude FX/FX 2*, Ares, Barricade II, Broadband, Battalion*, Express Pro, Everest GBX*, flucarbazone, Frontline*, Frontline 2,4-D*, imazamethabenz, imazethapyr, metsulfuron, Muster, Odyssey, Odyssey DLX*, Option, Pinnacle, Prepass*, Prism, Simplicity, Solo, Spectrum*, Stellar*, Thifensulfuron/tribenuron, tribenuron, Triton C* and K*, Velocity m3, Viper ADV

GROUP 3

(Grass & Broadleaf Control)
Edge, Fortress*, Trifluralin

GROUP 4 (Broadleaf Control)

2,4-D, 2,4-DB, Adrenalin*, Altitude FX*/FX 2, Axial Xtreme*, Battalion*, Bromoxynil + 2,4-D ester*, Bromoxynil + MCPA ester*, Curtail M, Dicamba, Dicamba + Mecoprop + MCPA, Dichlorprop + 2,4-D, Distinct*, DyVel, DyVel DSp, Eclipse*, Everest GBX*, Fluoxypyr+2,4-D, Frontline 2,4-D*, glyphosate+dicamba, Grazon, Lontrel, MCPA, MCPB+MCPA, Mecoprop, Optica trio, Prestige XC, Spectrum*, Stellar*, Tordon 22K, Triton C* and K*, Trophy



GROUP 6

(Broadleaf Control)
Basagran, Bromoxynil, Bromoxynil+2,4-D ester*, Bromoxynil+MCPA ester*, Horizon BTM*, Infinity*, Tundra*, Velocity m3* Viper ADV*

GROUP 5

(Grass & Broadleaf Control)
Atrazine, Primextra II Magnum*, Metribuzin, Simazine, Velpar

*Some products contain more than one active ingredient and therefore may appear in more than one group.

MANITOBA AGRICULTURE FOOD AND RURAL DEVELOPMENT <http://www.gov.mb.ca/agriculture/index.html>

\$9.82 (including GST)

GUIDE TO

FIELD CROP PROTECTION

2014

Weeds
Plant Diseases
Insects

Introduction

Weed Control

Disease Control

Seed Treatments

Insect Control

THIS PUBLICATION IS ONLY A GUIDE. ALWAYS REFER TO THE
PRODUCT LABEL FOR APPLICATION DETAILS AND PRECAUTIONS.

Manitoba 

Crop Protection Companies

AgWest Inc.
1-877-924-9378

Arysta LifeScience
www.arystalifescience.ca
1-866-761-9397

BASF
www.ag solutions.ca
1-877-371-2273

Bayer CropScience
www.bayercropscience.ca
1-888-283-6847

Brenntag Canada Inc.
www.brenntag.ca
1-204-233-3416

Cheminova
www.cheminova.com
1-888-316-6260

Chemtura
1-800-350-1745

Crop Production Services/
Loveland Products
www.cpsagu.com
1-800-328-4678

Degesch America Inc.
www.degeschamerica.com
1-800-330-2525

Dow AgroSciences
www.dowagro.ca
1-800-667-3852

E.I. duPont
www.dupont.ca/ag/
1-800-667-3925

Engage Agro
www.engageagro.com
1-866-613-3336

Gowan Company
www.gowanco.com
1-800-883-1844 ext. 2

Great Northern Growers
www.gng.ag
1-866-727-5226

Hedley Technologies
www.hedleytech.com
1-888-476-4473

IPCO
www.ipco.ca
(204) 233-3461

MANA Canada
www.manainc.ca
East: 903-304-1168
West: 1-855-264-6262

Monsanto
www.monsanto.ca
1-800-667-4944

Norac Concepts
www.noracconcepts.com
1-519-821-3633

Nufarm
www.nufarm.ca
1-800-868-5444

Peacock Industries
www.grasshoppercontrol.com
1-306-225-4691

Syngenta
www.farmassist.ca
1-877-964-3682

United Agri Products
www.uap.ca
1-800-561-5444

United Phosphorus Inc.
www.upi-usa.com
1-416-567-4102

Valent BioSciences
www.valent.com
1-800-898-2536

For more information on weeds, diseases and insects, visit the Manitoba Agriculture, Food and Rural Development website at www.manitoba.ca/agriculture

Manitoba Agriculture, Food and Rural Development GO Offices

Please dial area code 204 before the numbers below.

Central Plains GO Team

Carberry 834-8815
Gladstone 385-6633
Portage la Prairie 239-3352

Eastman GO Team

Beausejour 268-6094
St. Pierre 433-7749
Steinbach 346-6080
Vita 425-5050

North Interlake GO Team

Arborg 376-3300
Ashern 768-2782
Lundar 762-5649

North Parkland GO Team

Dauphin 622-2007
Roblin 937-6460
Ste Rose 447-4032

Pembina GO Team

Killarney 523-5260
Pilot Mound 825-3512
Somerset 744-4050

Red River GO Team

Altona 324-2804
Carman 745-5610
Morden 822-5461
Morris 746-2312

South Interlake GO Team

Teulon 886-2696

South Parkland GO Team

Hamiota 764-3010
Minnedosa 867-6572
Russell 773-5130

Southwest GO Team

Brandon 726-6482
Melita 522-3256
Souris 483-2153
Virden 748-4770

Urban GO Team

Winnipeg 945-4521

Valleys North GO Team

Swan River 734-3417
The Pas 627-8255

Manitoba Weed Supervisor Offices

Please dial area code 204 before the numbers below.

Central Region

Baldur (M/Th) 535-2176
Cartwright (T/Th) 529-2363
Crystal City (M/W/F) 873-2103
Holland 526-2732
Lettelier 746-0395
MacGregor 685-2050
Sanford 736-2331
Somerset (T/W/F) 744-2232

Eastern/Interlake Region

Arborg 376-3313
Beausejour 268-1764
Dugald 981-4098
Selkirk 785-3638
St. Pierre-Jolys 326-0280
Stonewall 467-4704
Vita 425-5050

Western Region

Carberry 834-6600
Dauphin 638-7050
Deloraine 747-2586
Glenboro 827-2602
Hamiota 764-2128
Hartney 858-2590
Killarney 523-5275

Miniota 567-3683
Pierson 634-2419
Reston 877-3327
Swan River 734-3344
Virden 851-0409

EMERGENCY NUMBERS

POISON CONTROL CENTRE
1-855-776-4766

PESTICIDE SPILL LINE
1-204-945-4888

Prepared by Manitoba Agriculture, Food and Rural Development in co-operation with Saskatchewan Agriculture and Food.




TABLE OF CONTENTS

1. Introduction.....	10
How to Use This Book.....	10
Product Labels and PCP Numbers.....	10
Safe Use of Herbicides, Fungicides and Insecticides	11
Protecting Yourself from Exposure to Herbicides, Fungicides, and Insecticides	11
Protective Clothing.....	12
Protecting Your Eyes, Face and Lungs.....	13
Avoiding Spray Drift	13
Herbicide Efficacy With Low-drift Nozzles.....	14
Handling a Drift Complaint.....	14
Mixing Pesticides.....	14
Container Disposal.....	15
Sprayer Tank Cleaning.....	15
How to Identify Crop and Weed Leaf Stages.....	17
Pesticide Formulations and Companies	20
Key to product pages.....	27
 2. Weed Control	 30
Integrated Weed Management.....	30
Making Spray Decisions	30
Weed Resistance to Herbicides.....	38
Adjuvants.....	42
Crop and Herbicide Recommendation Tables	43
Special Weed Problems	56
Soil Residual Herbicides	61
Effect of Rainfall on Herbicide Efficacy	64
Products Available as Prepackaged Tank Mixes.....	65
Herbicide Product Pages	66

3. Plant Disease Control	307
Integrated Plant Disease Management	307
Effects of Weather	310
Resistance Management	310
Fungicide Mode of Action	311
Foliar Fungicide Tables	314
Foliar Fungicide Product Pages	319
Seed Treatment Tables	384
Seed Treatment Product Pages	388
 4. Insect Control	 453
Insect Control Decisions	453
Field Scouting	453
Insecticides and Bees	454
Insecticide Poisoning in Humans	455
Resistance of Insects to Insecticides	455
Field Scouting and Insect Management Charts (for individual crops)	457
Insect Control in Stored Grain	494
Insecticide Product Pages	498
 Reference Material	
Pesticide Index	5
Crop Protection Companies	Inside Cover
Emergency Numbers	Inside Cover
Agricultural Offices	Inside Cover
Metric Conversion Factors	9
Tank Mix Cross-Reference Chart	Insert, Inside Back Cover
Forage Crops Chart - Recommended Herbicides	Insert, Inside Back Cover

Pesticide Index

Herbicide Directory

2,4-D.....	66	Clovitox Plus (see 'MCPB/MCPA').....	221
2,4-DB.....	69	Cobutox 625 (see '2,4-DB').....	69
AAtrex Liquid.....	70	Compitox (see 'mecoprop-p').....	223
Accent.....	72	Cordon (see 'fenoxaprop').....	156
Accurate (see 'metsulfuron').....	227	Cougar (see 'fenoxaprop').....	156
Achieve (Liquid Achieve) (see 'tralkoxydim').....	279	Credit 45 (see 'glyphosate').....	178
Achieve Liquid Gold.....	74	Crush'R Plus (see 'glyphosate').....	178
Adrenalin SC.....	75	Curtail M.....	125
Aim.....	77	Deploy (see 'thifensulfuron/tribenuron').....	272
Ally Toss-N-Go (see 'metsulfuron').....	227	Desica (see 'diquat').....	134
Altitude FX.....	78	Dicamba.....	126
Altitude FX2.....	80	Dicamba/Mecoprop/MCPA.....	130
Amitrol 240.....	82	Dichlorprop/2,4-D.....	132
Approve (see 'bromoxynil/2,4-D ester').....	112	Dichlorprop-D/ Dichlorprop DX (see 'dichlorprop/2,4-D').....	132
Arcs.....	83	Diquat.....	134
Arrow (see 'clethodim').....	119	Distinct.....	136
Assert 300 SC (see 'imazamethabenz').....	198	Dual II Magnum.....	137
Assert FL.....	85	DyVel.....	139
Assure II (see 'quizalofop').....	252	DyVel D5p.....	141
Attain XC (see 'fluroxypyr + 2,4-D').....	168	Eclipse III.....	143
Aurora (see 'clodinafop').....	122	Edge Granular.....	144
Authority.....	86	Embutox 625 (see '2,4-DB').....	69
Authority Charge.....	88	Enforcer D.....	145
Avadex Brands.....	88	Enforcer M.....	147
Avenge 200 C.....	92	Eptam 8-E.....	148
Avert (see 'imazamethabenz').....	198	Equinox.....	150
Axial/Axial BIA.....	93	Escort.....	151
Axial iPak.....	95	Estaprop XT (see 'dichlorprop/2,4-D').....	132
Axial Xtreme.....	96	Everest (see 'flucarbazone').....	164
Badge (see 'bromoxynil/MCPA ester').....	113	Everest GBX.....	153
Banvel II (see 'dicamba').....	126	Express Pro.....	155
Banvel VM (see 'dicamba').....	126	Express SG (see 'tribenuron').....	283
Barricade II.....	97	Fenoxaprop.....	156
Basagran Brands.....	99	FirstStep Complete (see 'tribenuron').....	283
Battalion.....	102	FlexStar GT.....	159
Bengal WB (see 'fenoxaprop').....	156	Florasulam + glyphosate.....	160
Bison (see 'tralkoxydim').....	279	Florasulam + MCPA.....	162
BlackHawk.....	104	Flucarbazone.....	164
Blazer.....	105	Flumioxazin.....	166
Bonanza (see 'trifluralin').....	286	Flurox 2,4 (see 'fluroxypyr + 2,4-D').....	168
Broadband.....	107	Fluroxypyr + 2,4-D.....	168
BroadSide (see 'thifensulfuron/tribenuron + MCPA').....	276	Focus.....	171
Bromotril (see 'bromoxynil').....	108	Foothills NG (see 'clodinafop').....	122
Bromoxynil.....	108	Fortress.....	172
Bromoxynil/2,4-D Ester.....	112	Frontier Max.....	174
Bromoxynil/MCPA ester.....	113	Frontline 2,4-D XC.....	176
Brotex 240 (see 'bromoxynil').....	108	Frontline XL (see 'florasulam + MCPA').....	162
Buctril M (see 'bromoxynil/MCPA ester').....	113	Gladiator (see 'imazethapyr').....	200
Bullwhip (see 'clodinafop').....	122	Glyfos (see 'glyphosate').....	178
Caliber 625 (see '2,4-DB').....	69	Glykamba (see 'glyphosate/dicamba').....	187
Casoron.....	116	Glyphosate.....	178
Centurion (see 'clethodim').....	119	Glyphosate/dicamba.....	187
Chateau (see 'flumioxazin').....	166	GoodHarvest (MPOWER).....	189
Cheminova Glyphosate (see 'glyphosate').....	178	Gramoxone.....	190
CleanStart.....	117	Grazon.....	191
ClearOut 41 Plus (see 'glyphosate').....	178	Harmony K.....	193
Clethodim.....	119	Harmony Max.....	193
Clodinafop.....	122	Harmony SG.....	195

Hawkeye Power (see 'dicamba')	126	Puma Advance (see 'fenoxaprop')	156
Heat	195	Pursuit (see 'imazethapyr')	200
Horizon NG (see 'clodinafop')	122	Quizalofop	252
Imazamethabenz	198	Reclaim	254
Imazethapyr	200	Refine M (see 'thifensulfuron/tribenuron + MCPA')	276
Impact	202	Refine SG (see 'thifensulfuron/tribenuron')	272
Inferno (see 'tribenuron')	283	Reflex	256
Inferno Duo	203	Reglone Desiccant (see 'diquat')	134
Infinity	205	Reglone Ion (see 'diquat')	134
Kerb	207	Restore II	257
Knockout Extra (see 'glyphosate')	178	Retain	259
Ko-Act	209	Reward	260
Koril 235 (see 'bromoxynil')	108	Rival (see 'trifluralin')	286
Ladder (see 'clodinafop')	122	Roundup brand products (R/T 540, Transorb HC, Ultra 2, WeatherMax) (see 'glyphosate')	178
Laji Plus (see 'glyphosate')	178	Rush 24 (see 'fluroxypyr + 2,4-D')	168
Leader (see 'bromoxynil/2,4-D ester')	112	Rustler (see 'glyphosate/dicamba')	187
Legend (see 'clodinafop')	122	Salvo (see '2,4-D')	66
Liberty 150 SN	209	Select (see 'clethodim')	119
Liberty 200 SN	212	Sencor (see 'metribuzin')	224
Linuron	213	Shadow RTM (see 'clethodim')	119
Logic M (see bromoxynil/MCPA ester)	113	Sharpshooter (see 'glyphosate')	178
Lontrel 360	216	Sharpshooter Plus (see 'glyphosate')	178
Lorox (see 'linuron')	213	Sierra (see 'flucarbazone')	164
Marengo (see 'tralkoxydim')	279	Signal (see 'clodinafop')	122
Matrix (see 'glyphosate')	178	Simazine	261
Maverick III (see 'glyphosate')	178	Simazine 480 (see 'simazine')	261
MCPA	218	Simplicity	263
MCPB/MCPA	221	Slam'R (see 'clodinafop')	122
Mecoprop-P	223	Solo	265
Metribuzin	224	Spectrum	266
Metsulfuron	227	Spike-Up (see 'tribenuron')	283
Mextrol 450 (see 'bromoxynil/MCPA ester')	113	StartUp (see 'glyphosate')	178
Momentum	230	Stellar	268
MPower Glyphosate (see 'glyphosate')	178	Sword (see 'dicamba/mecoprop/MCPA')	130
MPower HellCat (see 'fenoxaprop')	156	Tandem	269
MPower Kamikaze (see imazethapyr)	200	Target (see 'dicamba/mecoprop/MCPA')	130
MPower R (see 'thifensulfuron/tribenuron')	272	Tensile	271
MPower X (see 'tribenuron')	283	Thifensulfuron/tribenuron	272
MultiStar (see 'imazethapyr')	200	Thifensulfuron/tribenuron + MCPA	276
Muster Toss-N-Go	232	Thrasher (see 'bromoxynil/2,4-D ester')	112
NextStep NG (see 'clodinafop')	122	Thumper (see 'bromoxynil/2,4-D ester')	112
Nimble (see 'thifensulfuron/tribenuron')	272	Topline (see 'florasulam + MCPA')	162
Nuance (see 'tribenuron')	283	Topside (see 'MCPB/MCPA')	221
Nufarm Clodinafop (see 'clodinafop')	122	Tordon 22K	277
NuGlo (see 'glyphosate')	178	Touchdown Total (see 'glyphosate')	178
OcTtain XL (see 'fluroxypyr + 2,4-D')	168	Tracker XP (see 'dicamba/mecoprop/MCPA ester')	130
Odyssey	234	Tralkoxydim	279
Odyssey DLX	235	Traxion (see 'glyphosate')	178
Optica Trio	236	Traxos	281
Option 35 DF/Option 2.25 OD	238	Treflan (see 'trifluralin')	286
Oracle (see 'dicamba')	126	Tribenuron	283
Overdrive	240	TriCor (see 'metribuzin')	224
Pardner (see 'bromoxynil')	108	Trifluralin	286
Phantom (see 'imazethapyr')	200	Triton C	291
Pinnacle SG	241	Triton K	293
Poast Ultra	242	Trophy	295
PrePass XC (see 'florasulam + glyphosate')	160	Tropotox Plus (see 'MCPB/MCPA')	221
Prestige XC	245	Tundra	296
Primextra II Magnum	247	Turboprop (see 'dichlorprop/2,4-D')	132
Princep Nine-T (see 'simazine')	261	Ultim	298
Priority (see 'florasulam + glyphosate')	160	Valtera (see 'flumioxazin')	166
Prism SG	248	Vantage Plus Max II (see 'glyphosate')	178
Pulsar	250		

Varro	300	Viper ADV.....	305
Velocity m3.....	302	VMD 480 (see 'dicamba')	126
Velpar DF	304	WildCat (see 'fenoxaprop')	156
Vector (see 'glyphosate')	178	Wise-Up (see 'glyphosate').....	178
Vigil WB (see 'fenoxaprop').....	156	Yuma GL (see 'quizalofop').....	252

Foliar Fungicide Directory

Acapela	319	Overall 240 SC (see 'iprodione').....	345
Acrobat 50 WP.....	321	Palliser (see 'tebuconazole')	377
Allegro 500F.....	322	Parasol (see 'copper')	334
Astound.....	323	Penncozeb 75 DF (see 'mancozeb')	349
Azoxystrobin	324	Phostrol	350
Blanket	326	Pivot 418 EC (see 'propiconazole')	355
Bravo 500/Bravo Zn (see 'chlorothalonil').....	330	Priaxor DS.....	352
Bumper 418 EC (see 'propiconazole').....	355	Proline 480 SC	353
Cabrio Plus	327	Propel (see 'propiconazole')	355
Cantus WDG	328	Propiconazole.....	355
Caramba	329	Propulse	358
Chlorothalonil	330	Prosaro 250 EC	359
Contans WG	332	Quadris (see 'azoxystrobin').....	324
Copper.....	334	Quadris Top	361
Copper 53W/Copper Spray (see 'copper').....	334	Quash	362
Curzate 60 DF.....	336	Quilt	363
Cyazofamid	337	Ranman 400SC (see 'cyazofamid').....	337
Dithane Rainshield (see 'mancozeb').....	349	Reason 500SC	366
Echo 720/Echo 90DF (see 'chlorothalonil').....	330	Regalia Maxx	367
Folicur 432 F (see 'tebuconazole')	377	Revus	368
Folicur 250 EW (see 'tebuconazole')	377	Revus Top	369
Fontelis	338	Ridomil Gold (Bravo; SL/Bravo; 480EC; 480SL).....	369
Fuse (see 'tebuconazole')	377	Rovral Flo / Rovral RX (see 'iprodione').....	345
Gavel 75 DF	339	Scala SC	371
Headline DUO.....	340	Senator 70WP	372
Headline EC.....	341	Serenade Max / Serenade CPB	373
Inspire	344	Stratego 250EC	374
Iprodione	345	Tanos 50 DF.....	375
Kocide 2000 (see 'copper')	334	Tattoo C	376
Lance WDG.....	346	Tebuconazole	377
Luna Tranquility	348	Tilt 250E (see 'propiconazole')	355
MANA Azoxystrobin (see 'azoxystrobin')	324	Torrent 400 SC (see 'cyazofamid')	337
Mancozeb.....	349	Twinline.....	380
Manzate Pro-stick (see 'mancozeb').....	349	Vertisan.....	381
Nufarm Propiconazole (see 'propiconazole')	355	Zampro	383

Seed Treatment Directory

Acceleron DX-910	388	Cruiser Maxx Beans / Pulses	397
Acceleron Seed Treatment Package.....	389	Cruiser Maxx Vibrance Beans	397
Admire SPT (see 'imidacloprid').....	414	Cruiser Maxx Cereals / Cereals Commercial.....	399
Agrox B-2/Agrox CD	390	Cruiser Maxx Vibrance Cereals	399
Agrox FL	391	Cruiser Maxx D Potatoes	402
Alias 240 SC (see 'imidacloprid').....	414	DB-Red L.....	403
Alliance FL (see 'metalaxyl').....	424	DCT.....	404
Apron Advance/ Apron Maxx RTA/ Maxx RFC	392	Dividend XL RTA.....	406
Armour (see 'triticonazole')	444	EverGol Energy	407
Armour RTU (see 'triticonazole')	444	Gaucho CS FL.....	409
Belmont 2.7FS (see 'metalaxyl')	424	Gemini	410
Charter RTU (see 'triticonazole').....	444	General Storage Disinfectant.....	411
Confine	394	Heads Up Plant Protectant	412
Crown.....	395	Helix Vibrance	413
Cruiser 5FS.....	396	Helix XTra	413

Imidacloprid.....	414	Raxil PRO Shield	435
Insure Cereal	416	Raxil WW	436
Kocide 2000	418	Senator PSPT	437
Lumiderm	419	Solan MZ (see 'mancozeb')	420
Mancozeb	420	Sombrero 600 FS (see 'imidacloprid')	414
Maxim D / Liquid PSP/PSP/MZ PSP	421	StorOx	438
Maxim Quattro	422	Stress Shield 600	439
Mertect SC	423	Thiram 75WP	440
Metalaxyl	424	Titan ST (see 'Prosper FL')	428
Polyram 16D	426	Trilex AL	441
Potato ST 16 (see 'mancozeb')	420	Trilex EverGol	442
Proseed	427	Triticonazole	444
Prosper Evergol/Prosper FX	428	Tuberseal (see 'mancozeb')	420
Poncho 600 FS (see 'Prosper Evergol')	428	Vibrance 500 FS	446
Rancona Apex	430	Vibrance Maxx RTA/RFC	447
Rancona RS	431	Vibrance XL/Vibrance XL Co-Pack	448
Raxil MD	432	Vitaflo Brands	450
Raxil PRO	433		

Insecticide Directory

Actara	498	Malathion	522
Admire (see 'imidacloprid')	517	Malathion grain dust (see 'Stored Grain')	494
Alias (see 'imidacloprid')	517	Matador (see 'lambda-cyhalothrin')	519
Ambush (see 'permethrin')	527	Movento	523
Assail	499	MPOWER Krypton (see chlorpyrifos)	501
Beleaf	500	Nolo Bait	524
Chlorpyrifos	501	Nufos (see 'chlorpyrifos')	501
Citadel (see 'chlorpyrifos')	501	Oberon	525
Clutch	502	Orthene	526
Concept	503	Permethrin	527
Coragen	505	Perm-Up (see 'permethrin')	527
Cygon (see 'dimethoate')	510	Phostoxin (see 'Stored Grain')	494
Cypermethrin	506	Pounce (see 'permethrin')	527
Decis	507	Protect-IT (see 'Stored Grain')	494
Delegate	508	Pyrifos (see 'chlorpyrifos')	501
Dibrom	509	Pyrinex (see 'chlorpyrifos')	501
Dimethoate	510	Rimon	528
Dipel 2X DF	512	Ripcord (see 'cypermethrin')	506
Eco Bran	513	Sevin XLR	529
Endosulfan	514	Silencer (see 'lambda-cyhalothrin')	519
Entrust	515	Sluggo Professional	530
Fulfill	516	Success	531
Grapple / Grapple ₂ (see 'imidacloprid')	517	Tempo (see 'Stored Grain')	494
Imidacloprid	517	Thimet	532
Imidan	518	Thiodan (see 'endosulfan')	514
Insecto (see 'Stored Grain')	494	Thionex (see 'endosulfan')	514
Lagon (see 'dimethoate')	510	Up-Cyde (see 'cypermethrin')	506
Lambda-Cyhalothrin	519	Vydate	533
Lannate	521	Warhawk (see chlorpyrifos)	501
Lorsban (see 'chlorpyrifos')	501		

Table 1: Metric Conversion Factors* (Approximate)

<i>Metric Unit</i>	<i>Metric to Imperial Multiply By</i>	<i>Imperial Unit</i>	<i>Imperial to Metric Multiply By</i>	<i>Metric Unit</i>
LINEAR centimetre (cm)	x 0.39	inch	x 2.54	LINEAR centimetre (cm)
AREA square metre (m ²)	x 1.2	square yard	x 0.84	AREA square metre (m ²)
hectare (ha)	x 2.5	acres	x 0.4	hectare (ha)
VOLUME litre (L)	x 0.22	gallon	x 4.55	VOLUME litre
PRESSURE kilopascals (kPa)	x 0.14	psi	x 6.9	PRESSURE kilopascals (kPa)
WEIGHT gram (g)	x 0.035	oz	x 28.35	WEIGHT gram (g)
kilogram (kg)	x 2.2	lb	x 0.454	kilogram (kg)
AGRICULTURAL litres per hectare (L/ha)	x 0.089	gallons per acre	x 11.23	AGRICULTURAL litres per hectare (L/ha)
litres per hectare (L/ha)	x 0.357	quarts per acre	x 2.81	litres per hectare (L/ha)
litres per hectare (L/ha)	x 0.71	pints per acre	x 1.41	litres per hectare (L/ha)
millilitres per hectare (mL/ha)	x 0.014	fl.oz per acre	x 70.22	millilitres per hectare (mL/ha)
kilograms per hectare (kg/ha)	x 0.89	lb per acre	x 1.12	kilograms per hectare (kg/ha)
grams per hectare (g/ha)	x 0.014	oz per acre	x 70	grams per hectare (g/ha)
*EXAMPLE: To convert centimetres to inches, multiply by 0.39; conversely, to convert inches to centimetres, multiply by 2.54. CAUTION: Herbicide labels are in metric units only. Conversion between the Metric and Imperial system may result in confusion. It is recommended to use metric units only.				

Guide to Crop Protection 2014

For Reference Until December 31, 2014
This publication is updated annually and replaces
the 2013 and previous issues.

Introduction

How to Use This Book

This publication is only a guide. Always refer to the product label for application details and precautions. If the information in this publication differs from the label information, follow label instructions.

The *Guide to Crop Protection* is divided into four sections: (1) Weed Control; (2) Foliar Fungicides; (3) Seed Treatments; and (4) Insect Control.

To use the information in each of these sections, use the following process:

1. Turn to the charts at the beginning of each section. There is a set of charts for weeds, plant diseases and insect control. Select the chart for the crop you want or plan to grow. Use the chart to match your weed, disease or insect problems with the products available for that crop.
2. Once you have narrowed your product choices down to a few candidates, go to the recommendation section for that product. Products are listed alphabetically. Read the recommendations thoroughly for each product you are considering.
3. Read the product label attached to the container for detailed instructions on application.

This publication is intended to be used as a guide only. Information contained herein is that available at time of printing. While every effort has been made to ensure accuracy, the provincial government does not accept responsibility for label changes. When more than one trade name is listed, not all weeds or tank mixes may appear on all labels. Consult product labels attached to pesticide containers for final detailed instructions.

Certain recommendations in this publication are given in quantity of commercial product per acre (mL, L, g or kg/acre). Product labels are given in quantity of product per hectare (mL, L, g or kg/ha). To avoid application errors be sure to read and understand label recommendations.

The *Guide to Crop Protection* includes the most recent recommendations for weed, plant disease and insect

control in field and forage crops. These recommendations are based on the uses registered under the Pest Management Regulatory Agency's *Pest Control Products Act*. It is an offence under *The Pest Control Products Act* to apply any chemical in a manner not consistent with the product label. If you have any doubts regarding the instructions in this publication, or on the product label, contact the company representative, your local agricultural office or the Pest Management Regulatory Agency for further advice.

Product Labels and PCP Numbers

On each Product Page you will see a Registration or PCP number, so named because it is mandated by the *Pest Control Products Act*. Under the Act, every pesticide requires a unique identifier – the product's Registration or PCP number. That number must also appear on the product's label.

The pesticide label packaged with the product is the authoritative source of information on use of the product and will contain more detailed information than is included in this Guide. Some products have a number of trade names for the same active ingredient. However, each product will have its own Registration (PCP) number and these appear next to the registrants' names. Users who are seeking more detailed information than is provided in this guide, prior to purchase, can use the Registration (PCP) number to access a sample product label online through the Pest Management Regulatory Agency's (PMRA) website or they can contact the PMRA Hotline by phone at 1-800-267-6315.

Use this link – <http://www.hc-sc.gc.ca/cps-spc/pest/index-eng.php> – and select the link that reads 'Search Pesticide Labels' to access Electronic Label Search Tool. The PMRA Product Information database can be searched by a product's trade name, active ingredient, company name or Registration (PCP) number. Since several products can contain the same active ingredient and there are often several versions of the same or similar labels on this database, using the PCP number is the most direct route to finding the

label that links to the product page in this Guide. *There may be some differences between a label found on the package and the sample labels found on the PMRA-Label Search web site so always refer to the packaged product label when applying the product.*

Once the product is located, you may click on its number to view an Adobe Acrobat (PDF) document containing the label and any supplemental registrations. Some of these documents run to many pages but you can use the 'Find' capabilities of the Acrobat Reader plug-in for your browser to jump to specific areas of interest. If you do not have Adobe Acrobat Reader installed on your computer you can download a free version from www.adobe.com

Safe Use of Herbicides, Fungicides and Insecticides

Herbicides, fungicides and insecticides are classified according to the use hazard and risk involved. The categories of hazard are:

- toxicity
- flammability
- explosive potential
- corrosivity
- other

The degree of risk is represented by symbols taken from common traffic sign shapes represented by the stop, caution and yield signs. The signal word for each of the signs is danger (high risk), warning (moderate risk) and caution (low risk). Where the risk is minimal, no designation is required. The label on the container will carry the appropriate signs for the protection of the user. Degree of risk symbols for herbicides, insecticides and fungicides used in field and forage crops are included in the product directory. The symbols are illustrated in Figure 1.




Figure 1. Degree of Risk and Hazard Symbols



POISON CORROSIVE FLAMMABLE EXPLOSIVE OTHER

LD₅₀ values are used to rate the toxicity of pesticides. The LD₅₀ refers to the dose of pesticide (in mg per kg of the test animal's body weight) that is lethal to 50 percent of the group of test animals. For example, if a pesticide has an LD₅₀ value of 10 mg/kg, and the test animals each weigh 1 kg, then 50 percent of the animals would die if they each ate 10 mg of the pesticide.

Table 2. Oral LD₅₀ Values as they relate to the Risk/Hazard Symbols

	
DANGER POISON LD ₅₀ less than 500 mg/kg indicates high toxicity	WARNING POISON LD ₅₀ 500-1,000 mg/kg indicates moderate toxicity
	
CAUTION POISON LD ₅₀ 1,000-2,500 mg/kg indicates low toxicity	SYMBOL ABSENT LD ₅₀ greater than 2,500 mg/kg indicates very low toxicity

Different types of protective equipment are required for pesticides that differ in toxicity. Special equipment requirements are described on the product label, but in general the following precautions must be taken when using pesticides of different hazard ratings.

Danger Poison - requires goggles, respirator, gloves and skin protection, avoid fumes and spray mist.

Warning Poison - requires goggles, gloves and skin protection, avoid fumes and spray mist.

Caution Poison - requires gloves and skin protection, avoid fumes and spray mist.

The absence of a hazard symbol on a pesticide label indicates low toxicity to mammals. Nevertheless, protective clothing should be worn when using pesticides that do not have a hazard symbol.

Protecting Yourself from Exposure to Herbicides, Fungicides and Insecticides

The use of protective equipment and sound safety procedures will help minimize your exposure to herbicides, fungicides and insecticides. Follow the 10 rules for safe application listed below, and wear the safety equipment recommended.

10 Rules for Safe Application

1. Never smoke or eat while applying pesticides.
2. Avoid inhaling sprays or dusts. Wear protective clothing and a respirator.
3. Sprayer lines carrying chemicals should not enter the operator's cab.
4. Have soap, water and a towel available. Should concentrated product spill on skin, hands, face or eyes, wash immediately.
5. Wash hands and face when leaving the treated area, before break periods, lunch or urination.

6. Bathe or shower and change into clean clothing after working with pesticides. Wash clothing each day before re-use.
7. Call a physician or get the patient to a hospital immediately if symptoms of illness occur during or shortly after pesticide application. Be sure to take along the product label or container.
8. Store pesticides out of reach of children and where there is no chance of contact with human food or livestock feeds. Do not store herbicides with insecticides and avoid cross-contamination. Storage areas should be locked.
9. Keep chemicals in their original containers, never in unmarked containers or bottles used for food or drink.
10. Follow proper container disposal methods. All containers should be triple rinsed or pressure rinsed, punctured to render the container non-reusable, and delivered to designated disposal sites.

Protective Clothing

Wear protective equipment as described in the chart to reduce exposure.

EQUIPMENT	PROTECTION	HOW TO WEAR IT
Coveralls	<p>There are two types of coveralls: disposable and reusable. Disposable coveralls are lightweight and comfortable on warm days. They can be worn for mixing and applying pesticides, then discarded at the day's end. If they become contaminated, they should be discarded at once.</p> <p>The second type of coverall is made of washable fabric and may be reused many times. These fabric coveralls are adequate for use with all but the most highly toxic and concentrated pesticides.</p>	<p>Button (or zip) right up to the neck. Loose coveralls around the neck will suck and blow pesticide in and out of the interior of the coveralls as you bend and move.</p> <p>Wear coveralls over a long-sleeved shirt and pants.</p>
Aprons	<p>When pouring or otherwise handling concentrated pesticides, it makes good sense to wear protection in the form of an apron. The apron protects the front of your body from spills or splashes of the concentrate. The apron should be made of rubber or synthetic liquid-proof material that will resist the solvents used in formulating the pesticide.</p>	<p>Make sure the apron covers your body from your chest to your boots.</p>
Gloves	<p>Protect your hands by wearing chemical-resistant gloves. Neoprene gloves provide the best protection. Natural rubber gloves may be used when handling organo-phosphorus or carbamate pesticides. Be sure that they are designed for use with solvents and pesticides. Never use lined gloves, gloves with wristbands or leather gloves.</p>	<p>Put gloves on and roll up the first inch or two of the cuff. That way when you lift your hands, any liquid on the gloves won't drip down your arms.</p>
Hats	<p>Use a chemical-resistant hat, preferably made of washable plastic. The hat may be a hard hat or made of flexible plastic. In either case, it should have a plastic sweatband. Wash and dry entire hat after each use and before storing. Ordinary baseball caps with cloth sweatbands are dangerous as they absorb the pesticide and recontaminate the forehead each time you wear them. Even small amounts of moderately or slightly toxic pesticides may cause severe skin irritation or other illness if exposure continues for several days.</p>	
Boots	<p>Wear chemical-resistant, unlined boots. These boots are available in a variety of styles and materials. Neoprene boots are the best. Knee-length boots offer greater protection because they extend above the lower end of the apron. Avoid leather or fabric boots and shoes because these will absorb pesticides and cannot be cleaned effectively.</p>	<p>Wear your pant legs outside the top of your boots. This will prevent spills and splashes from running into the boot and onto your leg.</p>

Protecting Your Eyes, Face and Lungs

Wear the following equipment to protect your facial area from exposure:

EQUIPMENT	PROTECTION	HOW TO WEAR IT
Goggles	Chemical-resistant goggles keep your eyes safe from both splashing and, if using dry formulations, dusts or granules. Don't use goggles with cloth or elastic headbands as these will absorb pesticides.	Wear goggles snugly on your face so that the sides of your head are protected from splashes. If you wear glasses, make sure you purchase goggles that fit snugly over them. Never wear contact lenses when working around pesticides.
Respirators	Only NIOSH-approved respirators should be used. Do not exchange parts of different respirators. (For example, do not use a cartridge produced by Company "A" with a respirator produced by Company "B" as the combination may not provide adequate protection to the user). Dust masks are ineffective in protecting against herbicide vapours. Similarly, the filters on tractor cabs are intended to remove dust and are not designed to protect against herbicide vapours or mists. Chemical cartridge respirators are recommended for outdoor use when mixing and applying herbicides.	When carrying out operations, change filters each day. The cartridge should be replaced when chemical odour becomes apparent or when breathing becomes difficult. New cartridges should always be installed at the beginning of the spray season. Prior to commencing work, check the face seal while the respirator is on the wearer's face. Regardless of design, respirators cannot be worn securely by people wearing beards, moustaches or sideburns.
Face Shields	Goggles offer some protection, but frequently full-face protection is advised or required according to the pesticide label. It is especially important to protect your eyes and face when pouring or mixing liquid concentrates. Effective face shields are made of clear plastic.	Since the shield attaches to the hard hat, you can raise or lower it as needed.

Avoiding Spray Drift

To minimize the risk of drift, follow these guidelines:

- Do not spray in winds above 15 km/h (9 mph).
- Do not spray under dead calm conditions in early morning, night, or late evening. These are often associated with temperature inversions and the combination of these factors can result in long-distance spray drift (2 km or more). Fog or dust that seems to hang in the air is a good indicator of an inversion.
- Avoid nozzle pressures above 45 psi (310 kPa) for conventional flat fan tips.
- Use a minimum of 45 L/acre water for all pesticides unless otherwise specified for the product.
- Take note of buffer zones identified in the "Restrictions" section of this guide. Do not spray when the wind is blowing towards a nearby sensitive crop, shelterbelt, garden, or water body.
- Use amine formulations of 2,4-D or MCPA where possible. Use special care when applying volatile herbicides (most herbicides in Group 3 and Group 4, particularly ester formulations). Avoid spraying these products on or immediately before hot days.
- Ensure that air flow from air assisted sprayers is properly set to minimize airblast rebound and drift for different crop canopies.
- Operate nozzles at their minimum recommended height. For 80° tips, this is 18" (45 cm), and for 110° tips, this is 12" (35 cm). Orienting nozzles forward allows further height reductions.
- Special nozzles are now available that create coarse, low-drift sprays. Pre-orifice, Turbo-TeeJet, or venturi-type nozzles are available from a number of manufacturers, and these reduce drift by 50 to 95 percent. (Refer to the section entitled **Herbicide Efficacy with Low-Drift Nozzles**)
- Consider equipping your sprayer with protective shrouds. A number of different designs are available that can reduce drift between 35 and 75 percent.
- Reduce travel speeds. rapid air movement over nozzle tips increases the risk of fine droplets prone to drift and turbulence from the sprayer itself can increase the uncertainty of spray deposition.

For more information on reducing drift refer the Factsheet entitled "Spray Drift – Causes and Solutions" available at the Saskatchewan Ministry of Agriculture Website: www.agriculture.gov.sk.ca

Herbicide Efficacy with Low-drift Nozzles

A number of low-drift nozzles are now available from different suppliers. Well established nozzles, such as the Turbo Teejet, reduce drift by about 50 percent and provide equivalent efficacy to a standard flat fan nozzle. Newer nozzles ("venturi" types) are best known for their dramatic ability to reduce drift (50 to 95 percent). Research suggest that these nozzles perform well at conventional carrier volumes, travel speeds, and product rates. Some aspects require special attention:

Pressure: Some venturi-type nozzles require higher pressures to operate properly. Below 40 psi (275 kPa), patterns for these designs may deteriorate rapidly resulting in poor overlaps and erratic control. Design improvements have resulted in venturi nozzles that require less pressure to operate effectively. When using automatic rate controllers, make sure your pressures match the recommended pressure ranges for good nozzle performance.

Water Volume: Droplet size becomes more important at lower water volumes. Little is known about low-drift nozzle performance at or below 5 gal/acre (23 L/acre). Since low-drift nozzles generate fewer droplets than conventional nozzles, ensure that water volumes are high enough for coverage when using coarse sprays.

Weed Type: Difficult-to-wet weeds, such as wild oats, green foxtail, lamb's-quarters, and cleavers, typically require finer sprays for effective coverage. When using venturi nozzles on these weeds, make sure your pressure is high enough to achieve good coverage. Larger weeds and reduced product rates typically make chemical control more difficult, and these conditions may also reveal some performance differences between nozzles.

Herbicide Type: Herbicides that belong to herbicide Groups 2, 4, and 9 perform well with venturi nozzles, even at normal pressures (40 psi). Application of herbicides in Groups 1, 6, 8, 10 and 14 may require higher pressures with venturi nozzles to maintain good performance, especially under challenging conditions. Wild oat control may be reduced with the coarsest sprays, even when applied at high pressure.

Check with your chemical representative to see if the manufacturer supports the use of low-drift nozzles with their products.

More information is available in the factsheet "Pesticide Application and Choosing the Right Nozzles," available from your local extension office or at the Saskatchewan Ministry of Agriculture Website: www.agriculture.gov.sk.ca

Handling a Drift Complaint

When spray drift occurs, it is important to take the right steps to resolve the complaint. If you suspect that your crop or property has been damaged because of spray drift, use the following guidelines for resolving the situation.

1. Are you sure that the symptoms or damage you see has been caused by spray drift? Contact your local agricultural office to help determine if the damage is the result of spray drift.
2. Contact the suspected applicator as soon as possible. View the damage with the suspected applicator and determine if that person did, in fact, cause the damage.
3. If the damage was caused by the applicator, determine the extent of the damage and the level of compensation (if any) with the applicator.
4. If the situation cannot be resolved quickly because of disagreements on the extent of damage, cause of the damage, or level of compensation, contact your local agricultural office to discuss options on how to proceed. Documentation will be required, particularly if insurance companies are involved.
5. The involvement of a private consultant is recommended if documentation is required. Required documentation often includes samples of the damaged plants, photographs, and yield comparisons to determine losses. Your agricultural office can provide you with a list of private consultants in your area.
6. The best approach is to start an open and honest line of communication with the suspected applicator. The majority of drift complaints are resolved quickly and efficiently by communicating with the applicator, without the involvement of outside parties.

Mixing Pesticides

The ability to control a broad range of weeds or other pests in one pass is the advantage that a mix of two or more products allows. If tank mixing is not done in the correct order, the result could be a tank-load of material that may not control the target pests, cause injury to the crop, plug nozzles, or leave an undesirable residue in the tank that will require extensive cleaning. Mistakes like these are costly, could put the user at unnecessary risk of exposure to the products, or create an environmental disposal problem.

To avoid mixing that may result in incompatibilities, always consult the label of the products that are being used to learn the correct order. Remember to add all like components at the same stage of mixing. The list below is a general rule-of-thumb for mixing pesticides:

1. Fill the spray tank with 1/4 to 3/4 the amount of water required for the application and turn on the sprayer agitation. Check the products that are being used for the correct amount to add. Once agitation has begun, maintain until the tank is emptied.
2. Add any fertilizer or pH adjuster additives to the tank.
3. Add any wettable powders, water dispersible granules (DF, DG, or WDG), or flowable liquid suspensions to the tank. Add dry products slowly to prevent clogged return lines. Allow sprayer to agitate for a few minutes, allowing the product to become completely suspended in the tank, before adding the next component.

4. Shake any containers of liquid pesticide thoroughly before adding to ensure they are well mixed.
 5. Add any pesticides that are solutions (SN) (i.e. amines and salts)
 6. Add emulsifiable concentrates (EC, SC) (i.e. esters)
 7. Add any surfactants or other adjuvants.
- Remember to always consult the label for compatible mixes and recommended mixing order.

Many pesticides will break down if left in the tank for an extended period. Try not to mix any more than you can spray at one time. If you need to stop spraying for a short time, leave the sprayer agitation running to keep products from settling or separating in the tank.

Container Disposal

Proper disposal of used containers and unused pesticides is important to protect the environment and prevent contamination of soil and water resources. Rinse all containers prior to disposal to reduce environmental contamination caused by open dumping of unwanted containers. Only mix as much pesticide solution as is needed to treat the desired area.

Triple Rinsing

Triple rinsing renders used pesticide containers (metal, plastic, glass) more than 99.9 percent free of residues, in most cases.

Here are the steps that should be followed:

1. Empty contents of the container into the spray tank and drain in a vertical position for 30 seconds.
2. Add a measured amount of rinse water or other diluent until container is about one-fifth full.
3. Rinse the container thoroughly and pour the rinsate into the spray tank.
4. Repeat the procedure twice (it should take only about 5 minutes in total).
5. Puncture or break triple rinsed containers to render them non-reusable. Paper bags should be rinsed once prior to disposal.

Pressure Rinsing

Pressure rinsers can be used to rinse any size of empty pesticide container that can be lifted into position over the spray tank. A 30 second rinse with a pressure rinser is convenient and just as effective as triple rinsing. Pressure rinsers are constructed to be thrust into the bottom of a metal can or plastic jug. Holes, situated laterally in the rinser tip, direct water from a pressurized source against the inner sides of the container and effectively wash the residual pesticide into the spray tank. Some farmers have found it convenient to attach a rinser to the pump on their large water storage tank to minimize container handling. Pressure rinsers have the added advantage of rendering containers useless by automatically puncturing them.

Disposal of Containers

Properly rinsed containers should be delivered to a designated pesticide container disposal site. Contact your ag rep, municipal office or weed supervisor for the locations of pesticide container disposal sites in your municipality.

Sprayer Tank Cleaning

When pesticide application is completed each day it is important to empty and clean the sprayer thoroughly to prevent the breakdown of certain pesticides, prevent adhesion of the pesticide to the sprayer, and to maintain the sprayer parts in good condition. Certain pesticides break down very quickly when left in solution, and several pesticide solutions can be corrosive to sprayer parts. Tank cleaning is especially important when changing from one crop to another or from one pesticide to another. Each year several reports are logged of herbicide damage caused by carryover of product residue in the tank. To avoid the risk of contamination, sprayers should be cleaned as soon as possible after application is completed.

Do not clean sprayers where rinsate can run off into ditches or other water bodies, near sensitive plants or shelterbelts, or where other people or animals are likely to walk, to avoid unnecessary exposure to people, animals and the environment.

There are three basic types of rinse solution for cleaning sprayer tanks. Their recipes and basic procedures are outlined below:

The Ammonia Rinse – Fill spray tank and add 1 L of household ammonia (3%) for every 100 L of clean water needed for the rinse and begin agitation. Allow solution to flush through the booms until the boom is completely filled with ammonia solution and top up the tank with water. Circulate the ammonia solution through the tank and pump system for 15 minutes. Flush hoses and booms with ammonia rinse solution again (minimum 5 minutes) before emptying. Remove nozzles and screens and scrub with 0.1 L household ammonia per 10 L clean water and an old toothbrush. Perform clean water rinse to remove ammonia solution prior to next spray load. Some herbicides recommend leaving the ammonia rinse in the tank over night to improve cleaning potential.

The Fresh Water Rinse – The spray tank cleaning should begin and end with a fresh water rinse to remove the majority of potential contaminants prior to the cleansing process or prior to the next round of spraying. Drain the tank of its previous contents and fill the tank with clean water. Open nozzle valves and pump clean water through the booms and hoses. Top up the tank with more clean water and circulate/agitate for at least 10 minutes and empty the tank of waste water. If this is the first rinse after spraying, a high pressure hose could be used to clean residue from all surfaces in the tank. Do not enter the tank during the cleaning process.

The Detergent Rinse – After rinsing with clean water, fill spray tank and add a heavy-duty detergent at 0.25 L per 100 L of water (some suggest a non-ionic surfactant such as Agral 90 or Agsurf at 0.6 L per 100 L of water). Circulate the mixture for a minimum of 5 minutes and spray out through sprayer nozzles. Nozzles and screens are removed and cleaned individually with the same detergent solution in a small container. Soaking in this solution for several hours also helps to loosen any deposits.

The above solutions are just components of the overall sprayer cleaning process. Typical rinse instructions will repeat a combination of one or two or all of these basic rinses. Below we will give some generic rinse instructions utilizing the basic rinses as components of the larger cleaning procedure. Never enter the tank during the cleaning process as some cleansers may release dangerous gases.

Method A –

Drain contents of tank – 1 to 2X Water Rinse – 2X Ammonia Rinse – 2X Water Rinse (one just prior to the next spraying event)

Products: 2,4-D, Aatrex, Accent, Aim, Authority, Barricade, Battalion, Blazer, Broadband, CleanStart, dicamba, Dichlorprop/2,4-D, DyVel, Escort, Everest, GBX, Express Pro, florasulam + glyphosate, florasulam + MCPA, flucarbazone, flumioxazin, fluroxypyr + 2,4-D, Frontier, Frontline 2,4-D, glyphosate/dicamba, Impact, Inferno Duo, Infinity, MCPA, metsulfuron, Muster, Option, Pinnacle, Prism, Reclaim, Restore, simazine, Spectrum, Stellar, thifensulfuron/tribenuron Tordon 22K, Traxos, tribenuron, Triton C & K, Tundra, Ultim, Varro, Velocity.

Method B –

Drain Contents of tank – 2X Water Rinse – 2X Detergent Rinse – 2X Water Rinse

Products: Achieve Liquid Gold*, Avadex, Axial, Basagran, Bromoxynil/MCPA, Clethodim, diquat, Distinct, DyVel DSp, Enforcer, fenoxaprop, FlexStar, Focus, GoodHarvest, Gramoxone*, Heat, Liberty, Lontrel, metribuzin, Momentum, Overdrive, Poast Ultra, PrimExtra Magnum, Pulsar, quizalofop, Reflex, Reward*, tralkoxydim, trifluralin products, Trophy, Velpar, Viper ADV.

*Manufacturers of these products recommend adding a non-ionic surfactant such as Agral 90 or Agsurf II at 0.6 L per 100 L of water.

Method C –

Drain Contents of tank – Several repetitions of the Water Rinse with nozzles and screens removed and checked for debris.

Products: Adrenalin, Altitude, Amitrol 240, Ares, Clodinafop, Dual II Magnum, Eclipse, glyphosate products, imazamethabenz, imazethapyr, Odyssey, Prestige, Solo.

The above directions are general processes based on the similarities of tank cleaning recommendations between products in each of the herbicide groupings. Always follow the specific instructions on the product label.

Several products in the guide do not have label instructions regarding tank cleaning. In the case of products that have no cleaning recommendations on the label, there are some basic principals that can be applied. Products that are water based formulations can usually be cleaned from spray tanks using Method C above. Products that are formulated as an EC, SC or F (flowable) or use a petroleum based adjuvant should at least use Method B. The detergent breaks down the oil that may be sticking to the side of the tank. Products in Group 2 (most will already have a recommendation) with the exception of the 'IMI' products from BASF, will require the use of Method A. The ammonia in Method A either increases the solubility of the product allowing it to be easily removed from the tank surfaces or speeds the breakdown of these products in water. If the product that is to be cleaned out of the tank is a combination of these elements, use a combination of Methods to clean the tank. In these cases, use a good commercial tank-cleaning product from a recognized source, with both ammonia and detergent as components.

Group 2 compounds are highly active on sensitive plants so even a small amount remaining in the sprayer can present a risk of injury. They can also occasionally be trapped on the tank walls and plumbing by petroleum based formulations or adjuvants when tank mixed with other products, resulting in tank residues that may be tougher to remove. A way to reduce the chance of this occurring is to add detergent at 0.25 L per 100 L to the Ammonia Rinse portion to assist with the breakdown of the petroleum coating so that the ammonia may rid the tank of Group 2 product.

It is very important to clean sprayers immediately after every use. With a more diverse rotation, the likelihood of damage from lack of care increases dramatically.

How to Identify Crop and Weed Leaf Stages

Recognition of plant growth stages is essential for effective weed and disease control. Many herbicides and fungicides are safe on a crop only when applied at a specific growth stage. Similarly, weeds are controlled only when they are at certain growth stages.

For most post-emergent products, growth stages are described by the number of leaves. The following is a description of how to count leaves for staging.

Cereals and Annual Grass Weeds

Manufacturers generally use two different systems of staging for grasses. The minimum stages of application are similar, while the later stages may differ.

Some manufacturers use "Total Leaf Count" stages based on the number of leaves on the entire plant, including tillers or secondary shoots. Most recommendations however, are based on the number of main stem leaves and tillers. Tillers or stools are the secondary shoots or stems of a grass plant. Similar to the branches of a broadleaf plant, tillers will emerge from the axils between the leaf and main shoot.

Tillers usually begin to appear at the 3 or 4 leaf stage. When staging a plant in this manner, be sure to identify the tillers first, then count only leaves that originate from the main shoot.




Figure 2. Leaf Stages of Cereals and Annual Grass Weeds






Figure 3. Leaf Stages of Broadleaf Weeds and Crops



Broadleaf Weeds


Cotyledons - These are the seed leaves that usually emerge above ground. On some plants, such as fababeans, lentils and peas, they stay below the soil surface. Cotyledons are not true leaves and are not counted when determining leaf number. They are a different shape than the true leaves and may dry up and disappear at an early stage.

Alternate leaves - Some plants have one leaf at each node on the stem. The next leaf emerges at the next higher node and extends away from the stem in the opposite direction. These plants (lamb's quarters and wild mustard are good examples) are said to have alternate leaves. To determine the leaf stage, simply count the number of leaves present (Figure 3).

Opposite leaves - Plants with two leaves at each node, one on each side of the stem, are said to have opposite leaves. The next pair of leaves on the next node are rotated about 45° so that they are not directly over the previous pair. Plants with opposite leaves have even-leaf numbers only. When counting, the leaf number progresses from cotyledons to 2 leaf, 4 leaf, etc. These plants generally appear shorter than plants with alternate leaves at a similar leaf stage. **Be sure to count each pair as two leaves.** Hemp nettle is a weed that has opposite leaves (Figure 3).

Whorled leaves - More complex plants like cleavers may have whorled leaves. These plants have three or more leaves at each node on the stem. The leaf number in each whorl may vary, so be sure to count each individual leaf unless the Guide or label recommendation refers to the number of leaf whorls (Figure 3).

Figure 4. Leaf Stages of Certain Special Crops and Forages



PESTICIDE FORMULATIONS AND COMPANIES

Herbicides

Product	Common Name	Formulation	Company
2,4-D	2,4-D	600, 700 g/L SN, EC	Various
A/Atrex Liquid	atrazine	480 g/L SC	Syngenta Canada
Accent	nicosulfuron	75% DG	E. I. duPont Canada
Accurate	metsulfuron methyl	60% DG	Cheminova Canada
Achieve (Liquid Achieve)	tralkoxydim	400 g/L SC	Dow AgroSciences
Achieve Liquid Gold	tralkoxydim + bromoxynil : MCPA E	400 g/L SC + 225 : 225 g/L EC	Nufarm Agriculture
Adrenalin SC	imazamox : 2,4-D LV ester	20 : 560 g/L SC	BASF Canada
Aim	carfentrazone	240 g/L EC	FMC of Canada
Ally Toss-N-Go	metsulfuron methyl	60% DG	E. I. duPont Canada
Altitude FX	imazamox + fluroxypyr + MCPA 600 ester	120 g/L SN + 180 g/L + 600 g/L EC	BASF Canada
Altitude FX 2	imazamox + fluroxypyr	120 g/L + 180 g/L SN	BASF Canada
Amitrol 240	amitrole	231 g/L SN	Nufarm Agriculture
Approve	bromoxynil : : 2,4-D LV E	225 : 225 g/L EC	Nufarm Agriculture
Ares	imazamox : imazapyr	33 : 15% SN	BASF Canada
Arrow	clethodim	240 g/L EC	MANA Canada
Assert 300 SC	imazamethabenz	300 g/L SC	Nufarm Agriculture
Assert FL	imazamethabenz + florasulam + MCPA ester	300 g/L SC + 50 g/L SC + 500 g/L EC	Nufarm Agriculture
Assure II	quizalofop-p-ethyl	96 g/L EC	E. I. duPont Canada
Attain XC	fluroxypyr + 2,4-D E	333 g/L + 660 g/L	Dow AgroSciences
Aurora	clodinafop-propargyl	240 g/L EC	Farmers of North America
Authority	sulfentrazone	480 g/L SC	FMC of Canada
Authority Charge	sulfentrazone + carfentrazone	480 g/L SC + 240 g/L EC	FMC of Canada
Avadex MicroActiv	triallate	10% G, 480 g/L EC	Gowan Canada
Extra Strength BW			
Avenge 200-C	difenzoquat	200 g/L SN	Syngenta for AMVAC
Avert	imazamethabenz	300 g/L SC	Loveland (formerly Viterro)
Axial	pinoxaden	100 g/L EC or 50 g/L EC (BIA)	Syngenta Canada
Axial iPak	pinoxaden + pyrasulfotole : bromoxynil	100 g/L EC or 50 g/L EC (BIA) + 37.5 : 210 g/L EC	Syngenta Canada
Axial Xtreme	pinoxaden : fluroxypyr	50 : 87.5 g/L EC	MANA Canada
Badge	bromoxynil : MCPA E	225 : 225 g/L EC	BASF Canada
Barvel II	dicamba (diglycolamine salt)	480 g/L SN	Engage Agro (BASF)
Barvel VM	dicamba (diglycolamine salt)	480 g/L SN	E. I. duPont Canada
Barriade II	thifensulfuron : tribenuron + fluroxypyr	50% : 50% + 333 g/L EC	BASF Canada
Basagran/Basagran Forté	bentazon	480 g/L SN	E. I. duPont Canada
Battalion	rimsulfuron + metalachlor + dicamba	25% DF + 915 g/L EC + 480 g/L SN	Nufarm Agriculture
Blackhawk	Carfentrazone + 2,4-D Ester	240 g/L EC + 660 g/L EC	MANA Canada
Bengal WB	fenoxaprop-p-ethyl	120 g/L EC	MANA Canada
Bison	tralkoxydim	400 g/L SC	United Phosphorus Inc.
Blazer	acifluorfen	240 g/L SN	UAP
Bonanza	trifluralin	480 g/L EC, 10% G	Syngenta Canada
Broadband	pinoxaden : florasulam	92.7 : 7.7 g/L EC	Loveland (formerly Viterro)
BroadSide	thifensulfuron methyl : tribenuron methyl + MPCA	33.35% : 16.65% WSG + 500 or 600 g/L EC	
Bromotril	bromoxynil	240 g/L EC	MANA Canada
Brotex	bromoxynil	240 g/L EC	IPCO
Buctril M	bromoxynil : MCPA E	280 : 280 g/L EC	Bayer CropScience
Bullchips	clodinafop-propargyl	240 g/L EC	Cheminova Canada
Caliber	2,4-DB	625 g/L EC	UAP
Casoron	dichlobenil	4% G	UAP
Centurion	clethodim	240 g/L EC	Bayer CropScience
Chateau	flumioxazin	51% DG	Valent Canada
Cheminova Glyphosate	glyphosate IPA salt	356 g/L SN	Cheminova Canada
ClearOut 41 Plus	glyphosate IPA salt	360 g/L SN	Farmers of North America
CleanStart	glyphosate IPA + carfentrazone	356 g/L SN + 240 g/L EC	Nufarm Agriculture
Cloverox Plus	MCPB : MCPA	375 : 25 g/L SN	IPCO
Cobutox	2,4-DB	625 g/L EC	IPCO
Compitox	mecoprop-p	150 g/L SN	Nufarm Agriculture
Cordon	fenoxaprop-p-ethyl	120 g/L EC	Nufarm Agriculture
Cougar	fenoxaprop-p-ethyl	120 g/L EC	Cheminova Canada
Credit 45	glyphosate IPA salt	450 g/L SN	Nufarm Agriculture
Crush'R Plus	glyphosate IPA salt	360 g/L SN	AgWest/Agri-Star
Curtail M	clopyralid : MCPA E	50 : 280 g/L EC	Nufarm Agriculture
Deploy	thifensulfuron methyl : tribenuron methyl	50% : 25% DG	Aryla LifeScience
Desica	diquat	240 g/L SN	Syngenta Canada

Product	Common Name	Formulation	Company
Dichlorprop-D	dichlorprop : 2,4-D E	300 : 282 g/L EC	IPCO
Dichlorprop-DX	dichlorprop-P : 2,4-D E	210 : 400 g/L EC	IPCO
Distinct	dicamba : diflufenzopyr	50 : 20% DG	BASF Canada
Dual II Magnum	metolachlor	915 g/L EC	Syngenta Canada
DyVel	dicamba : MCPA K	84 : 336 g/L SN	BASF Canada
DyVel DSp	dicamba : mecoprop-p : 2,4-D A	110 : 80 : 295 g/L SN	BASF Canada
Eclipse III	glyphosate DMA + clopyralid	480 g/L + 360 g/L SN	Dow AgroSciences
Edge Granular	ethalfluralin	5% G	Dow AgroSciences
Embutox 625	2,4-DB	625 g/L EC	Nufarm Agriculture
Enforcer D	fluroxypyr : bromoxynil: 2,4-D	80: 190: 240 g/L EC	Nufarm Agriculture
Enforcer M	fluroxypyr : bromoxynil: MCPA	80: 200: 200 g/L EC	Nufarm Agriculture
Eptam-8E	EPIC	800 g/L EC	Gowan Company
Equinox	tepraloxym	200 g/L EC	BASF Canada
Escort	metsulfuron methyl	60% DG	E.I. duPont Canada
Estaprop XT	dichlorprop-P : 2,4-D E	210 : 400 g/L EC	Nufarm Agriculture
Everest 2.0	flucarbazone	397 g/L SC	Arysta LifeScience
Everest GRX	flucarbazone + fluroxypyr	66% DG + 180 g/L EC	Arysta LifeScience
Express Pro	tribenuron methyl : metsulfuron	42.9% : 8.6% SG	E.I. duPont Canada
Express SG	tribenuron methyl	50% SG	E. I. duPont Canada
FirstStep Complete	tribenuron methyl + glyphosate	75% DG + 540 g/L SN	Loveland (formerly Viterra)
FlexStar GT	Fomesafen : glyphosate	67 : 271 g/L SN	Syngenta Crop Protection
Flurox 2,4	fluroxypyr + 2,4-D E	180 g/L EC + 660 g/L EC	Nufarm Agriculture
Focus	Pyroxasulfone + carfentrazone	85% WDG + 240 g/L EC	FMC Canada
Foothills NG	clodinafop-propargyl	60 g/L EC	Loveland (formerly Viterra)
Fortress	triallate : trifluralin	10% : 4% G	Gowan Canada
Frontier Max	dimethanamid - P	720 g/L EC	BASF Canada
Frontline XL	florasulam : MCPA ester	4 : 280 g/L EC	Dow AgroSciences
Frontline 2,4-D XC	florasulam + 2,4-D E	50 g/L SC + 660 g/L EC	Dow AgroSciences
Gladiator	imazethapyr	240 g/L SN	Univar Canada
Glufos	glyphosate IPA salt	360 g/L SN	Chemnova Canada
Glykamba	glyphosate IPA + dicamba	194 g/L + 46 g/L SN	Nufarm Agriculture
Good Harvest (MPower)	glufosinate ammonium	150 g/L SN	Farmers of North America
Gramoxone	paraquat	200 g/L SN	Syngenta Canada
Grazon	picloram : 2,4-D E	65 : 240 g/L SN	Dow AgroSciences
Harmony K	clodinafop propargyl + thifensulfuron methyl + tribenuron methyl : dicamba	128 g/L EC + 7.7% : 3.9% : 54% DG	E. I. duPont Canada
Harmony Max	clodinafop propargyl + thifensulfuron methyl : tribenuron methyl + fluroxypyr	128 g/L EC + 33.35% : 16.65 SG + 180 g/L EC	E. I. duPont Canada
Harmony SG	clodinafop propargyl + thifensulfuron methyl : tribenuron methyl	128 g/L EC + 33.35% : 16.65 SG	E. I. duPont Canada
Harvey Power	dicamba (dimethylamine salt)	480 g/L SN	Adjuvants Plus (Gharda)
Heat	safufenacil	70% WSG	BASF Canada
Horizon NG	clodinafop-propargyl	60 g/L EC	Syngenta Canada
Impact	topremazone	336 g/L SC	UAP for AMVAC
Inferno	tribenuron	75 % WDG	Arysta LifeScience
Inferno Duo	Hucarbazone : tribenuron	45% : 25% WDG	Arysta LifeScience
Infinity	pyrasulfotole : bromoxynil	37.5 : 210 g/L EC	Bayer CropScience
Kerb SC	propyzamide	400 g/L SC	Dow AgroSciences
Kerb 50WSP	propyzamide	50% WP	Dow AgroSciences
Ko-Act	Tribenuron + 2,4-D	75% WDG + 660 g/L EC	Nufarm Agriculture
Koril 235	bromoxynil	235 g/L EC	Nufarm Agriculture
Knockout Extra	glyphosate IPA salt	360 g/L SN	Great Northern Growers
Ladder	clodinafop-propargyl	240 g/L EC	MANA Canada Inc.
Laji Plus	glyphosate IPA salt	360 g/L SN	Ray Glenn Commodities
Leader	bromoxynil : 2,4-D E	225 : 225 g/L EC	IPCO
Legend	clodinafop-propargyl	240 g/L EC	IPCO
Liberty	glufosinate ammonium	150 g/L SN	Bayer CropScience
Liberty 200 SN	glufosinate ammonium	200 g/L SN	Bayer CropScience
Linuron 400	linuron	400 g/L SC	UAP
Logic M	bromoxynil : MCPA E	225 : 225 g/L EC	IPCO
Lontrel 360	clopyralid	360 g/L SN	Dow AgroSciences
Lorox L	linuron	480 g/L SC	Tessenderlo Kerley Inc.
Marengo	tralkoxydim	400 g/L SC	Dow AgroSciences
Matrix	glyphosate DMA salt	480 g/L SN	IPCO
Maverick III	glyphosate DMA salt	480 g/L SN	Dow AgroSciences
MCPA	MCPA	300,400,500,564 g/L SN, EC	Various
Mecoprop-P	mecoprop-P	150 g/L SN	UAP
Mextrol 450	bromoxynil : MCPA E	225 : 225 g/L EC	Nufarm Agriculture
Momentum	clopyralid : fluroxypyr	90 : 90 g/L EC	Loveland (formerly Viterra)
MPower Glyphosate	glyphosate IPA salt	360 g/L SN	Farmers of North America
MPower HellCat	fenoxaprop-p-ethyl	120 g/L EC	Farmers of North America
MPower Kamikaze	imazethapyr	240 g/L SN	Farmers of North America
MPower R	thifensulfuron methyl: tribenuron methyl	50%:25% DG	Farmers of North America

Product	Common Name	Formulation	Company
MPower X	tribenuron methyl	75% DG	Farmers of North America
MultiStar	imazethapyr	240 g/L SN	Loveland (formerly Viterra)
Muster Toss-N-Go	ethametsulfuron-methyl	75% DG	E. I. duPont Canada
NextStep NG	clodinafop-propargyl	60 g/L EC	Arysta LifeScience
Nimble	thifensulfuron methyl :	50% : 25% DG	Cheminova Canada
Nuance	tribenuron methyl	75% DG	Cheminova Canada
Nufarm Clodinafop	tribenuron-methyl	240 g/L EC	Nufarm Agriculture
NuClo	clodinafop-propargyl	450 g/L SN	Nufarm Agriculture
OcTtain XL	glyphosate IPA salt	90 : 360 g/L EC	Dow AgroSciences
Odyssey	fluroxypyr : 2,4-D E	35% : 35% DG	BASF Canada
Odyssey DLX	imazamox : imazethapyr	35% : 35% DG +	BASF Canada
	imazamox : imazethapyr +	200 g/L EC	
	tepraloxym	160 : 130 : 310 g/L SN	UAP
Optica Trio	MCPA A : mecoprop-P : dichlorprop-P	35% DG	Bayer CropScience
Option 35 DF	foramsulfuron	22.5 g/L OD	Bayer CropScience
Option 2.25 OD	foramsulfuron	480 g/L SN	Adjuvants Plus (Gharda)
Oracle	dicamba (dimethylamine salt)	50% : 20% DG	Engage Agro (BASF)
Overdrive	dicamba : diflufenzopyr	280 g/L EC	Bayer CropScience
Pardner	bromoxynil	240 g/L SN	MANA Canada
Phantom	imazethapyr	50% WSG	E. I. duPont Canada
Pinnacle SG	thifensulfuron methyl	450 g/L EC	BASF Canada
Post Ultra	sethoxydim	480 g/L SN + 50 g/L SC	Dow AgroSciences
PrePass XC	glyphosate DMA + florasulam	333 g/L +	Dow AgroSciences
Prestige XC	fluroxypyr +	50 : 280 g/L EC	
	clopyralid : MCPA E	400 : 320 g/L SC	Syngenta Canada
Primextra II Magnum	metolachlor : atrazine	90% DG	True North
Princep Nine T	simazine	50 g/L SC	MANA Canada
Priority	florasulam	25% SG	E.I. DuPont Canada
Prism SG	rimsulfuron	87 : 113 g/L EC	Syngenta Canada
Pulsar	dicamba : fluroxypyr	90 g/L EC	Bayer CropScience
Puma Advance	fenoxaprop-p-ethyl	240 g/L SN	BASF Canada
Pursuit	imazethapyr	52.5% : 9.45% DG +	Dow AgroSciences
Reclaim	aminopyralid : metsulfuron-methyl +	564 g/L EC	
	2,4-D E	33.35% : 16.65% SG +	E. I. duPont Canada
Refine M	thifensulfuron methyl :	500 or 600 g/L EC	
	tribenuron methyl + MCPA E	33.35% : 16.65% SG	E. I. duPont Canada
Refine SG	thifensulfuron methyl :		
	tribenuron methyl		
Reflex	fomesafen	240 g/L SN	Syngenta Canada
Reglone Desiccant	diquat	240 g/L SN	Syngenta Canada
Reglone Ion	diquat	200 g/L SN	Syngenta Crop Protection
Restore	aminopyralid + 2,4-D A	240 g/L + 564 g/L SN	Dow AgroSciences
Restore II	aminopyralid : 2,4-D A	40 : 400 g/L SN	Dow AgroSciences
Retain SG	thifensulfuron methyl : tribenuron	50% : 25% DG +	Loveland (formerly Viterra)
	methyl : fluroxypyr : 2,4-D E	33.35% : 16.65% SG +	
Reward	diquat	240 g/L SN	True North
Rival	trifluralin	500 g/L EC, 10% G	Nufarm Agriculture
Roundup Transorb HC	glyphosate K+ salt	540 g/L SN	Monsanto
Roundup WeatherMax/Ultra 2	glyphosate K+ salt	540 g/L SN	Monsanto
R/T 540	glyphosate K+ salt	540 g/L SN	Monsanto
Rush 24	fluroxypyr : 2,4-D E	180 g/L EC + 660 g/L EC	MANA Canada
Rustler	glyphosate IPA + dicamba	194 g/L + 46 g/L SN	Monsanto
Salvo	2,4-D LV Ester	660 g/L EC	UAP
Select	clethodim	240 g/L EC	Arysta LifeScience
Sencor	metribuzin	75% DG	Bayer CropScience
Shadow RTM	clethodim	240 g/L EC	Loveland (formerly Viterra)
Sharpshooter	glyphosate IPA salt	356 g/L SN	UAP
Sharpshooter Plus	glyphosate IPA salt	360 g/L SN	UAP
Sierra 2.0	flucarbazone	397 g/L SC	Syngenta Crop Protection
Signal	clodinafop-propargyl	240 g/L EC	Nufarm Agriculture
Simazine 480	simazine	480 g/L SC	UAP
Simplicity	pyroxulam	30 g/L SC	Dow AgroSciences
Slam'R	clodinafop-propargyl	240 g/L EC	AgWest
Solo	imazamox	70% DG	BASF Canada
Spectrum	florasulam + clopyralid : MCPA E	50 g/L SC + 50 : 280 g/L EC	Dow AgroSciences
Spike-Up	tribenuron-methyl + glyphosate	75% DG + 540 g/L SN	Nufarm Agriculture
StartUp	glyphosate K+ salt	540 g/L SN	Loveland (formerly Viterra)
Stellar	florasulam + fluroxypyr	2.5 g/L SC + 100 g/L EC	Dow AgroSciences
Sword	dicamba : mecoprop : MCPA A	62.5 : 62.5 : 275 g/L SN	UAP
Tackle	glyphosate IPA + dicamba	140 g/L + 70 g/L SN	Syngenta Canada
Tandem	pyroxulam + fluroxypyr	30 g/L OD + 333 g/L EC	Dow AgroSciences
Target	dicamba : mecoprop-P : MCPA A	62.5 : 62.5 : 275 g/L SN	Syngenta Canada
Tensile	imazamox + clopyralid	70% DG + 75% DG	BASF Canada
Thrasher	bromoxynil : 2,4-D E	225 : 225 g/L EC	MANA Canada
Thumper	bromoxynil : 2,4-D E	280 : 280 g/L EC	Bayer CropScience
Topline	florasulam + MCPA ester	50 g/L SC + 600 g/L EC	MANA Canada

Product	Common Name	Formulation	Company
Topside	MCPB : MCPA A	375 : 25 g/L SN	UAP
Tordon 22K	picloram	240 g/L SN	Dow AgroSciences
Touchdown Total	glyphosate K + salt	500 g/L SN	Syngenta Canada
Tracker XP	dicamba : mecoprop-P : MCPA	62.5 : 62.5 : 275 g/L SN	IPCO
Traxion	glyphosate K + salt	500 g/L SN	Syngenta Canada
Traxos	pinoxaden : clodinafop-propargyl	25 : 25 g/L EC	Syngenta Canada
Treflan	trifluralin	480 g/L EC	Dow AgroSciences
TriCor	metribuzin	75% DG	United Phosphorus Inc.
Triton C	tribenuron methyl :	5.15% : 10.3% : 51.55% SG	E. I. duPont Canada
	thifensulfuron methyl : quinclorac		
Triton K	tribenuron methyl + dicamba + 2,4-D E or	50% SG + 480g/L SN + 660g/L EC	E. I. duPont Canada
	tribenuron methyl : dicamba + 2,4-D E	or 8.25% : 58.45% DG + 660g/L EC	
Trophy	fluroxypyr + MCPA E	180 g/L + 600 g/L EC	Nufarm Agriculture
Tropolox Plus	MCPB : MCPA A	375 : 25 g/L SN	Nufarm Agriculture
Tundra	fenoxaprop-p-ethyl : bromoxynil :	46 : 87.5 : 15.5 g/L EC	Bayer CropScience
	pyrasulfotole		
Turboprop	dichlorprop : 2,4-D E	300 : 282 g/L EC	UAP
Ultim	nicosulfuron : rimsulfuron	37.5% : 37.5% DG	E.I. duPont Canada
Vallera	flumioxazin	51% DG	Valent Canada
Vantage Plus Mux II	glyphosate DMA salt	480 g/L SN	Dow AgroSciences
Varro	thiencarbazone-methyl	10 g/L SC	Bayer CropScience
Vector	glyphosate DMA salt	480 g/L SN	Federated Coop
Velocity All-in-One	thiencarbazone-methyl :	5 : 31.3 : 175 g/L SC	Bayer CropScience
	pyrasulfotole : bromoxynil		
Velpar DF	hexazinone	75% DG	E I duPont Canada
Vigil WB	fenoxaprop-p-ethyl	120 g/L EC	IPCO
Viper ADV	imazamox : bentazon	20 g/L : 429 g/L SN	BASF Canada
VMD 480	dicamba (dimethylamine salt)	480 g/L SN	Adjuvants Plus (Gharda)
WildCat	fenoxaprop-p-ethyl	120 g/L EC	Loveland (formerly Viterra)
Wise-Up	glyphosate IPA salt	356 g/L SN	Adjuvants Plus
Yuma GL	quizalofop-p-ethyl	96 g/L EC	Gowan Canada

Fungicides

Product	Common Name	Formulation	Company
Acapela	picoxystrobin	250 g/L SC	E.I. duPont Canada
Acrobat 50 WP	dimethomorph	50% WP	BASF Canada
Allegro 500F	fluzazinam	40% SC	Syngenta Canada
Astound	cyprodinil : fludioxonil	37.5% : 25.0% DG	Syngenta Canada
Blanket	azoxystrobin + propiconazole	250 g/L SC + 418 g/L EC	MANA Canada
Bravo 500	chlorothalonil	500 g/L SC	Syngenta Canada
Bravo Zn	chlorothalonil	500 g/L SC	Syngenta Canada
Bumper 418 EC	propiconazole	418 g/L EC	Syngenta Canada
Cabrio Plus	pyraclostrobin : metiram	5% : 55% DG	MANA Canada
Canlus WDG Fungicide	boscalid	70% DG	BASF Canada
Caramba	metconazole	90 g/L EC	BASF Canada
Contans	Coniothyrium minitans strain	5.30% DG	BASF Canada
Copper 53W	tribasic copper sulphate	53% WP	UAP
Copper Spray	copper oxychloride	50% WP	UAP
Curzate 60 DF	cymoxanil	60% DF	UAP
Dithane Rainshield	mancozeb	75% DG	E.I. duPont Canada
Echo 720	chlorothalonil	720 g/L SC	Dow AgroSciences
Echo 90DF	chlorothalonil	90% DF	UAP
Folicur 423 F	tebuconazole	432 g/L F	Bayer CropScience
Folicur 250 EW	tebuconazole	250 g/L EW	Bayer CropScience
Fontelis	penthiopyrad	200 g/L SC	E.I. duPont Canada
I use	tebuconazole	432 g/L SC	Syngenta Canada
Gavel 75 DF	mancozeb : zoxamide	66.7% : 8.43% DF	Gowan Canada
Headline DUO ONE	boscalid : pyraclostrobin	25.2% : 12.8% DG	BASF Canada
Headline DUO TWO	boscalid	70% DG	BASF Canada
Headline EC	pyraclostrobin	250 g/L EC	BASF Canada
Inspire	difenoconazole	250 g/L EC	Syngenta Canada
Kocide 2000	copper hydroxide	35% DF	E.I. duPont Canada
Lance WDG	boscalid	70% DG	BASF Canada
Luna Tranquility	fluopyram : pyrimethanil	125 : 375 g/L SC	Bayer CropScience
MANA Azoxystrobin	azoxystrobin	250 g/L SC	MANA Canada
Manzate Pro-Stick	mancozeb	75% DF	United Phosphorus Inc.
Overall 240SC	iprodione	240 g/L SC	MANA Canada
Palliser	tebuconazole	432 g/L SC	Loveland (formerly Viterra)
Parasol WG	copper hydroxide	50% DG	Nufarm Agriculture
Penncozeb 75 DF	mancozeb	75% DG	UAP
Phostrol	mono- and dibasic sodium ,	53.6% F	Engage Agro Corporation
	potassium, and ammonium phosphites		
Pivot 418 EC	propiconazole	418 g/L EC	IPCO

Product

Priaxor DS
Proline 480 SC
Propel EC
Nufarm Propiconazole Fungicide
Propulse
Prosaro 250 EC
Quadris
Quadris Top
Quilt
Ranman 400SC
Reason 500SC
Regalia Maxx
Revus
Revus Top
Ridomil Gold 480EC/480 SL
Ridomil Gold/Bravo/
Ridomil Gold SL/Bravo
Rovral Flo
Rovral RX
Scala SC
Senator 70WP
Serenade Max/Serenade CPB
Stratego 250EC
Tanos 50 DF
Tattoo C
Tilt 250E
Torrent 400SC
Twinline
Vertisan
Zapro

Common Name

fluxapyroxad : pyraclostrobin
prothioconazole
propiconazole
propiconazole
prothioconazole : fluopyram
prothioconazole : lebuconazole
azoxystrobin
azoxystrobin : difenconazole
azoxystrobin : propiconazole
cyazofamid
fenamidone
Reynoutria sachalinensis
mandipropamid
mandipropamid + difenconazole
metalaxyl-M
metalaxyl-M + chlorothalonil

iprodione
iprodione
pyrimethanil
thiophanate-methyl
Bacillus subtilis
propiconazole : trifloxystrobin
famoxadone : cymoxanil
propamocarb : chlorothalonil
propiconazole
cyazofamid
pyraclostrobin : metconazole
penthiopyrad
dimethomorph : ametoctradin

Formulation

250 : 250 g/L SC
480 g/L SC
250 g/L EC
418 g/L EC
200 : 200 g/L SC
125 : 125 g/L EC
250 g/L SC
200 : 125 g/L SC
75 : 125 g/L SC
400 g/L SC
500 g/L SC
20% SC
250 g/L
250 g/L SC + 250 g/L EC
480 g/L EC, 480 g/L SC
480 g/L + 500 g/L SC

240 g/L SC
240 g/L SC
400 g/L SC
70% WP
14.6% WP / 1.34% SC
125 : 125 g/L EC
25% : 25% DF
375 : 375 g/L SC
250 g/L EC
400 g/L SC
130 : 80 g/L F
200 g/L EC
225 : 300 g/L SC

Company

BASF Canada
Bayer CropScience
Loveland (formerly Viterro)
Nufarm Agriculture
Bayer CropScience
Bayer CropScience
Syngenta Canada
Syngenta Canada
Syngenta Canada
FMC of Canada
Bayer CropScience
Engage Agro Corporation
Syngenta Canada
Syngenta Canada
Syngenta Canada
FMC of Canada
Monsanto
Bayer CropScience
Engage Agro Corporation
Bayer CropScience
Bayer CropScience
E.I. duPont Canada
Bayer CropScience
Syngenta Canada
Engage Agro Corporation
BASF Canada
E.I. duPont Canada
Engage Agro Corporation

Seed Treatments**Product**

Acceleron DX-910
Acceleron Seed Treatment
Package
Admire SPT
Agrox B-2
Agrox CD
Agrox FLn
Alias 240 SC
Allegiance FL
Apron Advance
Apron Maxx RTA
Apron Maxx RTA
Armour
Armour RTU
Belmont 2.7TS
Charter RTU
Confine

Crown
Cruiser SF5
Cruiser Maxx Beans
Cruiser Maxx Cereals/
Cruiser Maxx Cereals
Commercial
Cruiser Maxx D Potatoes

Cruiser Maxx Pulses

Cruiser Maxx Vibrance Beans
Cruiser Maxx Vibrance Cereals

DB-Red L
DCI
Dividend XL RTA
EverGol Energy
Gaucho CS II
Gemini
General Storage Disinfectant
Heads Up Plant Protectant

Common Name

Bacillus subtilis
Bacillus subtilis + thiamethoxam :
difenoconazole : metalaxyl-M : fludioxonil
imidacloprid
captan : diazinon
captan : diazinon
captan
imidacloprid
metalaxyl
fludioxonil : metalaxyl-M : thiabendazole
fludioxonil : metalaxyl-M
fludioxonil : metalaxyl-M
triticonazole
triticonazole
metalaxyl
triticonazole
mono and di-potassium salts of
phosphoric acid
carbathiin : thiabendazole
thiamethoxam
thiamethoxam : metalaxyl-M : fludioxonil
thiamethoxam : difenoconazole :
metalaxyl-M

fludioxonil : difenoconazole +
thiamethoxam
thiamethoxam +
fludioxonil : metalaxyl-M
thiamethoxam : metalaxyl-M :
fludioxonil + sedaxane
thiamethoxam : difenoconazole :
metalaxyl-M : sedaxane
maneb
diazinon : captan : thiophanate-methyl
difenoconazole : metalaxyl-M
penflufen : prothioconazole : metalaxyl
imidacloprid : carbathiin : thiram
triticonazole : thiram
dimethyl benzyl ammonium chloride
saponins of *Chenopodium quinoa*

0.18%
0.18% + 20.7% :
1.25% : 0.39% : 0.13% FS
240 g/L FS
33.5% : 11% WP
15% : 15% DS
30% FS
240 g/L FS
317 g/L FS
25 : 20 : 150 g/L FS
2.31% : 3.46% FS
0.73% : 1.10% FS
25 g/L FS
16.8 g/L FS
317 g/L FS
16.8 g/L FS
45.8% F

92 : 58 g/L FS
47.6% FS
22.6% : 1.70% : 1.12% FS
2.80% : 3.36% : 0.56% FS

19.4 : 19.4 g/L +
240 g/L FS
47.6% +
0.73% (or 2.31%) : 1.10% (or 3.46%) FS
22.6% : 1.70% : 1.12% + 500 g/L FS

30.7 : 36.9 : 9.5 : 8.0 g/L FS

323 g/L F
6% : 18% : 14% DS
3.37% : 0.27% FS
38.4 : 76.8 : 61.4 g/L FS
285.7 : 47.6 : 95.3 g/L FS
1.25% : 12.5% FS
10% F
63.02% WSP

Formulation**Company**

Monsanto
Monsanto

Bayer CropScience
Norac Concepts Inc.
Norac Concepts Inc.
Norac Concepts Inc.
MANA Canada
Bayer CropScience
Syngenta Canada
Syngenta Canada
Syngenta Canada
Loveland (formerly Viterro)
Loveland (formerly Viterro)
Chemtura Canada Co.
BASF Canada
Winfield Solutions

Chemtura Canada Co.
Syngenta Canada
Syngenta Canada
Syngenta Canada

Syngenta Canada

Syngenta Canada

Syngenta Canada

UAP
Norac Concepts Inc.
Syngenta Canada
Bayer CropScience
Bayer CropScience
BASF Canada
UAP
Engage Agro Corporation

Product	Common Name	Formulation	Company
Helix Vibrance	thiamethoxam : difenoconazole :	20.7% : 1.25% : 0.39% : 0.13% +	Syngenta Canada
Helix XTra	metalaxyl-M : fludioxonil + sedaxane	500 g/L FS	Syngenta Canada
Insure Cereal	thiamethoxam : difenoconazole :	20.7% : 1.25% : 0.39% : 0.13% FS	Syngenta Canada
Kocide 2000	metalaxyl-M : fludioxonil	17 : 17 : 10 g/L F	BASF Canada
Lumiderm	copper hydroxide	35% DF	E.I. duPont Canada
Maxim D	cyantraniliprole	625 g/L SC	E.I. duPont Canada
Maxim Liquid PSP	fludioxonil : difenoconazole	19.4 : 19.4 g/L FS	Syngenta Canada
Maxim MZ PSP	fludioxonil	40.3% F	Syngenta Canada
Maxim PSP	fludioxonil : mancozeb	0.5% : 5.7% DS	Syngenta Canada
Maxim Quattro	fludioxonil	0.5% DS	Syngenta Canada
Mertect SC	thiabendazole : fludioxonil :	26.5% : 3.32% : 2.65% : 1.33 % FS	Syngenta Canada
Polyram 16D	metalaxyl-M : azoxystrobin	500 g/L SC	Syngenta Canada
Poncho 600 FS	thiabendazole	16% DS	BASF Canada
Potato ST 16	metiram	16% DS	Bayer CropScience
Proseed	clothianidin	600 g/L FS	UAP
Prosper EverGol	mancozeb	16% DS	Syngenta Canada
Prosper FX	fludioxonil	40.3% LS	Bayer CropScience
Rancona Apex	clothianidin : penflufen :	290 : 10.7 : 7.15 : 7.15 g/L FS	Bayer CropScience
Rancona RS	trifloxystrobin : metalaxyl	285.7 : 50 : 7.14 : 5.36 g/L FS	Bayer CropScience
Raxil MD	clothianidin : carbathiin :		
Raxil PRO	trifloxystrobin : metalaxyl		
Raxil PRO Shield	ipconazole	4.61 g/L FS	Chemtura Canada Co.
Raxil WW	ipconazole : carbathiin	9.38 : 87.5 g/L FS	Chemtura Canada Co.
Senator PSPT	tebuconazole : metalaxyl	5.0 : 6.6 g/L FS	Bayer CropScience
Solan MZ	tebuconazole : prothioconazole : metalaxyl	3.0 : 15.4 : 6.2 g/L FS	Bayer CropScience
Sombrero 600 FS	tebuconazole : prothioconazole :	3.0 : 15.4 : 6.2 +	Bayer CropScience
StorOx	metalaxyl + imidacloprid	600 g/L FS	Bayer CropScience
Stress Shield 600	tebuconazole : metalaxyl + imidacloprid	5.0 : 6.6 + 480 g/L FS	Bayer CropScience
Thiram 75WP	thiophanate-methyl	10% DS	Engage Agro Corporation
Titan ST	mancozeb	16% DS	Norac Concepts Inc.
Trilex AL	imidacloprid	600 g/L FS	MANA Canada
Trilex EverGol	hydrogen peroxide	27% LS	Brenntag Canada
Tuberseal	imidacloprid	600 g/L FS	Bayer CropScience
Vibrance 500FS	thiram	75% WP	Chemtura Canada Co.
Vibrance Muxx RTA	clothianidin	600 g/L FS	Bayer CropScience
Vibrance Muxx RJC	trifloxystrobin : metalaxyl	13.5 : 10.8 g/L FS	Bayer CropScience
Vibrance XL	penflufen : trifloxystrobin + metalaxyl	154 : 154 + 317 g/L FS	Bayer CropScience
Vibrance XL Co-Pack	mancozeb	16% DS	Norac Concepts Inc.
Vitaflo 220	sedaxane	500 g/L FS	Syngenta Canada
Vitaflo 280	fludioxonil : metalaxyl-M + sedaxane	0.73% : 1.10% + 500 g/L LS	Syngenta Canada
Vitaflo Fungicide	fludioxonil : metalaxyl-M + sedaxane	2.31% : 3.46% + 500 g/L LS	Syngenta Canada
Vitaflo SP Fungicide	difenoconazole : metalaxyl-M : sedaxane	66.2 : 16.5 : 13.8 g/L FS	Syngenta Canada
	difenoconazole : metalaxyl-M + sedaxane	3.37% : 0.27% + 500 g/L FS	Syngenta Canada
	carbathiin : thiram	220 : 200 g/L FS	Chemtura Canada Co.
	carbathiin : thiram	15.59% : 13.25% FS	Chemtura Canada Co.
	carbathiin : thiram	15.59% : 13.25% FS	Loveland (formerly Viterro)
	carbathiin : thiram	15.59% : 13.25% FS	IPCO

Insecticides

Product	Common Name	Formulation	Company
Actara 240 SC/Actara 25 WG	thiamethoxam	240 g/L SC, 25% DG	Syngenta Canada
Admire 240 / SPT	imidacloprid	240 g/L SC	Bayer CropScience
Alias 240 SC	imidacloprid	240 g/L SC	UAP
Ambush	permethrin	500 g/L EC	Ambac Chemical Corp.
Assail	acetamiprid	70% WP	Engage Agro / Nippon
Beleaf	flonicamid	50% DG	Soda Co. Ltd
Citadel 480EC	chlorpyrifos	480 g/L EC	FMC of Canada / UAP
Clutch	clothianidin	50% DG	IPCO
Concept	imidacloprid : deltamethrin	75 g/L SC : 10 g/L SC	Valent Canada Inc.
Coragen	chlorantraniliprole	200 g/L SC	Bayer CropScience
Cygon 480EC / Cygon 480-Ag	dimethoate	480 g/L EC	E.I. duPont Canada
Decis	deltamethrin	50 g/L EC	IPCO, Cheminova Canada
Delegate	spinetoram	25% SG	Bayer CropScience
Dibrom	naled	864 g/L EC	Dow AgroSciences
Dipel 2X DF	Bacillus thuringiensis	32 billion CIU/kg DG	UAP
Eco Bran	carbaryl	2% spreadable bran bait	Valent BioSciences
Entrust	spinosad	80% WP	Peacock Industries
Fulfill	pymetrozine	50% DG	Dow AgroSciences
Grapple / Grapple ₂	imidacloprid	240 g/L SC	Syngenta Canada
Imidan	phosmet	50% WP	Chemimova Canada
Insecto	diatomaceous earth	90%	UAP
			Natural Insecto Products Inc.

Lagon 480E	dimethoate	480 g/L EC	UAP
Lannate	methomyl	90% WSP	E.I. duPont Canada
Lorsban 4E	chlorpyrifos	480 g/L EC	Dow AgroSciences
Malathion 85E / 500	malathion	85%, 500 g/L EC	UAP, IPCCO
Matador	lambda-cyhalothrin	120 g/L EC	Syngenta Canada
Movento 240 SC	spirotetramat	240 g/L SC	Bayer CropScience
MPOWER Krypton	chlorpyrifos	480 g/L	NewAgco Inc.
Nolo Bail	spores of <i>Nosema (Paranosema) locustae</i> Canning	Minimum of 2.2x10 ⁶ on coated wheat bran	M&R Durango, Inc.
Nufos 4E	chlorpyrifos	480 g/L EC	Cheminova Canada
Oberon	spiromesifen	240 g/L SC	Bayer CropScience
Orthene	acephate	75% WSP	UAP
Perm-LIP	permethrin	384 g/L EC	United Phosphorous Inc
Phostoxin	aluminum phosphide	55% tablets	Degesch America Inc
Pounce	permethrin	384 g/L EC	PMC of Canada
Protect-It	diatomaceous earth	90%	Hedley Technologies Ltd
Pyrifos 15G	chlorpyrifos	15% G	UAP
Pyrinex 480EC	chlorpyrifos	480 g/L EC	MANA Canada
Rimon 10 EC	novaluron	10% EC	Chemtura, MANA Canada
Ripcord	cypermethrin	407 g/L EC	Engage Agro Corp.
Sevin XLR PLUS	carbaryl	480 g/L SC	Bayer CropScience
Silencer 120 EC	lambda-cyhalothrin	120 g/L EC	MANA Canada
Sluggo Professional	ferric phosphate	0.76 % granules	Engage Agro Corp.
Success 480 SC	spinosad	480 g/L SC	Dow AgroSciences
Tempo	cyfluthrin	20% WP	Bayer CropScience
Thimet 15G	phorate	15% G	Amvac Chemical Corporation
Thiodan 4EC	endosulfan	400 g/L EC	Bayer CropScience
Thionex EC / 50 WP	endosulfan	400 g/L EC, 50% WP	UAP
UP-Cycle	cypermethrin	250 g/L EC	United Phosphorous Inc
Vydale L	oxamyl	240 g/L SN (water soluble liquid)	E.I. duPont Canada
Warhawk	chlorpyrifos	480 g/L EC	Loveland Products / UAP

Abbreviations

WP wettable powder	EC emulsifiable concentrate	WSP water soluble powder	SN solution	A amine
DG water dispersible granule	G granule	DF dry flowable	F flowable	E ester
WSG water soluble granule	SC suspension concentrate	SG soluble granule	OD oil dispersion	IS solution for seed treatment
DS powder/dust for dry seed treatment	IW emulsion, oil in water	FS flowable suspension concentrate for seed treatment	SL soluble concentrate	

KEY TO PRODUCT PAGES

Pesticide Product Name

This field lists the pesticide product name. Where there is only one product the commercial "trade" name is given. Where more than one company sells pesticides with the same combination of active ingredients the "generic" (active ingredient) name is given.

If the active ingredients are all in a common formulation (liquid, granule, etc.) the generic name will appear as 'Ingredient A/Ingredient B' and if the active ingredients are in separate containers to be mixed in the sprayer the names are given as 'Ingredient A + Ingredient B'.

Pesticide Resistance Group –

This area will the pesticide active ingredient(s) to the mode of action that ingredient uses and refer to a page number where more information can be found.

Company:

This section identifies the company (or companies) that manufacture or market this crop protection product (or generic equivalents) in Canada as well as the PCP# for that (those) product(s). See page 9 for more information on PCP numbers. PCP#s are given as '(PCP#XXXXXX)' where XXXXX is a four or five digit number unique to that product. In some cases, where there are multiple components with separate PCP numbers, the PCP number will be provided below under 'Formulation.'

Formulation:

This section gives information on the active ingredient and its concentration in the product as well as information on formulation type and packaging types and configurations. Formulation strength (or concentrations) are given in % by weight for dry formulations and g/L for liquid formulations. PCP numbers may also be give for some products (see above).

Crops and Staging:*

This section indicates on which crops the product may be used and what stage of crop development it should be applied at. Rates may also be included in this section if they vary between crop types or crop stage.

**This section will also indicate which crops are registered under the User Requested Minor Use Label Expansion (URMULE) program. Some companies, as a condition of placing these minor crops on their labels request, that users of their product on these crops do so at their own risk because the registration was approved with information the company did not produce.*

These crops will be flagged separately from the main crops.

Pest (Diseases, Insects, Weeds) and Staging:

This section indicates the pests (Diseases, Insects, or Weeds) that are indicated on the product label as controlled or suppressed, as well as any specifics on the timing of application relative to the pest stage if required. Rates may also be included in this section if they differ for different pests or stage of pest.

Rates:

The rates provided in this section are given in the amount of product required per acre and the number of acres treated per package unit where possible. This section will also indicate any adjuvants that are to be used in conjunction with the product and the rate of that adjuvant.

This section will not be present if rates have been integrated into either of the previous **Crops** or **Pest** sections.

Application Information:

Water Volume: This section indicates the minimum carrier water volume to be used to apply the product. Using less than the recommended minimum carrier application volume can negatively affect pesticide performance, particularly with contact pesticides and when using low drift nozzles.

Nozzles and Pressure: This section indicates if there are any particular nozzles that should or should not be used to apply the product. Pressures indicated reflect those for conventional nozzles. Low drift nozzles may require higher pressures for proper performance. A general statement of "Use nozzles and pressures designed to deliver proper coverage with ASABE ____ droplets" indicating the ideal droplet sizes to allow for the combination of lowest drift potential and best performance from the pesticide. ASABE

refers to the American Society of Agricultural Engineers who have set standards a series of droplet measurements (in microns or micrometres) that classify droplet sizes from 'fine' to 'very coarse'.

How it Works:

This section typically refers to the page where a general description of the various modes of action of either herbicide, fungicides or insecticides.

Effects of Growing Conditions:

This section summarizes any adverse conditions that will affect the biological function of the crop or the target pest and therefore possibly impact the product's performance. In most cases both crop and target pest must be growing or functioning normally for pesticides to provide expected performance and/or crop tolerance. Adverse weather conditions such as extreme heat, cold, drought or flooding can slow or stop the biological processes in the crop or pest. These biological processes in the crop allow the pesticide to be degraded quickly. If biological processes that are attacked by the pesticide, and under normal conditions would kill the pest, are not functioning normally the pest may be able to rid itself of the pesticide before dying and recover from the application.

Tank Mixes:

This section indicates which other pesticides the pesticide label indicates are registered for use as tank mix combinations with this pesticide.

Common mixes may include:

Herbicides:

(Subtitles may indicate specific crops or condition restrictions.)

Insecticides:

Fungicides:

Fertilizers:

There may be additional pesticides that are registered but not listed on this product's label. Other pesticides may have this product listed as a mix option on their labels. The note below (in bold) directs users to a chart inside the back cover that show all available mixes for this pesticide. The product listed on the left column of the chart is the product that supports the mix. Mixes supported by both products are marked with an 'X'. Mixes supported by only one of the products is indicated by an arrow pointing to the left column.

Included in the tank mix section in non-bolded italics may be any precautions against the mixing of pesticides which will have adverse reactions such as crop injury, reduced pest control or unusual increased danger in the use of the product.

Note: The above mixes are those listed on the pesticide label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Since most pesticides have a capacity to injure neighboring plants, wildlife or people, they will come with restrictions on their use in order to prevent this unintentional damage. Misuse of pesticides may result in as little as temporary or superficial damage to plants or a slight irritation to the eyes or nose, or could also result in poor performance of the pesticide, severe injury and/or yield loss to very sensitive plants and/or unacceptable residues in agricultural commodities, and/or serious illness or death of non-target organism or people. It is important to comply with product restrictions in order to minimize the impact of the pesticide used on non-target organisms and people. A selection of common restrictions and precautions found on product labels are provided in this section, *but it is important to read the label carefully in order to understand how to use the product properly.*

Rainfall: This section indicates the required delay between application and rainfall to avoid reductions in the performance of the product or the unintentional movement of the product.

Re-entry: This section indicates when it is safe for a person to re-enter treated field following an application of a particular pesticide without the same personal protection used to apply the product.

Resistance Management: This section highlights products where an increased risk of the target pests developing resistance to the group of products (typically fungicides) has been identified. If no specific risk has been identified the reader is referred to a general resistance section. All pesticides have some risk of the target pest developing resistance. Rotating pesticide groups using as many different resistance groups as possible in the rotation is one way to avoid or delay resistance development.

Grazing: This section indicates whether and how soon treated crops may be grazed by livestock or otherwise fed to livestock. This restriction is in place to avoid residues of the pesticide from being detected in milk or meat from animals consuming forage, greenfeed or straw from treated crops or forage.

Pre-harvest interval: Is the time that must be left between application of a pesticide and the harvest of a crop in order to prevent greater than allowable residues of the pesticide in the harvested material. Harvest is the cutting of the crop (i.e. combining or hay cut) or removal of the harvestable material from the plant (i.e. picking fruit or stripper header). Maximum Residue Limits (MRLs) are set for commodities based on registered rates and staging of pesticides used in the production of those commodities. Disregarding these intervals can result in residues over the MRLs, which can lead to market disruptions.

Re-cropping: This section indicates how soon specific crops may be seeded into treated fields. Failure to adhere to these delays could result in injury to the following crop.

Aerial Application: This section indicates whether the product may be applied by aircraft and any special conditions that may be necessary.

Labelling: In addition to other precautions and warnings, seed treatment products will also have statements about how seed treated with the product should be labeled.

Storage: This section indicates how the product must be stored. As a general rule, unused pesticides should always be stored in their original containers in a secure, dry area, away from other pesticides, food or feed.

Buffer Zones: This section will indicate any setback distances that are required from sensitive aquatic or upland habitats. Newer labels may indicate that these distances are from the downwind edge of the boom but older labels may not. Examples of aquatic habitats are lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands. Examples of terrestrial habitats are grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands.

In addition to the set back or 'buffer' distances indicated on product labels, provincial environment departments may also have additional restrictions or requirements for permits to apply pesticides to or near water. Check with the provincial environment department/ministry for more information.


Tank Cleaning:

This section describes the measures that are required to properly clean out spray tanks. A general overview of sprayer cleaning is given on page 14, but products where there is a high risk of crop damage as a result of very low level contamination of the spray solution, will have specific measures indicated.

Hazard Rating:

This section indicates the relative toxicity of the pesticide, formulations or components. For an explanation of the symbols used here see page 10. An additional symbol that is used that is not a standard symbol is the (!) exclamation mark which indicates an otherwise undefined risk factor (i.e. irritation).

Example:

 Caution – Eye Irritant

Some older products have not had hazard ratings developed while others products have very low toxicity and do not have hazard warnings. Even in the absence of a hazard rating users should wear a minimum of nitrile gloves and an apron as well as long sleeved apparel during mixing and avoid unnecessary exposure.

Weed Control

The use of in-crop herbicides to control weeds is often very important in determining the success or failure of a crop. However, many other practices that can be implemented before and after a herbicide application can help to reduce weed competition. The use of these practices is termed Integrated Weed Management.

Integrated Weed Management

A farming system that utilizes an array of inter-dependent cultural, biological and herbicidal weed control practices is implementing Integrated Weed Management (IWM). It is essential that IWM involves an array of tools including the rotation of available herbicide groups, ensuring that weeds are exposed to a diverse range of control mechanisms. The principal aim of IWM is to improve the health and vigour of crops so that they may out-compete weeds emerging in the stand. This helps to reduce selection for resistance to any single control agent and to delay or prevent the development of herbicide resistant weeds.

Practising IWM does not mean abandoning chemical weed control, just relying on it less exclusively. For example:

- You may decide to choose a taller wheat variety or a tall, viny pea variety for a certain field. These crop selections will compete strongly with weeds, possibly allowing you to skip a spray operation in more competitive crops.
- You could insert a short-term forage crop into your crop rotation. Studies show that short-term alfalfa stands can reduce wild oat and green foxtail populations by up to 80 percent the year after breaking.
- Early sown barley may give you enough of a "jump" on the weeds that you can avoid herbicide applications.
- Use of vigorous, high-quality seed, sown shallow, can give you better crop competition than poor-quality or deeply sown crop seed.
- Banding nitrogen near the seed can give your crop an advantage over weeds.

For more information, refer to "Integrated Weed Management: Making it Work on Your Farm" factsheet, available from both Manitoba Agriculture, Food and Rural Development and Saskatchewan Ministry of Agriculture.


Making Spray Decisions

Field Scouting

Field scouting is an important tool for making informed spray decisions. Accurately assessing the type and number of weeds in the field will help you determine if a spray operation is necessary. The scouting pattern diagram on this page provides a guideline for scouting a field. The entire field should be walked to get a feel for the distribution and species of the weeds present. A minimum of 20 weed counts

should be taken across the field. A smaller number may be used, but be aware that accuracy decreases as the number of counts gets smaller. Count the number of weeds in a 1 m² or a 0.25 m² area and divide the total number of weeds by the number of counts taken to obtain an average for the field. If using 0.25 m² samples, make sure to multiply by four so your average is for a 1 m² area.

Some weeds are not distributed uniformly and may be found in patches (for example, Canada thistle) or in low spots. As well, the type and number of weeds found along the field edges may be very different from those found inside the field. These areas should be considered separate from the rest of the field. If possible, patches, low spots, and field borders should be treated separately, as field wide spraying may not be required. Look out for new invading weeds and patches of herbicide-resistant weeds. Herbicide-resistant weeds and new invaders should be removed (manually if necessary), regardless of their number, to prevent them from spreading and becoming a serious control problem. Mapping your field's weed problems will allow you to monitor the spread of weed patches over time and help you assess the effectiveness of your control program.



Yield Losses Caused by Weeds

Knowing the amount of crop yield loss caused by a given weed density will help you decide if a spray operation is required. The tables on the following pages give an indication of the yield loss caused by some of the important grassy weeds.

THESE TABLES SHOULD BE USED ONLY AS A GUIDE. The figures are based on Western Canadian research trials and will not be accurate all of the time. The yield loss values apply only to healthy, well fertilized crops with good stand establishment. Crops that are diseased or emerged unevenly will not compete well with weeds and will suffer larger yield losses than indicated in these tables. The yield loss figures are based on competition from single weed species only. Other weeds, such as wild mustard or Canada thistle, must be controlled if the figures are to be accurate. As well, the tables are based on competition from normal height crops. Semi-dwarf or hybrid varieties may not compete as well with weeds and the figures may not be accurate in these cases.

Table 1. Yield Losses (Percent) in Wheat Caused by Wild Oats.

	Wild Oat Density – Number Per Square Metre																
	1	2	4	6	8	10	12	14	16	18	20	25	30	35	40	45	50
Wild Oats are 1 Leaf Stage Ahead of the Crop	1	2	4	6	8	10	12	14	15	17	19	22	26	29	32	34	37
Wild Oats are the Same Leaf Stage as the Crop	1	1	2	4	5	6	7	8	9	10	11	14	16	18	20	22	24
Wild Oats are 1 Leaf Stage Behind the Crop	0	1	1	2	3	3	4	5	5	6	7	8	10	11	13	14	15

Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Figure 1. Spray Decision Guideline for Wild Oats in Wheat.





Table 2. Yield Losses (Percent) in Wheat Caused by Green Foxtail (Wild Millet).




Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Table 3. Yield Losses (Percent) in Barley Caused by Wild Oats.

Crop Density (plants/m ²)	Relative Emergence	Wild Oat Density (plants/m ²)											
		1	5	10	15	20	25	30	40	50	70	100	150
300	Wild Oats are 1 Leaf Stage Ahead of the Crop	0.3	1.4	2.8	4.1	5.4	6.7	8.0	10.3	12.6	16.8	22.4	30.2
	Wild Oats are the Same Leaf Stage as the Crop	0.3	1.3	2.5	3.7	4.8	6.0	7.1	9.2	11.3	15.1	20.3	27.6
	Wild Oats are 1 Leaf Stage Behind the Crop	0.2	0.9	1.7	2.6	3.4	4.2	5.0	6.6	8.1	11.0	15.0	20.9
225	Wild Oats are 1 Leaf Stage Ahead of the Crop	0.4	1.9	3.6	5.4	7.0	8.6	10.2	13.1	15.9	20.9	27.4	36.2
	Wild Oats are the Same Leaf Stage as the Crop	0.3	1.6	3.1	4.6	6.1	7.5	8.8	11.4	13.9	18.4	24.4	32.6
	Wild Oats are 1 Leaf Stage Behind the Crop	0.2	1.0	2.0	3.0	4.0	4.9	5.8	7.6	9.3	12.6	17.1	23.6
175	Wild Oats are 1 Leaf Stage Ahead of the Crop	0.5	2.3	4.6	6.7	8.7	10.7	12.5	16.1	19.3	25.1	32.3	41.8
	Wild Oats are the Same Leaf Stage as the Crop	0.4	1.9	3.8	5.6	7.3	8.9	10.5	13.6	16.4	21.6	28.2	37.1
	Wild Oats are 1 Leaf Stage Behind the Crop	0.2	1.1	2.3	3.4	4.4	5.5	6.5	8.5	10.4	14.0	18.9	25.9

Table 4. Yield Losses (Percent) in Barley Caused by Green Foxtail (Wild Millet).



Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Table 5. Yield Losses (Percent) in Canola Caused by Wild Oats and Volunteer Cereals.

	Weed Density – Number Per Square Metre												
	1	2	4	6	8	10	12	14	16	18	20	25	30
Wild Oats	3	5	6	8	9	10	11	12	13	14	15	16	18
Volunteer Wheat	1	3	6	8	10	11	12	14	15	16	17	19	21
Volunteer Barley	3	5	8	10	12	14	15	17	18	19	20	23	25

Sources: Dew and Keys, Agriculture Canada (Lacombe, Alberta)
O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)

Figure 2. Spray Decision Guideline for Wild Oats and Volunteer Cereals in Canola.





Table 6. Yield Losses (Percent) in Canola Caused by Green Foxtail (Wild Millet)



Source: O'Donovan, Alberta Environmental Centre (Vegreville, Alberta)


Table 7. Yield Losses (Percent) in Flax Caused by Wild Oats and Volunteer Cereals.

	Weed Density – Number Per Square Metre									
	1	2	3	4	5	6	7	8	9	10
Wild Oat	6	8	10	12	13	15	16	17	18	19
Volunteer Wheat	6	11	15	18	22	24	27	29	31	33
Volunteer Barley	6	12	16	21	24	28	31	34	36	39

Sources: Dew and Keys, Agriculture Canada (Lacombe, Alberta)

Friesen et al., University of Manitoba (Winnipeg, Manitoba)

Figure 3. Spray Decision Guideline for Wild Oats and Volunteer Cereals in Flax.



Deciding to Spray – Economic Thresholds and Herbicide Resistance

An **economic threshold** is the level of infestation at which lost yield exceeds the cost of the chemical and its application. Determining the economic threshold will help you decide if a spray operation is necessary.

The following example outlines how to determine an economic threshold:

You have a wild oat problem in your wheat. After a thorough field scouting you have determined that your field has an average density of 35 wild oats per square metre. You know that the crop and weeds are at the same leaf stage. Using Table 1, choose the "Same Leaf Stage" row and read across to 35 wild oats per square metre. You will find that your yield loss will be about 18 percent.

You think it could be a 40 bushel per acre wheat crop, and expect to get \$4 per bushel for it. Therefore:

$$\begin{aligned} 40 \text{ bushels} \times 0.18 (\text{percent of expected yield loss}) &= \\ 7.2 \text{ bushels per acre of lost yield} & \\ 7.2 \text{ bushels} \times \$4 \text{ per bushel} &= \\ \mathbf{\$28.80 \text{ per acre of lost income}} & \end{aligned}$$

Now find out the price of your herbicide. Most wild oat herbicides for wheat cost between \$20 to \$25 per acre. In this case, lost income exceeds the cost of the herbicide and application, so spraying would be justified.

Alternatively, you may want to use the figures provided with some of the yield loss tables. These figures provide flowcharts to assist you in making spray decisions. In some cases the flowcharts may indicate to spray when you do not have an economic threshold weed density, but most times they will prevent you from spraying unnecessarily.

Another factor to consider when deciding whether to spray is your herbicide rotation. A one in three rotation of herbicide groups is currently recommended to delay the development of herbicide resistance for weeds such as wild oats and green foxtail. Skipping a spray operation will give you an extra year of flexibility in your herbicide rotation. This means that you have one extra herbicide group to choose from the year after you skipped the spray operation. When making spray decisions, the ability to rotate herbicides should be considered in addition to the economics of spraying.

Making the Spray Decision

Remember that economic thresholds should be used only as guides when making a spray decision. Lost income caused by dockage or downgrading must also be considered. **FIELDS THAT ARE NOT SPRAYED THIS YEAR HAVE A HIGHER POTENTIAL FOR PROBLEMS THE FOLLOWING YEAR BECAUSE OF WEED SEED RETURN.** A farmer's experience and common sense play an important role when deciding to spray. Used properly, however, the economic threshold can be an important tool in making spray decisions.

Weed Resistance to Herbicides

In recent years, the number of herbicide-resistant weeds and the areas they infest have increased.

Most herbicide-resistant weed infestations have developed following repeated use of the same herbicide (or herbicide group) for a number of years on the same field. Growers who have developed weed resistance on their farms will typically see a weed, which is normally controlled by a herbicide, escape uncontrolled after a number of years of use of the same product or product group. Individual plants may be resistant to 1.5 up to 10 or more times the normal field rate.

Herbicide Groups

To help you plan your herbicide program, the following table lists "herbicide groups." To slow down the process of developing weed resistance, use products from different groups from year to year on your fields.

Table 1: Herbicide Groups Based on Mode of Action

Group 1 (contain ACCase grasskillers) Achieve Liquid Gold*, Axial, Axial iPak*, Axial Xtreme*, Broadband*, clethodim, clodinafop, Equinox, fenoxaprop, Harmony (SG*, K* and Max*), Odyssey DLX*, Poast Ultra, quizalofop, tralkoxydim, Traxos, Tundra
Group 2 (contain ALS/AHAS inhibitors) Accent, Adrenalin*, Altitude FX/FX2*, Ares, Assert FL*, Barricade*, Battalion*, Broadband*, Escort, Everest GBX*, Express Pro, florasulam + glyphosate, florasulam + MCPA ester, flucarbazone, Frontline 2,4-D*, Harmony (SG*, K* and Max*), imazethapyr, imazethabenz, Inferno Duo, Ko-Act*, metsulfuron, Muster, Odyssey, Odyssey DLX*, Option, Pinnacle, Prism, Retain*, Simplicity, Solo, Spectrum*, Stellar*, Tandem*, Tensile*, tribenuron, thifensulfuron/tribenuron, thifensulfuron/tribenuron + MCPA*, Triton C*, Triton K*, Ultim, Varro, Velocity m3*, Viper ADV*
Group 3 (contain mitotic inhibitors) Bonanza, Edge, Fortress*, Rival, Treflan
Group 4 (contain growth regulator herbicides) 2,4-D, 2,4-DB, Achieve Liquid Gold*, Adrenalin*, Altitude FX/FX2*, Assert FL*, Axial Xtreme*, Barricade*, Battalion*, BlackHawk*, bromoxynil/2,4-D ester*, bromoxynil/MCPA ester*, Curtail M, dicamba, dicamba/mecoprop-P/MCPA, dichlorprop/2,4-D, Distinct*, DyVel, DyVel DSp, Eclipse III*, Enforcer D & M*, Everest GBX*, florasulam + MCPA ester, fluroxypyr + 2,4-D, Frontline 2,4-D*, glyphosate/dicamba*, Grazon, Harmony K*, Harmony Max*, Ko-Act*, Lontrel, MCPA, MCPB/MCPA, mecoprop-P, Momentum, Optica Trio, OverDrive*, Prestige XC, Pulsar, Restore, Retain*, Spectrum*, Stellar*, Tandem*, Tensile*, thifensulfuron/tribenuron + MCPA*, Tordon 22K, Triton C*, Triton K*, Trophy

Continued on next page

Group 5 (contain photosynthetic inhibitors – triazines) AAtrex, metribuzin, Primextra II Magnum*, simazine, Velpar
Group 6 (contain photosynthetic inhibitors – nitriles/benzothiadiazoles) Achieve Liquid Gold*, Axial iPak*, Basagran / Basagran Forte, bromoxynil, bromoxynil/2,4-D ester*, bromoxynil/MCPA ester*, Enforcer D & M*, Infinity*, Tundra*, Velocity m3*, Viper ADV*
Group 7 (contain photosynthetic inhibitors – ureas/amides) linuron
Group 8 (unknown mode of action) Avadex BW, Avenue, Eptam-8E, Fortress*
Group 9 (contain inhibitors of EPSP synthase) Glyphosate - several brands. Cleanstart*, Eclipse III*, FlexStar GT*, florasulam + glyphosate, glyphosate/dicamba*
Group 10 (contain inhibitors of glutamine synthetase) GoodHarvest, Liberty
Group 11 (inhibit carotenoid synthesis - triazoles) Amitrol 240
Group 14 (contain inhibitors of protoporphyrinogen oxidase/PPO/Protox) Aim, Authority, BlackHawk*, Blazer, CleanStart*, FlexStar GT*, flumioxazin, Focus*, Heat, Reflex
Group 15 (inhibit cell division - benzamides, chloroacetamides) Dual II Magnum, Focus*, Frontier, Kerb, Primextra II Magnum*
Group 19 (inhibits auxin transport) Distinct*, OverDrive*
Group 22 (membrane rupture, photosynthetic inhibitors) Diquat, Gramoxone, Reward
Group 27 (HPPD inhibitors – isoxazole) Axial iPak*, Impact, Infinity*, Tundra*, Velocity m3*

*Products contain more than one active ingredient and appear in more than one group. In some instances, both active ingredients act to kill the same weed using different modes of action. In these instances, use of tank mixes may slow down the process of developing weed resistance.

New herbicides do not necessarily have a unique mode of action and may fall within the groups listed in the charts.

Herbicides that have the same mode of action may not control the same weed spectrum or have the same crop safety. For example, Assert and Ally have the same mode of action; however, Assert controls wild oats while Ally does not.

How Herbicides Work

After applying a herbicide, fields can be scouted to determine the effectiveness of the treatment. The symptoms of different herbicide groups, and the approximate time it takes to develop these symptoms, is listed in the table below. Weed patches that are not affected should be noted and checked, as they may be herbicide resistant. Note that symptoms may take longer to develop when conditions are not conducive to rapid plant growth.

The following table gives a brief description of symptoms that may be exhibited if plants are injured by a herbicide. The symptoms of each group are addressed for both foliar and soil exposures.

Table 2: The Mode of Action, Site of Uptake and Symptoms of Different Herbicide Groups.

Herbicide Group	Mode of Action	Site of Uptake	Weed symptoms/timing	
			Grass weeds	Broadleaf weeds
1	Systemic	Foliar	Reduced growth, yellowing of growing point in 1 to 3 weeks. Newest leaf of affected plant pulls out easily in 3 to 5 days.	Tolerant
2	Systemic	Foliar/Soil	Newest leaves yellowed in 3 to 10 days, dead in 1 to 3 weeks.	Discolored (red/yellow/purple) at the growing point and spreading to the whole plant in 1 to 3 weeks.
3	Systemic	Soil	Reduced emergence, poor root development of emerged plants. Roots often swollen/stunted and root tips darkened.	Reduced emergence, poor root development of emerged plants.
4	Systemic	Foliar	Tolerant to moderate rates. High rates cause symptoms similar to drought. Poor timing may cause kernal abortion.	Abnormal growth (twisted stems, cupped leaves) in 2 to 10 days.
5	Systemic	Soil	Wilted and yellowed oldest leaves, death in 7 to 10 days.	
	Contact	Foliar	Yellowed oldest leaves, death within days.	Bleached oldest leaves, death within days.
6	Contact	Foliar	Some leaf burn possible.	Yellowed leaves in 2 to 4 days, death in 1 to 2 weeks.
7	Systemic	Soil	Yellowed and stunted plants, death in 10 to 14 days.	
	Contact	Foliar	Interveneal yellowing of oldest leaves, death within days.	
8	Contact	Foliar	Yellowed leaves in 3 to 7 days, stunted plants.	Tolerant
	Systemic	Soil	Reduced emergence, emerged leaves dark green/blue.	
9	Systemic	Foliar	Wilted, yellowed leaves in 7 to 10 days. Newest growth is impacted first followed by the rest of the plant.	
10	Contact	Foliar	Wilted, bleached leaves in 3 to 5 days, death in 1 to 2 weeks.	
11	Systemic	Foliar	Plants wilted in 2 to 3 days, bleached and purpling leaves in 1 to 2 weeks.	Plants wilted in 2 to 3 days, bleached leaves in 1 to 2 weeks. Perennial plants die slowly.
14	Contact	Foliar	Some leaf burn at contact points or leaf edges.	Leaves yellowed and desiccated in 1 to 3 days. (Post applications)
	Systemic	Soil	Bleaching and yellowing, death prior to or shortly following emergence	
15	Systemic	Soil	Reduced emergence, emerged plants stunted.	
19	Systemic	Foliar	Twisting of older leaves, new leaves fail to expand, plant death in 2 to 4 weeks.	
22	Contact	Foliar	Leaves wilted within hours, desiccated in 1 to 3 days.	Leaves wilted in 1 to 3 days, desiccated and dead in 3 to 7 days.
27	Systemic	Foliar	Some bleaching and whitening of leaves.	Leaves bleached and whitened in 2 to 10 days and death in 7 to 10 days.

How to Identify Weed Resistance

It is important to avoid confusing herbicide failure caused by resistance with herbicide failure caused by various other factors (such as weather or application errors). When a herbicide fails to control weeds because of weather or application factors, that herbicide may work in the field the next season. But when herbicides fail because of the development of resistance, they will fail in subsequent years, regardless of weather or application procedures.

Herbicide resistance should be suspected under the following conditions:

- A weed species that the herbicide controlled in previous seasons now escapes the treatment, while other weeds that appear on the label continue to be controlled in the field.
- The escapes cannot be attributed to adverse weather or emergence after application (if a post-emergence product is in question).
- Irregular-shaped patches of a weed develop where the herbicide gives little or no control.
- Records of the past history of the field show repeated use of the same herbicide, or combinations of herbicides, that kill the weed in question in the same way.

Table 3: Herbicide-Resistant Weeds in Western Canada

WEED	HERBICIDE GROUP	LOCATIONS CONFIRMED
Canada Fleabane	Group 9	Occurs in several US states
Cleavers	Group 2	AB, SK
	Group 4	AB
	Multiple combinations of: Groups 2 & 4	AB
Chickweed	Group 2	AB, SK
Cow Cockle	Group 2	AB
Green Foxtail	Group 1	AB, MB, SK
	Group 2	Occurs in MB & Ontario
	Group 3	AB, MB, SK
	Multiple combinations of: Groups 1 & 3	MB, SK
Hemp-nettle	Group 2	MB
	Group 4	AB
	Multiple combinations of: Groups 2 & 4	AB
Kochia	Group 2	AB, MB, SK
	Group 4	Occurs in North Dakota and Montana
	Group 5	
	Group 9	AB, SK and occurs in North Dakota. Suspected in MB.

Lamb's-quarters	Group 2	Occurs in Ontario
	Group 5	Occurs in Ontario
Marshelder (False ragweed)	Group 2	Occurs in North Dakota
Mustard, Ball	Group 2	AB
Mustard, Wild	Group 2	AB, MB, SK
	Group 4	MB
	Group 5	MB
Narrow-leaved hawk's-beard	Group 2	AB
Persian Darnel	Group 1	SK
Redroot pig-weed	Group 2	Occurs in MB, Ontario and North Dakota
	Group 5	Occurs in Ontario
Russian thistle	Group 2	Occurs in SK, North Dakota and Montana
Shepherd's-purse	Group 2	AB, SK
Smartweed, pale	Group 2	MB
Spiny Annual Sow-thistle	Group 2	MB
Stinkweed	Group 2	AB
Wild buck-wheat	Group 2	AB
Wild oat	Group 1	AB, MB, SK
	Group 2	AB, MB, SK
	Group 8	AB, MB, SK
	Multiple combinations of Groups: 1 & 2, 1 & 8, 2 & 8, 1, 2 & 8	AB, MB, SK

See Table 1 on page 38 for a complete list of products in each Herbicide Resistance Group.

If Weed Resistance Develops on Your Farm

It is important to identify weed resistance before it spreads across your farm. Plan on conducting a "patch watch" scouting program this summer to identify suspicious patches before they become difficult to manage. Resistant weed patches have been identified on fields where producers were unaware of their existence.

Your patch watch program should begin shortly after spraying and continue through July after the crop has headed out and most weeds are visible from a distance. If you find suspicious looking patches, contact your

local agricultural office or crop protection company representative to assist you in confirming weed resistance. If resistance is suspected:

1. Map the location of the patches and mark them with stakes so you will remember their location.
2. Mow, cultivate or spot spray the patches. Resistant patches should not be allowed to produce seed.
3. Patchy areas should NOT be harvested with the rest of the field. Harvest these areas separately, and make sure to clean all harvesting equipment before leaving the area to prevent the spread of seeds across the field or to a neighbouring field.
4. Check patches each year to monitor their spread. Keeping your resistant weeds isolated to a manageable patch is easier than dealing with an entire field of resistant weeds.

Adjuvants and Registered herbicides:

Note – some products are specific about the concentration of active ingredient in the surfactant for product performance. Check with the herbicide page in this guide or the product label.

TRADE NAME	COMPOSITION	REGISTERED HERBICIDES
<i>Addit</i> (PCP# 29263)	37% Surfactant blend	<i>Bison</i>
<i>Adigor</i> Adjuvant	48.8% methylated rapeseed oil 28.2% ethoxylated alcohol	<i>Axial 100EC*</i>
<i>Agral 90</i> (PCP# 11809 or 24725),	90% nonylphenoxy polyethoxy ethanol	<i>Accent, Adrenalin SC, Altitude FX, Battalion, diquat, Escort, flucarbazone, glyphosate, Gramoxone, Muster, Pinnacle, Prism, imazethpyr, metsulfuron, thifensulfuron / tribenuron, Reflex, Reward, Triton K, Ultim</i>
<i>Agsurf II</i> (PCP# 30071, 30265, 30266 or 30267)	92% Alcohol ethoxylate	
<i>Amigo</i> (PCP# 22644), <i>X-Act</i> (PCP# 28225)	30% phosphate ester surfactant	<i>clethodim†*</i>
<i>Assist</i> (PCP# 16937), <i>XA oil Concentrate</i> (PCP# 11769), <i>Score</i> (PCP# 12200), <i>Signal</i> (PCP# 29173), <i>Steppe</i> (PCP# 29281)	83% paraffin based mineral oil 17% surfactant blend	<i>AAtrex, Basagran</i> (all crops), <i>Blazer</i> , <i>clodinafop†*</i> , <i>clodinafop + bromoxynil / MCPA ester†*</i> , <i>Harmony K*</i> , <i>Harmony SG*</i> , <i>Simplicity</i> , <i>Yuma GL</i> . (Not all adjuvants may be used with all herbicides listed)
<i>Citovett Plus</i> (PCP# 12766), <i>Super Spreader</i> (PCP# 17402)	50% Octylphenoxypolyethoxy ethanol	<i>Accent, AAtrex, Basagran</i> (peas), <i>Battalion, Escort, metsulfuron, Muster, Pinnacle, Prism, thifensulfuron / tribenuron, Triton K, Ultim</i>
<i>Companion</i> (PCP# 15882)	70% Octylphenoxypolyethoxy-(9)-ethanol	<i>Glyphosate, metsulfuron, Muster</i>
<i>Corn oil</i> (PCP# 18473)	99% paraffin based mineral oil 1% surfactant blend	<i>AAtrex, linuron</i>
<i>Enhance</i> (PCP# 29270 or 29952), <i>MANA Adjuvant 80</i> (PCP# 30419)	80% triglyceride ethoxylate 10 POE	<i>Accent, Battalion, diquat, Escort, glyphosate, Gramoxone, Lontrel, Muster, Pinnacle, Prism, imazethapyr, metsulfuron, thifensulfuron / tribenuron, Reflex, Reward, Ultim</i>
<i>Hasten</i> (PCP# 27420)	77.4% methyl and ethyl oleate(esterified vegetable oil)	<i>Option 35DF</i>
<i>LI700</i> (PCP# 23026)	80% surfactant blend	<i>Diquat, flucarbazone, glyphosate, Fulfill insecticide</i>
<i>Liberate</i> (PCP# 29491)	100% lecithin, methyl esters of fatty acids and alcohol ethoxylate	<i>Battalion, Pursuit, Reflex, Odyssey, Everest, thifensulfuron / tribenuron, tribenuron, metsulfuron (Accurate)</i>
<i>Merge</i> (PCP# 24702)	50% solvent 50% surfactant blend	<i>Ares, Equinox, Odyssey, Odyssey DLX, Poast Ultra, quizalofop, Solo, Tensile, Triton C</i>

Continued on next page

TRADE NAME	COMPOSITION	REGISTERED HERBICIDES
<i>Sure-Mix</i> (PCP# 25467)	60% paraffin based mineral oil 40% surfactant blend	quizalofop
<i>Turbocharge</i> (PCP# 23135)	50% mineral oil 39.5% surfactant blend	<i>Achieve Liquid Gold</i> *, tralkoxydim [†] *

* The adjuvant is packaged with the product.

[†] Note: All products may not be registered with all adjuvants. See product page in the following sections to determine which adjuvants are registered for each herbicide.

Adjuvants and Your Herbicide

Adjuvants are important ingredients in chemical weed control. Many herbicides must be applied with an adjuvant. If it is forgotten, the level of weed control can vary widely, and re-spraying may be necessary.

Most products have adjuvants built into the formulation. Other require adjuvant addition (e.g. Refine SG). Some adjuvants were developed specifically for one herbicide, and these are either pre-packaged with the herbicide, or are identified by name on the label (e.g. Turbocharge for Achieve, Amigo for Select).

With some products, adjuvants need to be added only under certain conditions. For example, glyphosate products have built-in adjuvants, but require additional adjuvant when low rates (pre-seeding or chem-fallow), high water volumes, or certain tank mixes are used.

Adjuvants should be added only when required. If one is not required, addition can reduce weed control or injure crops. Product labels will describe when an adjuvant is required, and what type should be used.

There are two main classes of adjuvants: "activators or spray modifiers" (these include surfactants and crop oils), and "utility modifiers" (these include pH adjusters, water conditioners, low-drift adjuvants, and anti-foaming agents). The most important class of adjuvants is the activators. Surfactants, the main group within the activators, are "surface active agents." These chemicals produce effects at points where two substances touch, such as between two liquids (herbicide and water) or between a solid and a liquid (herbicide and leaf surface). Some surfactants act as dispersing agents, helping to keep a pesticide suspended in water. Others work on the plant, improving the wetting, sticking and penetrating characteristics of the herbicide droplets. Oil-based adjuvants contain petroleum or vegetable oil plus an emulsifier that suspends the oil in tiny droplets within the spray solution. Oil-based adjuvants typically assist in herbicide penetration into the leaf.

There are two basic types of surfactants (ionic and non-ionic), of which the non-ionic are most common. The tables on the previous page list the surfactants registered for use with herbicides in Western Canada.

Crop and Herbicide Recommendation Tables

The ratings in the following charts give general comparisons based on rates, timing and other application instructions and precautions as outlined in this Guide.

The ratings for crop tolerance and weed control are explained below. Where ratings are not given, a dot (•) will indicate if the weed is listed on a product label. Where rate ranges are listed for controlling a given weed, ratings are based on results achieved with the higher rate.

RATING	CROP TOLERANCE	WEED CONTROL
E - Excellent	Consistently exhibits a high degree of tolerance over a wide range of growing conditions.	Consistently provides a high degree of control over a wide range of growing conditions.
G - Good	Exhibits good crop tolerance under most growing conditions.	Provides good control under most growing conditions.
F - Fair	Tolerance is acceptable depending on growing conditions.	Control is acceptable depending on growing conditions.
P - Poor		Provides suppression, but control is often unacceptable.

All ratings are primarily based on the use of the herbicides on their own. Tank mixing herbicides may reduce the control of one or both tank mix partners, or result in a reduced level of crop safety.

Weed Control Tables

Table 1. Weed Control in Barley

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Stinkweed	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canada	Volunteer Sunflowers
2,4-D	66																											
Achieve Liquid Gold	74	• ³	•			•	•	•			•	• ⁵	•		•	•		•	•	•	•	•		•	• ¹⁰		•	•
Avadex	88					•																						
Avenge	92					•																						
Axial	93	•	•			•																						
Axial Xtreme	96	•	•			•	S		•						•													
Barricade II	97						•	•	•				•	•								•		•	•	•	•	
Bromoxynil	108						•				•				• ³			• ⁷	• ⁷	•		•		• ⁷				
Bromoxynil/2,4-D	112						•	•			•		•		•			•	•	•	•	•				•	•	
Bromoxynil/MCPA	113						•	•			•		•		•			•	•	•	•	•	• ¹⁰		• ¹⁰		•	•
Curtail M	125						•				•	• ³	•		S			•	• ⁷		• ⁷	• ⁷	• ¹⁰	•	•	• ⁷	•	
Dicamba + MCPA/2,4-D	126						•			•	•		•	• ³	•			•	•	•	•	•	• ¹⁰		• ¹⁰		•	•
Dicamba/Mecoprop/MCPA	130						•	•	•				•	• ³	•			•	•	•	•	•		•	• ¹⁰		•	•
Dichlorprop/2,4-D	132						•	• ³			•		•		•		•	•	•	•	•	•	•				•	•
DyVel	139						•			S	•		•	• ³	•			•	•	•	•	•	•		•		• ³	•
DyVel DSp	141									S	•		•				S	•	•	•	•	•			• ¹⁰		•	
Enforcer D	145						• ⁷								•				• ⁷	• ⁷							•	
Enforcer M	147						• ⁷		• ⁷	• ⁷					•	•			S								•	
Fenoxaprop	156	•	•			•																						
Florasulam + MCPA	162						•		•	•		• ⁵	•	•				•	•		•	•	• ¹⁰	•	• ¹⁰		•	•
Fluroxypyr + 2,4-D	168						•	S	•	•	• ³	•	S	•	•	•	•	•	•	•	•	•	S	•	S	•	•	•
Fortress	172		•			•	S								S	S			S	S								
Imazamethabenz	198					•	S											•										
Infinity	205							•	•			• ⁴	•	•	•	•	S	•	•	•	•	•	S	•	S		•	
Linuron + MCPA amine	213		S				•				•		•	•	•			•	•		•	•						
Lontrel 360	216						•																•		•			
MCPA	218										•	• ¹⁰	•		•			•			•			•				
MCPA K	218										•		•					•			•			•				
Mecoprop-p	223								•	•								•										
Metribuzin	224							•	•					• ⁷				•	•	•		•		•			•	
Metsulfuron	227								•				•	•			•	•	•	•	•	•	• ³				•	
Momentum	230						•			•					•											•	•	
Prestige XC	245							S				• ³	•	S	•		•	•	•	•	•	•	•	•	•	•	•	
Spectrum	266						•	•	•				• ³	•				•	•	•	•	•	•	•	•	•	•	
Thifensulfuron/tribenuron	272						•	•					•	•				•	•	•	•	•	•				• ³	•
Thifensulfuron/tribenuron + MCPA	276						•	•				• ⁶	•	•	•	•		•	•	•	•	•	•	•			•	•
Tralkoxydim	279	•	•			•																						
Trifluralin (foxtail control)	286		•																									
Trifluralin (grassy and broadleaf)	286	•	•			•	•	•									•		•									
Triton C	291						•	•	•				•	•	•	•	•	•	•	•	•	•	•			• ³	•	
Triton K	293						•						•	•	•	•	•	•	•	•	•	•				•	•	
Trophy	295						S		•	•		•	• ³	•	•			•	•		•	S		•		•	•	
Tundra	296	•	•			•	•	•	•			• ⁴	•	•	•	•	S	•	•	•	•	•	S	•	S		•	•

¹ MCPA K mixes only. ² Will not control CLEARFIELD canola varieties. ³ Spring seedlings only.

⁴ Up to 25 cm diameter. ⁵ Seedlings and overwintered rosettes. ⁶ Less than 15 cm diameter ⁷ Controlled at higher rates ¹⁰ Top growth control only.

• Control. S - Suppression.

Table 2. Weed Control in Oat

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow thistle (Perennial)	Stinkweed	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
Bromoxynil	108					•				•				• ³	•		• ⁶	• ⁶	• ³		•		• ⁶				
Bromoxynil/MCPA	113					•	•			•		•		•	•		•	•	•	•	•	• ⁷	•	• ⁷		•	•
Curtail M	125					•				•	• ³			• ⁵			•	• ⁶		•	• ⁶		•		• ⁶		
Dicamba + MCPA	126					•		•	•	•			• ⁵	•	•		•	•	•	•	•	• ⁷	•	• ⁷		•	
Dicamba/Mecoprop/MCPA	130					•	•		•				• ³	•	•		•	•	•	•	•	•	•	• ⁷		•	•
DyVel	139					•			•	•			• ³	•	•		•	•	•	•	•	•	•		• ⁶	•	
Florasulam + MCPA	162					•		•			• ⁵	•	•	•	•		•	•		•	•	• ⁷	•	• ⁷		•	•
Linuron + MCPA amine	213	S				•	•	•		•		•	•		•		•	•		•	•	•	•				
Lontrel 360	216					• ⁶																• ⁷		•			
MCPA	218									•		•					•			•		•					
MCPA K	218									•		•	• ⁵	•	•		•			•			•				
Mecoprop-p	223							•	•						•		•							• ⁷			
Prestige XC	245					•		• ⁵	•		• ³	•	• ⁵	•	•	•	•	•		•	•	•	•	•	•	•	
Spectrum	266					•		•			• ⁵	• ³	•		•		•	•		•	•	•	•	•	•	•	
Thifensulfuron/tribenuron	272					•		•				•	•		•		•	•	•	•	•	•	•		• ⁶	•	
Thifensulfuron/tribenuron + MCPA	276					•		•			•	•	•		•	•	•	•	•	•	•	• ⁷	•	• ⁷		•	•

¹ MCPA K mixes only. ² Will not control CLEARFIELD canola varieties. ³ Spring seedlings only. ⁵ Seedlings and overwintered rosettes. ⁶ Controlled at higher rates.

⁷ Top growth control only. S - Suppression. • Registered for control.

Table 3. Weed Control in Rye or Triticale

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Stinkweed	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
2,4-D ^{3,4}	66										•	•			•	•		•	•	•	•	• ¹⁰	•			•	•	
Achieve Liquid Gold ^{1,4}	74	•	•			•	•	•			•	•	•		•	•		•	•	•	•	• ¹⁰	•	• ¹⁰		•	•	
Avenge ¹	92					•																						
Bromoxynil ¹	108						•				•				• ⁹	•		• ⁹	• ⁹	•		•		• ⁹				
Bromoxynil/MCPA ^{1,4}	113						•	•			•		•		•	•		•	•	•	•	• ¹⁰	•	• ¹⁰		•	•	
Dicamba + 2,4-D ^{2,4}	126					•			•	•	•		•		•	•		•	•	•	•	• ¹⁰	•	• ¹⁰		•	•	
Infinity ⁵	205					•		•	•		• ⁶		•	•	•	•	S	•	•	•	•	S	•	S		•		
MCPA ^{3,4}	218										•		•		•	•		•		•				•				
Tralkoxydim ³	279	•	•			•																						

¹ Fall rye only. ² Spring rye only. ³ Fall and Spring rye. ⁴ Rye only; not registered for triticale. ⁵ Triticale only. ⁶ Up to 25 cm diameter.

⁸ Seedling stage only. ⁹ Controlled at higher rates. ¹⁰ Top growth control only.

• Registered for control. S - Suppression.

Table 4. Weed Control in Wheat (Spring and Winter)

HERBICIDE	PAGE	Crop	W-Winter, S-Spring	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Stinkweed	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers
2,4-D	66	W/S									•		•	•			•	•			•	•	•	•			•		•	•
Achieve Liquid Gold	74	W/S	• ⁶	•				•	•	•			•	•	•		•	•			•	•	•	•			•		•	•
Adrenaline SC ¹¹	75	S ¹¹	•	•				•	•	•		•	•	•		•	•	•			•	•	•	•	•		•		•	•
Altitude FX ¹¹	78	S ¹¹	•	•				•	•	•		•	•	•		•	•	•			•	•	•	•			•		•	•
Altitude FX2 ¹¹	80	S ¹¹	•	•	•			•	•	•							•	•	•	•	•	•	•	•			•		•	•
Avadex	88	S						•																						
Avenge ^{7,8}	92	W/S						•																						
Axial ⁷	93	S		•				•																						
Axial Xtreme	96	S	•	•				•	•			•					•			•										
Barricade II	97	S							•	•		•								•										
Basagran + 2,4-D ⁷	99	S											•					•			•	•	•	•			•		•	•
Bromoxynil	108	W/S						•					•				• ⁶	•		• ⁶	• ⁶	•	•	•		• ¹¹			•	•
Bromoxynil/2,4-D	112	S						•	•				•				•	•		•	•	•	•	•			• ¹¹		•	•
Bromoxynil/MCPA	113	W/S						•	•				•				•	•		•	•	•	•	•	• ¹¹	• ¹¹			•	•
Clodinafop	122	S	•	•				•													•	•	•	•						
Clodinafop + Bromoxynil/MCPA	122	S	•	•				•	•	•			•				•	•		•	•	•	•	•	• ¹¹	• ¹¹			•	•
Curtail M	125	S						•					•	• ⁶	•		•			• ¹¹		•	•	•	• ¹¹	• ¹¹			• ¹¹	•
Dicamba + MCPA/2,4-D	126	W/S						•					•			• ⁶	•	•		•	•	•	•	•	• ¹¹	• ¹¹			•	•
Dicamba/Mecoprop/MCPA	130	W/S						•	•				•			• ⁶	•	•		•	•	•	•	•	• ¹¹	• ¹¹			•	•
Dichlorprop/2,4-D	132	W/S						•	• ⁶				•			•	•	•		• ⁶	•	•	•	•	• ¹¹	• ¹¹			•	•
DyVel	139	W/S						•			•	•	•			• ⁶	•	•		•	•	•	•	•	• ¹¹	• ¹¹	• ¹¹	• ¹¹	• ⁶	•

Continued on next page.

Table 4. Weed Control in Wheat (Spring and Winter) *continued*.

HERBICIDE	PAGE	Crop	W-Winter	S-Spring	Barley	Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Corn	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Stinkweed	Thistle, Canada	Volunteer Flax	Volunteer Mustard, Canola	Volunteer Sunflowers		
DyVel DSp	141	W/S								•				S	•				•	•	S			•	•		•	• ¹⁴		• ¹⁴		•	•	
Enforcer D	145	S								• ⁹										•	•			• ⁹	• ⁹							•		
Enforcer M	147	W/S								• ⁹		• ⁹	• ⁹						•	•			S									•		
Fenoxaprop	156	S			•	•			•														S										•	
Florasulam + MCPA	162	S								•		•	•			• ²	•	•		•			•		•	•	•	• ¹⁴		• ¹⁴		•	•	
Flucarbazone	164	S				• ¹			•													•	•		•	•						• ⁵		
Fluroxypyr + 2,4-D	168	W/S								•		S	•	•	•	• ⁶	•	S	•	•	•	•	•	•		•	•	S	•	S	•	•	•	
Fortress	172	S			•				•										S	S			S	S		•	•							
Frontline + 2,4-D	176	S								•		•	•	•	•	• ⁷	•	S	•	• ¹³		•	•	•	•	•	•	S	•	S		•	•	
Harmony K	193	S		•	•				•	•	•	•							• ¹³	•		•	•	•	•	•	•	•				• ⁵	•	
Harmony SG	195	S		•	•				•	•	•	•								•		•	•	•	•	•	•	•				• ⁵	•	
Imazamethabenz	198	S							•													•		•	•	•	•	•					• ⁵	•
Infinity	205	W/S								•	•	•				S	•	•	•	•	•	S	•	•	•	•	•	S	•	S		•		
Linuron + MCPA amine	213	S								•	•			•	•		•	•	•	•		•	•		•	•	•	•						
Lontrel 360	216	S								•																	• ¹⁴		•					
MCPA	218	W/S													•		•	S		•		•			•				•					
Mecoprop-p	223	S										•	•	•												•								
Metribuzin	224	S									•	•								•		•	•	•	•	•	•					•		
Metsulfuron	227	S										•	•							•	•	•	•	•	•	•	•	• ¹³	•	• ¹³		•	•	
Momentum	230	S								•				•						•						•	•	•					•	
Prestige XC	245	W/S								•		S	•	•		• ⁶	•	S	•	•	•	•	•		•	•	•	•	•	•	•	•	•	•
Simplicity	263	W/S	•	S ¹					•	S	•	•	•		S				•	•		•		•	S	•	•	•	•	S		• ⁵		
Spectrum	266	S								•	•	•	•	•		• ²	• ⁶	•	•	•	•		•	•	•	•	•	•	•	•		•	•	
Thifensulfuron/tribenuron	272	W/S								•	•						•	•	•	•	•	•	•	•	•	•	•	•				• ⁵	•	
Thifensulfuron/tribenuron + MCPA	276	W/S								•	•					• ¹⁵	•	•	•	•		•	•	•	•	•	•	•				•	•	
Tralkoxydim	279	W/S	•	•					•																									
Trifluralin (foxtail control)	286	S		•																														
Triton C	291	S								•	•	•							•	•	•	• ⁶	•	•	•	•	•	•					• ⁵	•
Triton K	293	S								•									•	•	•	•	•	•	•	•	•					•	•	
Trophy	295	S								S			•	•	•				•	•	•	•	•	•	•	•	S	•	•		•	•	•	
Tundra	296	S	•	•					•	•	•	•	•			S	•	•	•	•	•	S	•	•	•	•	•	S	•	S		•	•	
Varro	300	W/S	•	S					•	•	•	•	•	•	•			•	•	S	S	•	•	S	•	•						S		

¹ Green foxtail only. ² Spring seedlings and overwintered rosettes. ³ MCPA K mixes only. ⁵ Will not control CLEARFIELD canola varieties. ⁶ Spring seedlings only.⁷ Not registered for durum wheat. ⁸ Not recommended for all spring wheat varieties. Check product listing for details. ⁹ Controlled at the higher rates.¹⁰ Tank mix with 2,4-D ester when applying to durum wheat. ¹¹ For use on CLEARFIELD wheat varieties only. ¹² Up to 25 cm diameter. ¹³ Less than 15 cm diameter.¹⁴ Top growth only.

• Registered for control. S - Suppression

Table 5. Weed Control in Corn

HERBICIDE	PAGE	Barnyard Grass	Volunteer Cereals	Foxtail, Green	Foxtail, Yellow	Wild Oats	Quackgrass	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed Annual Species, Lady's-thumb	Sow-thistle (Perennial)	Stinkweed	Sunflower, Volunteer	Thistle, Canada	Volunteer Canola
2,4-D	66								P		G	P	E	G	E	E		•	G	E	F	P	E	P	P	E
Accent	72	•		•	S	•	•																			
Atrazine	70					•		•								•	•	•			•					
Basagran/Basagran Forté	99								•	•	•					•	•	S	S	•	•		•		S	•
Battalion ⁵	102	•		•	•		S	•								•	•	•			•					
Bromoxynil	108						G				F			F	G	F	F	G		G		F				
Bromoxynil/MCPA	113						•	•			•		•	•	•	•	•	•	•	•	S	•	•	S	•	
Dicamba	126						•			•				•	•	•	•	•		•	S			S		
Dicamba + 2,4-D amine	126						•			•	•	S	•	•	•	•	•	•	•	•	S	•	•	•	S	•
Dual II Magnum	137	•		•	•													S								
DyVel DSp	141						G			S	G		E	G	E	E	G	G	E	G	F	E		F	G	
Focus ⁸	171	• ⁹		• ⁹	• ⁹						•			•	•		•					•				•
Frontier Max	174			•																						
Glyphosate ^{1,4}	178	E	E	E	E	E	F	G	G	E	E	•	F	E	E	E	E	E	E	E	E	F	E	E	F	• ⁷
Impact ¹⁰	202	S		S	S				S							S	•	S			S					
Impact + Atrazine	202	S		S	S				S							•	•	•			•					
Liberty 200 SN ³	212	•		•	•	•	S	•	•		•					•	•	•		•	•	•	•		S	
Linuron	213	S	S	S	S			•	•			•		•	•	•	•	•		•	•	•	•			
MCPB/MCPA	221															•	•	•		•		S	•		S	•
Option 35 DF/Option 225 OD ⁶	238	•		•	•				•							•	•	•								
Primextra II Magnum	247	•		•	•			•								•	•	•			•					
Ultim ^{2,6}	298	•	•	•	•	•	•											•								

¹ For use on Roundup Ready varieties only. ² See product page for registered corn varieties. ³ For use on Liberty 200 SN tolerant corn varieties only.

⁴ Not all glyphosate products are registered for use on Roundup Ready Corn. ⁵ Registered for use in Red River Valley region of Manitoba ONLY.

⁶ For use in Manitoba only. ⁷ Except glyphosate tolerant varieties. ⁸ Apply prior to seeding or up to 3 days after seeding.

⁹ Control of the following weeds emerging from seed (not controlled if emerged at application). ¹⁰ Must be applied with a tank mix partner.

• Controlled S - Suppression

Table 6. Weed Control in Pea

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green and Yellow	Quackgrass	Volunteer Barley	Volunteer Wheat	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's-purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Sunkweed	Thistle, Canada
Authority/Authority Charge ³	86							G								E	E		E						
Avadex	88						G																		
Basagran/Basagran Forté	99									F	F	F							S	F	F	F		F	P
Clethodim	119	E	E	F	E	E	G								S	G	E		G	S		S			
Edge Granular	144	E	E		S	S	S	G		G	S														
Equinox	150	• ¹	•	•	•	•																			
Imazethapyr	200	E ¹					F	S		G	G				G			E	G		G	G		E	
MCPA Sodium Salt/Amine	218													P			P	G			P		P	P	
MCPB/MCPA	221														F		G	G	G		G		P	G	P
Metribuzin	224									G					F		G	G				G		G	
Odyssey	234	G	E ¹		G	G ²	G	S		E	E			E	S	S	S	E	E	E	E	E	G	E	
Odyssey DLX	235	•	• ¹		•	•	•	S		E	E			E	S		S	E	E	E	E	G		E	
Poast Ultra	242	E	E	F	E	E	G																		
Quizalofop	252	G	E	F	E	E	G											G		G	G				
Trifluralin (broadleaf & grassy weeds)	286	G	G				F	F		G															
Viper ADV	305	•	•		•	•	•	S			S						•	•	•		•	•		•	

¹ Green foxtail only. ² Excluding CLEARFIELD varieties. ³ For in season activity only. For initial burn down of other weeds see table 13b.

• Registered for control. S - Suppression.

Table 7. Weed Control in Other Pulses

HERBICIDE	CROP							ANNUAL WEEDS																	PERENNIALS							
	PAGE	Bean, Dry	Faba bean	Lentil	Chickpea	Soybean	Sweet White Lupin	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Volunteer Barley	Volunteer Wheat	Wild Oat	Buckwheat, Wild	Chickweed	Cleavers	Cocklebur	Hemp-nettle	Kochia	Lamb's-quarters	Mustard, Wild	Nightshade, Hairy	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Stinkweed	Canada Thistle	Dandelion	Perennial Sow-thistle	Quackgrass	
Authority/Authority Charge ¹⁹	86				X									G					E	E		E										
Basagran/Basagran Forte	99	X ⁶	X			X									F	F	F			F	F	F	S	F	F	F	F	F	P			
Blazer	105					X											•			•	•		•			•		S				
Clethodim	119	X		X	X	X		E	E	E	E	E	G																			
Dual II Magnum	137	X ⁷				X	X	G	G	G													P									
Edge Granular	144	X ¹	X	X ⁸		X		E	E	E	S	S	S	G	G	S		S	G	E		G	S		S							
Eptam 8-E	148	X						G	G	G	F	F	G		G				G			G										
Equinox	150			X ¹⁰					•		•	•	•																		•	
Flexstar ⁴	159					X		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Focus ¹⁷	171					X		• ¹⁸	• ¹⁸	• ¹⁸							•		•	•			•			•						
Frontier Max	174	X ¹							•																							
Glyphosate ^{11, 12}	178				X ¹¹			E	E	E	E	E	E	G	E	E	•	E	E	E	E		E	E	E	E	E	E	F	F	F	F
Imazethapyr	200	X ¹			X																	G										
Linuron	213				X	X		S	S	S				•	•								•		•	•	•					
Metribuzin (post-emergence)	224			X	X										F			F		F	F					F	F					
Metribuzin + Trellan (PTI)	224		X			X		G	G	G			F	F	G			F		G	G	G	G	G	G	G	G					
Odyssey	234			X ⁹		X		G	G		G	G ¹⁵	G	F ¹³	E	E		F ¹³	S	E		E	E	E ¹³	E	G	E					
Odyssey DLX	235			X ⁹				•	•		•	•	•	S	E	E		S	S	E		E	E	E	E	G	E					
Pinnacle	241					X														•	•		•			•						
Poast Ultra	242	X	X	X	X	X	X	E	E	E	E	E	G																		F	
Quizalofop	252	X ⁶		X	X	X		G	E	•	E	E	G																		F	
Reflex + Basagran	256	X ⁶															•			•	•	S		•	•	•						
Solo	265			X ⁹				•	•	•	•	• ¹⁵	•	S		S			•	•		•		•	•	•	•					
Trifluralin (broadleaf & grassy weeds)	286	X	X	X		X		G	G	G			F	F	G				G			G	G									
Viper ADV	305	X				X		•	•	•	•	•	•		•				•	•		S	•	•	•	•						

¹ White and kidney beans only. ² Pre-emergent surface treatments only. ³ Pinto, pink and red beans only. (Refer to Imazethapyr section for full list of weeds controlled).

⁴ For glyphosate tolerant soybeans in the Red River Valley of Manitoba. ⁵ For use on navy beans in the Red River Valley of Manitoba. Does not include weeds controlled by Basagran Forté. ⁶ Not all dry bean types have been tested for tolerance to this herbicide. ⁷ White, kidney and pinto beans only. ⁸ Fall applications only.

⁹ For use ONLY on CLEARFIELD lentil varieties. ¹⁰ Including Clearfield lentil varieties. ¹¹ For use on glyphosate tolerant varieties only.

¹² Not all glyphosate products are registered for use on glyphosate tolerant soybeans. ¹³ Suppression in CLEARFIELD lentils.

¹⁴ Not controlled in CLEARFIELD lentils. ¹⁵ Not including CLEARFIELD wheat varieties. ¹⁷ Apply prior to seeding of or up to 3 days after seeding.

¹⁸ Control of the following weeds emerging from seed (not controlled if emerged at application).

¹⁹ For in season activity only. For initial burn down of other weeds see table 13b.

• Registered for control. S - Suppression.

Table 8. Weed Control in Flax and Solin (Low Linolenic Acid Varieties).
Not all products are registered for use on Solin.

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Quackgrass	Volunteer Cereals	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Stinkweed	Thistle, Canada
Authority/Authority Charge ⁵	86							G								E	E			E						
Avadex	88						G																			
Basagran ¹ /Basagran Forté	99								F	F	F						F	F	F	P	F	F	F	F	F	F
Bromoxynil ¹	108							G				F				F	G		F	F	G		G		F	
Bromoxynil/MCPA ¹	113							E	F			G		G		G	E		E	S	G	G	F	P	G	P
Clethodim ¹	119	E	E	E	F	E	G																			
Curtail M ¹	125							F				•	C ²	E			E		E	P		E	P	G	E	G
Eptam ³ 8-E	148	G	G	G		F	G			G							G			G						
Equinox ¹	150		•		•	•	•																			
Fortress	172		G	G			G	P								P	P			P	P					
Lontrel 360 ¹	216							G																F		G
MCPA/MCPA K	221											G	P	E	F		E		E			E		P	E	P
Poast Ultra ¹	242	E	E	E	F	E	G																			
Quizalofop ¹	252	G	E	•	F	E	G																			
Trifluralin (broadleaf and grassy weeds) ⁴	286	G	G	G			F	E		G							G			G	G					

• Registered for control.

¹ Registered for use on Solin (low linolenic acid varieties). ² Spring seedlings only. ³ Not recommended for use on flax in Saskatchewan. ⁴ Fall application only.

⁵ For in season activity only. For initial burn down of other weeds see table 13b.

Table 9. Weed Control in Canola

HERBICIDE	PAGE	Barnyard Grass	Foxtail, Green	Foxtail, Yellow	Quackgrass	Volunteer Barley	Volunteer Wheat	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Sow-thistle (Perennial)	Stinkweed	Thistle, Canada
Ares ²	83	•	•	•		•	•	•	•			•				•		•	•	•	•	•	•	•	•	•	
Avadex	88							G																			
Clethodim	119	E	E	E	F	E	E	G																			
Eclipse ⁴	143		E		F	E	E	E	G	G	E	E		F		E	E	E		E	E	E	E	•	G ⁷	E	G ⁷
Edge Granular	144	E	E	E		S	S	S	G		G	S				S	G	E			E	S		S			
Equinox	150		•		•	•	•	•																			
Fortress	172		G	G				G	P								P	P			P	P					
Glyphosate ^{4,5}	178	E	E	E	F	E	E	E	F	G	E	E		F	E	E	E	E	G	E	E	E	E	E	F	E	F
Liberty ¹	209	G	G		F	F ³	G ³	G	G		G	G		F	G	G	G	G	G	G	G	G	G	G	G	G	F
Lontrel 360	216							G																	G		G
Muster Toss-N-Go	232														G	G				G	P			G		F	
Odyssey ²	234	G	E			G	G ⁵	G	F		E	E			E	G		S		E	E	E	E	G		E	
Poast Ultra	242	E	E	E	F	E	E	G																			
Quizalofop	252	G	E	•	F	E	E	G																			
Solo ²	265	•	•	•		•	• ⁶	•	S			S						•		•	•		•	•		•	
Tensile ²	271	•	•	•		•	•	•	•			S						•		•	•		•	•	• ⁹	•	• ⁹
Trifluralin	286	G	G	G				F	F		G						G				G	G					

¹ For use **only** on Liberty Link varieties. ² For use **only** on CLEARFIELD canola varieties. ³ Ratings based on 1.35 L./acre rate of Liberty. Control may be reduced at lower rates. ⁴ For use **only** on glyphosate tolerant canola varieties. ⁵ Will not control CLEARFIELD wheat volunteers. ⁶ Season long control. ⁷ Top growth control only.

• Registered for control. S—Suppression.

Table 10. Weed Control in Potatoes*

HERBICIDE	PAGE	Barley Grass	Footail, Green and Yellow	Volunteer Canola	Volunteer Corn	Volunteer Barley	Volunteer Wheat	Wild Oats	Quackgrass	Chickweed	Dandelion	Hemp-nettle	Lamb's-quarters	Mustard, Wild	Nightshade	Pigweed, Redroot	Pigweed, Prostrate	Purslane	Smartweed (Annual)	Shepherd's-purse	Sunkweed
Chateau	166																				
Clethodim	119	•	•		•	•	•	•					•		•	•	•				
Diquat	134																				
Dual II Magnum	137	•	•																		
Eptam 8-E	148	•	•												2	S					
Glyphosate (preplant use only)	178			• ¹	• ¹	•	•	•	S	•			•		3	•	•	•			
Gramoxone (pre-emergent use only)	190	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•		•	•	•
Linuron (pre-emergent use only)	213	S	•										•	•	•	•	•	•	•	•	•
Metribuzin ³	224			•						•			•			•	•	•	•	•	•
Poast Ultra	242	•	•		•	•	•	•	S			•	•	•		•			•	•	•
Prism	248	•	•					•					S			•					

• Controlled S - Suppression

¹ Will not control glyphosate tolerant varieties. ² American and Eastern black nightshades. ³ Hairy nightshade.⁴ Consult manufacturer or seed provider for varietal tolerance to Metribuzin.**Note:** Before using any pesticides on potatoes, consult the list of Agricultural Pesticides Approved for Use, available from Simplot Canada and McCain Foods (Canada).

Table 11. Weed Control in Sunflowers

HERBICIDE	PAGE	Barley Grass	Footail, Green and Yellow	Quackgrass	Volunteer Barley	Volunteer Wheat	Wild Oats	Buckwheat, Wild	Catchfly, Night-flowering	Chickweed	Cleavers	Cocklebur	Dandelion	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mallow, Round-leaved	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Smartweed, Annual Species	Sow-thistle (Perennial)	Sunkweed	Thistle, Canada
Authority / Authority Charge ⁴	86							G																	
Clethodim	119	E	E	F	E	E	G									E	E			E					
Edge Granular	144	E	E		S	S	S	G		G	S														
Eptam 8-E	148	G	G		F	F	G			G					S	G	E			E	S	S			
Imazamethabenz	198																G			G					
Muster Toss-N-Go	232																		G					G	
Poast Ultra	242	E	E	S	E	E	G						G	G					G			G		F	
Solo ^{2,3}	265	•	•		•	• ¹	•	S		S							•								
Trifluralin	286	G	G				F	F		G						G				G	G		•		

• Registered for control. S - Suppression.

¹ Will not control CLEARFIELD volunteers. ³ Apply only on CLEARFIELD sunflower varieties.⁴ For in season activity only. For initial burn down of other weeds see table 13b.

Table 12. Weed Control in Special Crops

HERBICIDE	PAGE	CROP							ANNUAL WEEDS																	PERENNIALS										
		Canaryseed	Safflower	Caraway	Coriander	Buckwheat	Mustard	Cheesed mustard (<i>Braister juncea</i>)	Barley Grass	Footail, Green	Footail, Yellow	Volunteer Barley	Volunteer Wheat	Wild Oat	Buckwheat, Wild	Catchfly, night-flowering	Chickweed	Cleavers	Cocklebur	Flaxweed	Hemp-nettle	Kochia	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Russian Thistle	Shepherd's Purse	Smartweed, Annual Species	Stinkweed	Volunteer Flax	Volunteer Mustard, Canola	Canada Thistle	Dandelion	Perennial Sow-thistle	Quackgrass	
Avadex	88	✓ ¹					✓							G																						
Avenge	92	✓												G																						
Bromoxynil	108	✓																	F			F	G	F	F	G			G	F						
Bromoxynil/MCPA	113	✓													E	F			G	G		G	E	E	G	G	G	G	G	G		G	P		P	
Clethodim	119				✓		✓		E	E	E	E	E	G																					F	
Curtail M	125	✓													F					•	E		S	E	E	P		E	P	E		F	G	G ⁴	G	
Dicamba + MCPA	126	✓													G			G	G	E	F		E	E	G	G	E	G	E	E	P	G	F	P	F	
Dicamba/Mecoprop/MCPA	130	✓													G	G			G	E	F	G	E	E	G	G	E	G	E	E		G	F		F	
Edge Granular	144		✓	✓	✓		✓ ¹		E	E	E	S	S	S	G		G	S			S	G	E		G	S		S								
Fortress	172						✓			G	G				G	P						P	P		P	P										
Linuron	213			✓	✓																															
Muster Toss-n-Go	232						✓ ³												G	G			G	P			G	F								
Odyssey	234						✓ ⁴	•	•		•	• ⁸	•	•			•	•		•	•		•	•	•	•	•	•	•	•	•	•	• ⁸			•
Odyssey DLX	235						✓ ⁴	•	•		•	•	•	•			•	•		•	•		•	•	•	•	•	•	•	•	•	•	• ⁸			•
Poast Ultra	242		✓	✓	✓	✓	✓		E	E	E	E	E	G																					F	
Prestige XC	245	✓													E		S	E		E	S	E	E	E	G		E	G	E	E	E	E	E	G ⁴	E	
Quizalofop	252						✓	✓	G	E	•		E	G									•	•	•			•	•	•						
Solo	265						✓	✓	•	•	•	•	• ⁸	•	S		S						•	•	•			•	•	•		• ⁸				
Trifluralin	286		✓				✓		G	G	G				F	F	G						G		G	G										
Trophy	295	✓													S				•	•	•	•	•	•				•	S	•	•	•				

¹ Granular formulation only. ² Yellow mustard only. ³ Brown and oriental mustards only. ⁴ Spring seedlings only. ⁵ Oriental mustard only. ⁶ For use in CLEARFIELD varieties only. Non-CLEARFIELD varieties will be severely injured by this treatment. ⁸ CLEARFIELD varieties not controlled. • Registered for control. S - Suppression.

Table 13a. Herbicides to Control Weeds Before Seeding or After Seeding but Prior to Crop Emergence

HERBICIDE	PAGE	Pre-seeding	Pre-emergent	Barley	Canaryseed	Canola	Chickpea	Corn, Field	Corn, Sweet	Dry Bean	Field Pea	Flax	Forage Grasses	Lentil	Oat	Rye	Soybean	Wheat
Aim	77	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Amitrol	82	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Authority Charge	88	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
CleanStart	117	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Florasulam + glyphosate	160	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate	178	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + 2,4-D	178	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + Bromoxynil	178	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + Bromoxynil/MCPA	178	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate/dicamba	187	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + Heat	195	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + MCPA ¹	218	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓
Glyphosate + tribenuron	283	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓	✓

¹ Rates of application varies among brands. Consult the product page for application rates.

Table 13b. Weed Control Before Seeding or After Seeding but Prior to Crop Emergence

HERBICIDE	PAGE	Brome, Downy	Buckwheat, Wild	Dandelion	Flaxweed	Foxtail Barley	Foxtail, Green	Hemp-nettle	Kochia	Lady's-thumb	Lamb's-quarters	Mustard, Wild	Narrow-leaved Hawk's-beard	Pigweed, Redroot	Quackgrass	Russian Thistle	Shepherd's-purse	Stinkweed	Volunteer Cereals	Volunteer Canola (including glyphosate tolerant varieties)	Volunteer Flax	Wild Oats
Aim/Authority Charge ³	77	•	•						•		•	•		•				•		•		
Amitrol 240	82		•	•																		
CleanStart	117	•	•	• ²	•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Florasulam + glyphosate	160	•	•	•	•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate	178	•	•	•		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Glyphosate ¹ + 2,4-D	178	•	•		•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + Bromoxynil	178	•	•				•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate ¹ + Bromoxynil/MCPA	178	•	•		•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate/dicamba	187	•	•		•		•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + Heat	195	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Glyphosate + MCPA	218	•	•				•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + tribenuron	283	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•

• Controlled S - Suppression

¹ Rates of application varies among brands. Consult the product page for application rates.² Spring seedlings only. ³ Initial burndown only. For extended in season control see Authority Charge in crop tables 6,7,8 and 11.

Table 13c. Herbicides for Use as Harvest Aid or Dessicant Before Crop Harvest

HERBICIDE	PAGE	Alfalfa ⁷	Barley	Canola	Chickpea	Dry bean	Faba bean	Forage	Field Pea	Flax	Lentil	Oat	Potato	Soybean	Sunflower	Wheat
Aim ^{3,4}	77		•			•			•			•	•	•		•
CleanStart	117		•			•			•			•	•	•		•
Diquat ^{3,5}	134			•	•	•		•	•	•	•	•	•	•	•	
Humioxazin ³	166					•										
Glyphosate	178		•	•	•	•	•	•	•	•	•	•		•		•
Good Harvest	189	•									• ⁶		• ⁶			
Heat ^{3,8,4}	195					•			•		•			•	•	

¹ Rates of application vary among brands. Consult glyphosate page for specific application rates. ² For pre-harvest perennial weed control and may provide harvest management benefit. ³ For rapid plant tissue dry down to facilitate harvest. ⁴ May be tank mixed with glyphosate when used prior to harvest. ⁵ Refer to product page for surfactant requirements. ⁶ Not for crops grown for seed. ⁷ Seed production only. ⁸ Red lentil only.

Table 14. Weed Control in Summerfallow (Chem Fallow)

HERBICIDE	PAGE	Brome, Downy	Buckwheat, Wild	Dandelion	Flaxweed	Foxtail Barley	Foxtail, Green	Kochia	Lady's-thumb	Lamb's-quarters	Mustard, Wild	Pigweed, Redroot	Quackgrass	Russian Thistle	Stinkweed	Thistle, Canada	Sow-thistle (Perennial)	Volunteer Cereals	Volunteer Canola	Wild Oats
Dicamba + 2,4-D ³	126	•	•	•				•	•	•	•	•		•	•	•	•			
Dicamba/Mecoprop/MCPA	130	•	•	•				•	•	•	•	•		•	•	•	•			
Distinct	136		•	• ¹				•	•	•	•	•		•	•	•	•			
DyVel DSp	141	•	•	•				•	•	•	•	•		•	•	•	•		•	•
Express SG	283	•	•	•				•	•	•	•	•		•	•	•	•			
Express PRO	155	•	•	•				•	•	•	•	•		•	•	•	•			
Florasulam + glyphosate	160	•	•	•	•		•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate	178	•	•	•	•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + 2,4-D amine (500 g/L) ³	178		•	•				•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + Bromoxynil	178	•	•				•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate/dicamba	187	•	•	•	•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Glyphosate + Heat	195	•	•	•	•	•	•	•	•	•	•	•		•	•	•	•	•	•	•
Lontrel 360	216	•	•	•				•	•	•	•	•		•	•	•	•	•	•	•

• Controlled S - Suppression

¹ Top growth control only. ² Not including glyphosate tolerant canola.³ Rates of application varies among brands. Consult the product page for application rates. ⁴ Fall rosettes and spring seedling.

Table 15. Fall Weed Control in Stubble

HERBICIDE	PAGE	Flaxweed	Narrow-leaved Hawk's-beard	Shepherd's-purse	Stinkweed	Thistle, Canada	Quackgrass	Dandelion
2,4-D	86	•	•	•	•	•	•	•
Amitrol 240	82					•		
Dicamba	126					•		•
Dicamba + Glyphosate	126	•	•	•	•	•	•	•
Dicamba/ mecoprop/MCPA	130	•	•	•	•	•		
Distinct ⁶	136					• ¹		• ¹
DyVel DSP	141	•	•	•	•	•		
Florasulam + glyphosate	160	•	•	•	•			•
Glyphosate	178					•	•	•
MCPA	218	•	•	•	•	•	•	•

• Control S = Suppression. Levels of suppression vary depending on the product and growing conditions in the fall. Regrowth and in-crop treatments can be expected.

¹ Top growth control only.

⁶ To be used only in a mix with glyphosate.

Table 16. Weed Control in Grass Pastures and Hayfields

HERBICIDE	RATE (per acre)	PAGE	Absinth	Birdweed, Field	Burdock	Thistle, Canada	Dandelion	Dock, Curled	Daisy, English	Flaxweed	Footail Barley	Gumweed	Narrow-leaved Hawk's-beard	Knapweed	Leafy Spurge	Nodding Thistle	Poplar	Pussy Toes	Red Bartsia	Sage, Pasture	Snowberry	Sow-thistle, Perennial	Stinkweed	Tansy, Common	Wild Rose	Willow	Wormwood, Biennial
2,4-D (500 g/L)	0.57 - 1.82 L ¹	86	•	•	•	•	•	•		•		•	•		•	•	•	•	•	•	•	•				•	
2,4-DB	0.71 - 1.72 L ¹	89		•		•	•	• ¹					• ¹			•	•				•	•					
Dicamba	0.85 - 1.86 L ¹	126		•		•	•	•	•					•	•	•	•			•	•	•	•	•			
Dicamba + 2,4-D	0.85 + 0.89 L ¹	126	•	•	•	•	•	•	•	•	•				•	•	•		•	•	•	•	•	•	•	•	
Escort	10 - 12 g	151				•	•										• ¹					•		•	•	• ¹	
Grazon	2.8 L	191			•	•	•	•																			
Kerb	0.36 - 0.45 kg	207									•																
MCPA (500 g/L)	0.7 - 1.9 L ¹	218	•	•	•	•	•	•		•	•				• ¹							•	•	•		•	
MCPB/MCPA	1.11 - 1.72 L	221	•	•		•																•	•				
Overdrive	115 g	240					•								•											•	
Reclaim	20 acre per case	254				•	•													•					•		
Restore II	15 acre per case	257	•	•	•	•	•	•	•	•		•		•	•	•	•	•	•	•	•	•	•	•			
Tordon 22K	0.45 - 1.8 L	277		• ²		•				•					•	• ²				•							

• Controlled. S = Top growth suppression only. ¹ Rates may vary between different brands. Check product page for specific rate for product and use.

² May require multiple applications for complete control. ³ Controlled by the highest rate within this range.

Table 17. Weed Control in Shelterbelts

HERBICIDE	USE	SHELTERBELT SPECIES	WEEDS
	Before Planting		
	After Planting (Cuts & Establish)		
	Established		
	American Elm		
	Birch		
	Caragana		
	Crabapple		
	Green Ash		
	Juniper		
	Lilac		
	Manitoba Maple		
	Poplar		
	Scots Pine		
	Siberian Elm		
	Willow		
	Barnyard Grass		
	Footail, Green and Yellow		
	Wild Oats		
	Buckwheat, Wild		
	Chickweed		
	Cleavers		
	Cocklebur		
	Dandelion		
	Flaxweed		
	Hemp-nettle		
	Kochia		
	Lamb's-quarters		
	Mustard, Wild		
	Pigweed, Redroot		
	Quackgrass		
	Russian Thistle		
	Shepherd's-purse		
	Sow-thistle, Annual Species		
	Sow-thistle (Perennial)		
	Stinkweed		
	Thistle, Canada		
Amitrol 240	82		
Casoron	116		
Glyphosate	178		
Gramoxone	190		
Linuron	213		
Simazine	261		
Trifluralin Liquids	286		

• Controlled

¹ Yellow foxtail only.

Special Weed Problems

This section identifies specific weeds and some herbicides recommended for control. Refer to the particular crop section or the product label for information on specific products that may be used on the crops and for application instructions.

Absinthe

2,4-D LV Ester (500 g/L) - In grass pastures with no legumes, spray 1.82 L/acre in late June, prior to flowering. Re-treat regrowth in late summer when plants have 6 to 10 inches (15 to 25 cm) of new growth. More than 1 season of treatment may be required.

dicamba - In grass pasture and rangeland only, apply 0.5 L/acre in 20 to 30 gallons (90 to 135 L) per acre for top-growth control when leaves are fully expanded.

Restore/Restore II - In grass pastures and rangeland, apply *Restore A* at 0.2 L and *Restore B* at 1 L per acre (one package treats 15 acres) or *Restore II* at 0.97 L/acre (one 10 L jug treats 10 acres) when actively growing.

Alders

2,4-D LV Ester (500 g/L) - In grass pastures and non-crop land, apply 1.78 L/acre to the foliage of actively growing brush.

dicamba + 2,4-D amine or LV ester (500 g/L) - In grass pastures, rangeland and non-crop land, apply dicamba at 2.1 L per 1,000 L of water with 2,4-D LV ester or amine at 4.0 L per 1,000 L of water to the foliage of actively growing brush in the spring or early summer and wet the foliage until the point of runoff.

Aspen Poplar (Trembling Aspen)

dicamba + 2,4-D amine or LV ester (500 g/L) - In grass pastures, rangeland and non-crop land, apply dicamba at 1.32 L/acre with 2,4-D LV ester or amine at 1.78 L/acre in 20 gallons/acre (90 L/acre) water to the foliage of actively growing brush in spring or early summer.

Baby's Breath (Perennial)

dicamba - In grass pastures with no legumes, apply 3.72 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water when actively growing.

Biennial Wormwood

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing plants.

Overdrive - In grass pastures and non-crop land, apply at 115 g per acre for control.

Restore II - In grass pastures, apply 0.97 L/acre (one 10 L jug treats 10 acres).

Black Medic

Bromoxynil/MCPA ester; Dichlorprop/2,4-D; Mecoprop-p; dicamba/mecoprop/MCPA; 2,4-D amine or LV ester - Apply in registered crops at registered rates to black medic in the 1 to 4 leaf stage for suppression only.

Chokecherry

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing brush.

Common Tansy

Glyphosate - Apply at 1.9-2.8 L/acre in 10 gallons of water/acre (40 L/acre) to actively growing plants that are 8-10 inches (20-25 cm) tall (summerfallow, stubble and noncropland).

Escort - In pastures, rangeland and rough turf, apply 8 g/acre in 10 to 20 gallons/acre (45 to 90 L/acre) of water to actively growing plants of less than 4 inches (10 cm) tall. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Restore/Restore II - In grass pastures and rangeland, apply *Restore A* at 0.2 L and *Restore B* at 1 L per acre (one package treats 15 acres) when actively growing for season long control. Apply *Restore II* at 0.97 L/acre (one 10 L jug treats 10 acres) for suppression.

Curled Dock

dicamba - As a patch treatment or in pasture and rangeland, apply 0.92 L/acre Banvel II in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds for top growth control.

Glyphosate - As a spot treatment, apply 2.83 to 4.86 L/acre (360 g/L formulations or equivalent of other formulations) in 10 gallons/acre (45 L/acre) water when most plants have reached the early bud stage. Do not disturb treated plants for at least 10 days following treatment.

MCPA amine, 2,4-D amine - Apply 0.445 to 0.69 L/acre of formulations containing 500 g/L MCPA or 2,4-D amine to give top growth control.

Dichlorprop/2,4-D - 0.71 L/acre for suppression before plants are 2 inches (5 cm) tall.

Diffuse and Spotted Knapweed

dicamba - In grass pastures, rangeland and non-crop land, apply at 1.86 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds.

Restore - In grass pastures and rangeland, apply *Restore A* at 0.2 L and *Restore B* at 1 L per acre (one package treats 15 acres) when actively growing.

Tordon 22K - In rangeland and grass pasture, apply 0.91 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Downy Brome and Japanese Brome

Adrenalin, Altitude FX/FX2 - Apply at label rates to suppress Japanese brome to the 4 leaf stage in CLEARFIELD wheat.

Glyphosate - Prior to crop emergence, apply 0.51 to 0.77 L/acre (360 g/L formulations or equivalent of other formulations) in 5 to 10 gallons/acre (23 to 45 L/acre) water before downy brome is 6 inches (15 cm) in height.

glyphosate /dicamba - Prior to crop emergence, apply 1.0 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water between emergence and heading of downy brome.

Odyssey DLX - control spring seedlings of Japanese brome in registered crops.

PrePass - Apply in spring or fall, prior to seeding cereal crops or in fallow at registered rates to control downy brome up to the 4 leaf stage.

Simplicity - Suppression of downy brome and control of Japanese brome up to the 6 leaf stage when applied at 0.2 L per acre in the fall in winter wheat. Apply in spring at 0.2 L per acre to control Japanese brome up to the 6 leaf stage in winter or spring wheat (including durum).

Solo - Apply at label rates to suppress Japanese brome to the 4 leaf stage in registered crops.

Tandem - Applied at the maximum labelled rate in spring wheat (including durum) will control Japanese brome up to the 6 leaf stage.

Trifluralin - Apply at recommended rates for weed control in broadleaf crops prior to emergence.

Velocity m3/All-in-One - Apply at registered rates in registered crops to suppress Japanese brome.

Viper - In field peas, at registered rates to suppress Japanese brome.

Field Bindweed

dicamba - As a patch treatment or in rangeland, apply 1.0 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water. Apply when field bindweed is in the flowering stage and allow 3 weeks after treatment before resuming normal summerfallow tillage.

Basagran - In labelled crops, apply 0.71 L/acre followed by 0.71 L/acre 7 to 10 days later. Apply in 20 to 35 gallons/acre (90 to 160 L/acre) water before field bindweed has developed a dark green colour and before it has begun trailing. Use a recommended surfactant (see recommendations under the appropriate crop).

2,4-DB - As a spot treatment in labelled crops apply 2.83 to 4.86 L/acre in 10 gallons/acre (45 L/acre) water at the bud stage. Do not disturb plants for at least 10 days following treatment. Heavy rainfall within 2 hours of application may wash chemical off the foliage and a repeat treatment may be required. Rainfall occurring within 6 hours after application may reduce control.

2,4-D amine - In grass pastures containing no legumes or as a spot treatment, apply 1.82 L/acre of formulations containing 500 g/L 2,4-D amine at early flowering stage.

Glyphosate - As a spot treatment, apply 2.8 to 4.9 L/acre (360 g/L formulations or equivalent of other formulations) in 10 to 30 gallons/acre (45 to 135 L/acre) at the full bloom stage or beyond. Allow 7 or more days after application before tillage.

glyphosate/dicamba - Prior to crop emergence, apply 1.26 L/acre in 5 to 10 gallons/acre (23 to 45 L/acre) water to foxtail barley before initiation of the seed head for suppression only.

Restore II - For season long control in grass pastures and rangeland only, apply Restore II at 0.97 L/acre.

Tordon 22K - In rangeland and grass pasture, apply 3.6 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Field Horsetail

Amitrol 240 - Apply 5.0 to 6.7 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures when the weed is young and actively growing.

MCPA amine, potassium and sodium salt mixtures - Apply 0.57 L/acre of formulations containing 500 g/L MCPA after the weeds have fully emerged for top growth control. May be used in wheat, oats, barley, flax and rye.

Foxtail Barley

Glyphosate - Prior to crop emergence, apply 1 to 2 L/acre (360 g/L formulations or equivalent of other formulations) in 5 to 10 gallons/acre (23 to 45 L/acre) water to foxtail barley at the seedling to heading stage. Late fall applications may provide better control of established plants than spring applications.

Glyphosate - In glyphosate tolerant canola, apply 2 applications, each at 0.5 L/acre (360 g/L formulations or equivalent of other formulations), for season long control.

Gramoxone - Apply 2.23 L/acre in 98 gallons/acre (445 L/acre) water or 75 mL in 2.2 gallons (10 L) water/1076 square feet (100 sq. m) for top growth control only.

Kerb - Apply registered rates in 20 gallons/acre (90 L/acre) water between October 1 and freeze-up. Use the lower rate on grey-wooded soils or where perennial bluegrass or fescues are the predominant pasture species. Do not use Kerb for foxtail barley removal in seed grass stands or desired foliage stands of timothy or fescue grass species. At recommended rates, pasture stands of perennial bluegrass and fescue may be reduced by 10 to 15 percent. Where perennial bluegrass and fescues are the dominant pasture species, use the lower rate of Kerb. Spray overlaps may seriously harm desirable pasture grass species. Where the grass stand comprises mostly foxtail barley and reseeding to a desirable grass species is required, delay seeding into the Kerb-treated soil until the end of June. Do not harvest or graze within 60 days of application with Kerb. Avoid using Kerb on soils having more than 6 percent organic matter.

Goat's-Beard

2,4-D amine - Apply 0.91 L/acre of formulations containing 500 g/L in early fall or early spring.

dicamba - In grass pasture and rangeland only, apply 1.86 L/acre in 20 to 30 gallons (90 to 135 L) per acre when leaves are fully expanded.

Dichlorprop + 2,4-D - Apply 1.62 L/acre in early spring or fall.

Gumweed

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 0.89 L/acre to the foliage of actively growing plants.

Hemp Dogbane

2,4-D amine or LV ester - Apply 1.38 to 1.82 L/acre of formulations containing 500 g/L 2,4-D in fall before frost and while plant leaves are green.

Glyphosate - Apply 2.83 to 4.86 L/acre (360 formulations - see glyphosate page for other rates) when hemp dogbane is in the early bud stage. Apply in 10 gallons/acre (45 L/acre) water. Do not disturb treated plants for at least 7 days after application.

Hoary Cress

Amitrol 240 - For non-selective patch treatment in pastures and non-crop land, apply 8.9 to 13.8 L/acre.

Glyphosate - As a spot treatment in labelled crops, apply 2.83 to 4.86 L/acre (360 g/L formulations or equivalent of other formulations) in 10 to 30 gallons/acre (45 to 135 L/acre) water when most plants have reached the early bud stage. Do not disturb treated plants for at least 10 days following treatment.

Leafy Spurge

Amitrol 240 - Apply 15.2 to 18.5 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures when the weed is between the late stages of flowering and early seed development.

dicamba - Apply 0.84 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water for top growth control when the weed is actively growing. Patch treatment or pasture.

2,4-D amine - Apply 1.82 L/acre of formulations containing 500 g/L 2,4-D at early flowering stage. Repeat at least once to new growth later in the season. Control of established plants and new seedlings will require continued applications for a period of at least 4 to 5 years.

Tordon 22K - In rangeland and grass pasture, apply 3.6 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Overdrive - In grass pastures and non-crop land, apply at 115 g per acre for top-growth control.

Locoweeds, Lupines, and Milk-vetches

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 to 2.75 L/acre at the full bloom stage.

Milkweed

Amitrol 240 - Apply 7.6 to 11.3 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures in the early summer when all the shoots have emerged.

Glyphosate - When making Preharvest applications, use 1.0 L/acre (360 g/L formulations or equivalent of other formulations). For patch treatments, apply 4.86 L/acre (360 g/L formulations or equivalent of other formulations) in 10 gal-

lons/acre (45 L/acre) water. Apply when most plants have reached the bud to bloom stage. Reduced results may occur on plants treated after full bloom as not all milkweed plants reach the required stage of growth at the same time. Repeat treatments may be required. Do not disturb plants for 10 days following treatment. Do not apply to plants covered with dust.

Narrow-leaf Hawk's-beard

2,4-DB - Apply to forage legume crops at recommended rates at the 2 to 4 leaf stage of narrow-leaf hawk's-beard, after legume growth in the fall has stopped.

2,4-D LV ester (600 g/L) - In fall stubble, apply 0.57 to 0.90 L/acre to fall rosettes. Apply to fall seedlings or spring seedling to the 2 leaf stage at 0.22 to 0.38 L/acre or 0.4 to 0.6 L/acre in spring prior to bolting to control.

Adrenalin SC - Up to the 4 leaf stage in registered crops.

Barricade, thifensulfuron/tribenuron, Triton C, Triton K - Up to 4 inches tall in registered crops.

Express Pro - Up to 3 inches tall with residual activity, prior to the seeding of registered crops.

Express SG - For season long control in range and pasture at the early bud - pre-bloom stage.

Florasulam + glyphosate - up to 8 cm tall prior to seeding registered crops.

Frontline 2,4-D - Up to 2 leaf stage in registered crops.

Glyphosate - Prior to crop emergence, apply 0.51 to 0.77 L/acre (360 g/L formulations or equivalent of other formulations) in 5 to 10 gallons/acre (23 to 45 L/acre) water. Use the high rate if narrow-leaf hawk's-beard is between 3 and 6 inches (8 to 15 cm) in height.

Glyphosate - In glyphosate tolerant crops, apply 0.5 L/acre (360 g/L formulations or equivalent of other formulations) at the 0 to 6 leaf stage. Not all products are registered. Check glyphosate pages.

Heat - Apply in a mix with glyphosate for rapid burndown prior to seeding.

Tribenuron - Up to in a mix with glyphosate prior to seeding.

Velpar - Apply in late fall or early spring for control in established alfalfa in forage and seed production.

Pasture Sage

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.2 L/acre to the foliage of actively growing plants.

dicamba - In grass pastures, rangeland and non-crop land, apply dicamba at 1.86 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds.

Reclaim - In grass pastures and non-crop land, apply Reclaim A at 92 g per acre plus Reclaim B at 0.8 L per acre (20 acres per case) for 2 years of control.

Tordon 22K - In rangeland and grass pasture, apply 1.82 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to

permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Perennial Smartweed

Glyphosate - Apply 2.0 L/acre (360 g/L formulations or equivalent of other formulations) in 10 gallons/acre water. Apply when vines are a minimum of 8 inches (20 cm) tall, but before flowering.

Poplar

dicamba + 2,4-D - In grass pasture and rangeland only, apply dicamba at 2.1 L plus 2,4-D 500 amine at 4 L or 2,4D 600 ester at 3.3 L per 220 gallons (1000 L) of water and apply by wand to the point of runoff when leaves are fully expanded.

Glyphosate - As a non-selective spot treatment, apply 1.21 to 2.43 L/acre (360 g/L formulations or equivalent of other formulations) in 10 to 30 gallons/acre (45 to 135 L/acre) water in the summer through early fall when brush is actively growing.

Poverty Weed

dicamba - As a spot treatment or in grass pasture or rangeland apply 1.86 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water when weed is actively growing. Dicamba at 0.61 L/acre will provide only top growth control.

Restore II - For season long control in grass pastures and rangeland, apply Restore II at 0.97 L/acre (on 10 L container treats 10 acres).

Tordon 22K - In rangeland and grass pasture, apply 1.82 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Prairie Everlasting, Prairie Sage

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing plants in the early fall, and repeat in the spring.

Purple Loosetrife

(dryland situations only)

Glyphosate - Apply 2.43 L/acre (360 g/L formulations or equivalent of other formulations) in 30 to 60 gallons/acre (135 to 270 L/acre) water when purple loosetrife is actively growing and at or beyond the bloom stage. If using hand held equipment, apply a 1 to 2 percent solution until plants are wet. Use a 33 percent product solution if using a wiper applicator. Do not treat plants over open water. If possible, remove and destroy the flower heads before treatment to ensure prevention of seed set. For large monocultures of purple loosetrife, gradually work from the periphery inward over a number of years to allow competing vegetation to invade the treated area. Sprayed areas should be monitored for new seedlings to prevent re-infestation of purple loosetrife.

Red Bartsia

2,4-D amine or LV ester - Apply 0.57 L/acre of formulations containing 500 g/L 2,4-D in 10 gallons/acre (45 L/acre) water. On hayland, treat within 10 days after first cutting. Roadsides and pastures should be sprayed as soon as the red bartsia appears, usually in early July. Repeat treatment if necessary for later germination.

Roses

dicamba + 2,4-D amine or LV ester (500 g/L) - In grass pastures, rangeland and non-crop land, apply dicamba at 1.48 L/acre with 2,4-D LV ester or amine at 1.78 L/acre to the foliage of actively growing brush in the spring or early summer.

Escort - In pasture and rangeland, apply Escort at 0.012 kg/acre with non-ionic surfactant at 0.2 L per 100 L spray solution in 10 to 20 gallons/acre (45 to 90 L/acre) water. Apply between mid-June and mid-August after the brush has leafed out, but before the leaves begin to turn their fall colours.

Reclaim - In grass pastures and non-crop land, apply Reclaim A at 92 g per acre plus Reclaim B at 0.8 L per acre (20 years of control) for 2 years of control.

Russian Knapweed

Tordon 22K - In rangeland and grass pasture, apply 1.82 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

dicamba - In grass pasture and rangeland only, apply 3.72 L/acre in 20 to 30 gallons (90 to 135 L) per acre when leaves are fully expanded.

Restore II - In grass pastures and rangeland, apply Restore II 0.97 L/acre (one 10 L jug treats 10 acres) when actively growing for suppression.

Saskatoon

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing brush.

Scentless Chamomile

Bromoxynil/MCPA ester - Apply in registered crops at label rates when scentless chamomile is in the 2 to 4 leaf stage.

Curtail M - In registered crops, apply 0.81 L/acre in 10 gallons/acre (45 L/acre) water when scentless chamomile is actively growing and in the 2 to 4 leaf stage.

dicamba - Apply 0.51 L/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water to actively growing weeds for top growth control.

Escort - In pastures, rangeland and rough turf, apply 8 g/acre in 10 to 20 gallons/acre (45 to 90 L/acre) of water to actively growing plants of less than 4 inches (10 cm) tall. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Express SG - Apply in a mix with glyphosate prior to seeding registered follow crops.

Liberty - In registered crops, apply 1.1 L/acre to plants up to 4 inches (10 cm) in height.

Lontrel - In registered crops, apply 0.23 L/acre in 10 gallons/acre (45 L/acre) water when scentless chamomile is actively growing and in the 2 to 4 leaf stage.

metsulfuron plus 2,4-D - Apply 3 g/acre metsulfuron plus 0.34 to 0.45 L/acre 2,4-D LV ester or amine (500 g/L formulations) in 10 gallons/acre (45 L/acre) water for control of scentless chamomile up to the 4 leaf stage in wheat, barley, and creeping red fescue. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Restore/Restore II - In grass pastures and rangeland, apply Restore A at 0.18 L and Restore B at 0.6 L per acre (one package treats 16.7 acres) or Restore II at 0.86 L/acre (one 10 L container treats 11.6 acres) when actively growing.

Thifensulfuron/tribenuron - Apply 8 g/acre of DG formulations or 12 g/acre of Refine SG in 10 gallons/acre (45 L/acre) water to actively growing seedlings for suppression. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Tordon 22K - In rangeland and grass pasture, apply 0.445 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Triton C - Apply at label rates to suppress scentless chamomile up to 10 cm across or high.

Stinging Nettle

2,4-D amine - Apply 0.91 to 1.82 L/acre of formulations containing 500 g/L 2,4-D amine.

Stork's Bill

Altitude FX, Everest GBX, florasulam + MCPA, Prestige XC, Pulsar, Stellar, Tandem, thifensulfuron/tribenuron, Trophy - Apply at label rates to provide suppression in registered crops. See product pages for Crops, Rates and Staging.

Basagran - In registered crops apply 0.91 L/acre at the 2 to 6 leaf stage.

Dichlorprop/2,4-D - Apply at 0.71 L/acre to registered crops when stork's-bill is in the 2 to 4 leaf stage.

Fluroxypyr + 2,4-D, OctTain - Apply at the maximum labelled rate to registered crops when stork's-bill is in the 1 to 8 leaf stage.

Liberty - in registered crops apply 1.35 L/acre to plants in 1 to 3 leaf stage.

Linuron - Apply with MCPA amine in registered crops at registered rates to stork's-bill in the 2 to 4 leaf stage.

Metsulfuron - Apply with 2,4-D or MCPA amine or LV ester in registered crops at registered rates to stork's-bill in the 2 to 4 leaf stage.

Odyssey - in registered crops, apply 17 g/acre of the Odyssey component plus adjuvant.

Glyphosate - in glyphosate tolerant crops, apply 0.5 L/acre (360 g/L formulations or equivalent of other formulations) from emergence to the 6 leaf stage.

Spectrum - In registered crops apply at 20 acres per case to control from the 2 to 4 leaf stage.

Toadflax (Yellow)

Amitrol 240 - Apply 7.6 to 11.3 L/acre in 10 to 30 gallons/acre (45 to 135 L/acre) water in non-cropped areas and pastures when the weed is in the advanced rosette to prebud stage.

Dichlorprop/2,4-D - Apply 0.71 L/acre in 10 to 18 gallons/acre (45 to 80 L/acre) water in wheat or barley for toadflax suppression. Apply when majority of toadflax is no taller than 6 inches (15 cm). The use of Dichlorprop/2,4-D for suppression of toadflax in wheat or barley should be part of a long-term planned approach for toadflax control, which includes spring and fall tillage, fall patch spraying, summerfallow or chemical fallow.

Glyphosate - Apply 2.83 to 4.86 L/acre (360 g/L formulations or equivalent of other formulations) when most plants have reached the early bud stage of growth. Allow 7 more days after application before tillage. A rate of 1.0 L/acre may be used with Preharvest applications or when controlling in summerfallow situations.

Metsulfuron plus 2,4-D - Apply 2 to 3 g/acre Ally plus 0.34 to 0.45 L/acre 2,4-D LV ester or amine (500 g/L formulations) in 10 gallons/acre (45 L/acre) water for toadflax suppression in wheat, barley, and creeping red fescue. Add non-ionic surfactant at 0.2 L per 100 L of spray solution.

Thifensulfuron/tribenuron - In registered crops, apply 8 g/acre of DG formulations or 12 g/acre of Refine SG in 10 gallons/acre (45 L/acre) water for suppression of toadflax. Apply when toadflax is less than 15 cm (6 inches) in height. Add non-ionic surfactant at 0.2 L per 100 L spray solution.

Tordon 22K - In rangeland and grass pasture, apply 3.6 L/acre in 90 to 180 gallons/acre (400 to 800 L/acre) of water to actively growing weeds. **WARNING** - Picloram is a very persistent and water-soluble herbicide. Do not apply to permeable soil. Do not apply to irrigated areas. Take special precautions to prevent drift.

Western Snowberry (Buckbrush)

2,4-D amine or LV ester (500 g/L) - Apply 1.82 L/acre 2,4-D amine or LV ester in a minimum of 20 gallons/acre (90 L/acre) water in spring or early summer. Retreatment may be necessary the following year.

Dicamba plus 2,4-D LV ester (500 g/L) - Apply 1.48 L/acre dicamba tank mixed with 1.82 L/acre 2,4-D LV Ester in 20 gallons/acre (90 L/acre) water in spring or early summer after the leaves are fully expanded.

Escort - Apply 10 g/acre in 10 to 20 gallons/acre (45 to 90 L/acre) water between mid-June and mid-August after the brush has leafed out, but before the leaves turn their fall colours.

Reclaim - In grass pastures and non-crop land, apply Reclaim A at 92 g per acre plus Reclaim B at 0.8 L per acre (20 acres per case) for 2 years of control.

White Cockle

2,4-DB - Apply Embutox 625 at 1.1 L/acre or Caliber 400 at 1.7 L/acre or Cobutox 600 at 1.1 L/acre for top growth control to registered crops only.

Express SG - Apply in a mix with glyphosate prior to seeding registered follow crops to control spring rosettes.

Mecoprop - Apply 2.2 L/acre in 18 gallons water/acre (*80 L/acre) for top growth control of established plants. Will also control seedlings. Apply to registered crops only.

Wolf Willow (Silverwillow)

Dicamba + 2,4-D amine or LV ester (500 g/L) - In grass pastures with no legumes, apply dicamba at 2.1 L per 1000 L of water with 2,4-D LV ester or amine at 4.0 L per 1000 L of water to the foliage of actively growing brush in the spring or early summer and wet the foliage until the point of runoff.

Reclaim - In grass pastures and non-crop land, apply Reclaim A at 92 g per acre plus Reclaim B at 0.8 L per acre (20 acres per case) for 2 years of control.

Wild Tomato

2,4-D or MCPA amine or ester (500 g/L) - Apply 0.34 to 0.45 L/acre to registered crops up to the 8 leaf stage of wild tomato.

Bromoxynil+MCPA ester - Apply 0.40 L/acre to registered crops from the 1 to 6 leaf stage of wild tomato.

Willow

2,4-D LV ester - In grass pastures and non-crop land, apply 2,4-D LV ester (500 g/L) at 1.78 L/acre to the foliage of actively growing brush.

Dicamba + 2,4-D - In grass pasture and rangeland only, apply dicamba at 1.7 L plus 2,4-D 500 amine at 3.24 L per acre in 20 to 30 gallons (90 to 135 L) of water per acre when leaves are fully expanded.

Glyphosate - As a non-selective spot treatment, apply 1.21 to 2.43 L/acre (360 g/L formulations or equivalent of other formulations) in 10 to 30 gallons/acre (45 to 135 L/acre) water in the summer through early fall when brush is actively growing.

Soil Residual Herbicides

When applied at recommended rates in a crop, most herbicide residues will disappear within a few weeks after application and impose no restriction on cropping options the next year. However, some herbicide residues do not degrade quickly, and can persist in the soil for months or years following application, thereby restricting the crops that can be grown in rotation. Herbicide residues in the soil are deactivated in various ways including:

- Break down by chemical reactions,
- Break down by soil microbes,
- Escape to the atmosphere as a gas (volatilization),
- Break down by light (photodegradation),
- Leaching,
- Binding to soil particles.

Herbicides often disappear from the environment by more than one of these mechanisms. Many herbicides considered to be non-residual are bound temporarily to soil particles while they are broken down gradually by either soil microbes or chemical reactions. The binding action insures that the herbicide is not available to the crop in quantities that will cause damage.

As a general rule, breakdown processes are favoured by warm, moist soil conditions. During the winter, when the ground is frozen, and in the summer when the soil is dry, herbicide degradation is reduced. The residual activity of certain herbicides is also affected by soil organic matter and soil pH. These soil factors are seldom uniform across a field.

Herbicide carryover is aggravated by low levels of organic matter and is more likely to occur on eroded hilltops than in other parts of a field. The risk of herbicide carryover will also be greater in sprayer overlaps which are most common around headlands and slough margins.

Growers should be aware of the residual properties before applying any herbicide if they are to avoid cropping restrictions in following years. Knowledge of the limitations associated with herbicides that leave a soil residue, along with an accurate record of application (i.e. rates, locations) will serve to minimize rotational problems. Each herbicide used in mixes should be considered separately.

Soil tests using chemical extraction cannot always give a good indication of the potential injury risk from herbicide residue because of the influence of organic matter, clay and pH. Because of this, a field bioassay or laboratory bioassay, where plants are grown directly in the treated soil are best for detecting the potential for injury. These tests are not intended to be used to shortcut restrictions on the label, but provide information on rotational crops where none is available.

Injury symptoms from other causes can resemble herbicide carryover injury (i.e. cold weather, flooding, drought, insects, diseases, etc.). Consult with your local agronomist on potential causes before spending money on testing.

Herbicides that leave a soil residue and are of particular concern in Western Canada are found in the following chart.

Re-cropping Restrictions for Residual Herbicides:

Figures listed are the number of cropping seasons before each crop can be grown ("1" means that the crop can be grown the year following application). For plant-back restrictions less than one season; the delay is indicated with a "d" for number of days or with "mths" for the number of months. A blank space means that there are no recommendations given on the product label and a field bioassay is recommended by many product manufacturers to determine if these crops are safe to plant. A field bioassay is a strip of a test crop that covers an area of the field that is representative of the field variation and should include an untreated area.

PRODUCT	Alfalfa	Barley	Canaryseed	Clearfield canola	Non-Clearfield canola	Fababeans	Field corn	Dry beans	Field peas	Flax	Forage grasses	Lentils	Mustard†	Oats	Potatoes	Rye	Soybeans	Sunflowers	Wheat (durum)	Wheat (spring)	Wheat (winter)
2,4-D*	1	1	1	1	1		1	1	1	1	1	1		1		1			1	1	1
Accent		1					1													1	
Adrenalin, Solo, Viper ADV		1	1	1	1				1	1		1		1					1	1	
Altitude IX/IX2		1	1	1	1				1	1		1		1						1	
Amitrol		1d	1	1d	1d		10d*	10d*	5d*	1		1	1	1			6d	1	1d	1d	1d
AAtrex, Primextra II Magnum						1*	1		1*	1*											
Ares		1	1	1	2		1		1	2		1		1				2	2	1	
Authority / Authority Charge	1	1		2	2		1		0	0		3					0	0	1	1	1
Avadex	0	0	0	0	0	1	1	1	0	0		1	0	2		1	1	1	0	0	0
Barricade, Fluroxypyr + 2,4-D, Retain, Trophy	2	1	2	1	1	2	2	2	1	1	1	1	1	1	2	1	2	2	1	1	1
Battalion		1					1														1
Curtail M, Prestige XC	2	1	2	1	1	2	1	2	1*	1	1	2	1	1		1	2	2	1	1	1
Dicamba*		1		1*	1*		1	1*						1			1		0*	0*	1
Dual II Magnum							1								1		1				
Eclipse III, Lontrel 360		1		1	1					1	1		1	1		1			1	1	
Edge	0		2	0	0	0		0	0		2	0	0	2			0	0	1*	1*	
Express Pro		1d		10 mths	10 mths				10 mths										1d	1d	1d
Flucarbazon, Everest GBX (Brown soils)																				1	
Flucarbazon, Everest GBX (Dark Brown soils)		1*		1*	1*				1*	1*									1	1	
Flucarbazon, Everest GBX (Black soils)		1*		1*	1*			1*††	1*	1*									1	1	
Flucarbazon, Everest GBX (Grey-Wooded soils)		1*		1*	1*				1*											1	
Flumioxazin	1	1		1	1		1	1							1		0	1	1	1	1
Focus							1														
Frontier Max							0*	0*													1
Florasulam + MCTA, Frontline 2,4-D, Spectrum		1		1	1				1*					1					1	1	1

PRODUCT	Alfalfa	Barley	Canaryseed	Clearfield canola	Non-Clearfield canola	Fababeans	Field corn	Dry beans	Field peas	Flax	Forage grasses	Lentils	Mustard†	Oats	Potatoes	Rye	Soybeans	Sunflowers	Wheat (durum)	Wheat (spring)	Wheat (winter)
Imazamethabenz (Black and Grey Wooded soils)		1	2	1	1				1	1				2				1	1	1	
Imazamethabenz (Brown and Dark Brown soils)		1	2	1	2				2	2				2				1	1	1	
Imazethapyr	1	1		0					0			1								1	
Infinity / Velocity m†	1	1	1	1	1		1		1	1		2		1			1		1	1	
Kerb	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	2
Metribuzin				2	2	0			0						0*		0*	2			
Metsulfuron (pH less than 7, Brown and Dark Brown)		1	4		2					2		3	4	1					1	1	
Metsulfuron (pH less than 7, other soils)		1	4		1					1		3	4	1					1	1	
Metsulfuron (pH 7 to 7.9, Brown and Dark Brown soils)		1	4		3					3		4	4	2					1	1	
Metsulfuron (pH 7 to 7.9, other soils)		1	4		2					3		4	4	1					1	1	
Mustar	2	1	2		2	2		2	2	1	2*	2	2	1					1	1	1
Odyssey*, Odyssey DLX*	1	1	1	1	2		1		1	2		1		1				2	1	1	
Option	1	1		1	1		1	1	1					1	1		1			1	1
Florasulam + glyphosate (prior to Aug 1)		0d		1	1				1					0d					0d	0d	0d
Prism		1					1								1		1				0
Pulsar	2	1	2	1	1	2	2	2	1	1	1	1	1	2	2	1	2	2	1	1	1
Reflex*							1	0									1			1	0
Simplicity		1		1	1				1	1		1	1	1			1			1	
Stellar		1		1	1				1					1					1	1	1
Tandem		1		1	1				1	1		1	1	1						1	
Trifluralin	0	1*	2	0	0	0	1*	0	0	1*	2	0	0	2		0	0	0	1*	0*	1*
Triton C**		0		1	1				1	2*		2*						1	0	0	
Ultim		1					1														1
Varro	1	1	1	1	1		1	1	1	1		1		1			1	1	1	1	10m

* The minimum re-cropping intervals are listed. These intervals may be longer than those listed depending on the use rates, region, province, soil types, environment, time of application and crop variety. Refer to product page for more information.

** Drought restrictions apply to drought conditions (80% of normal June to September rainfall) for high pH soils (greater than pH 7.5) and severe drought (less than 65% of normal June to Sept. rainfall) for all soils.

† May not be valid for all varieties or crop types. See product page for details.

†† DO NOT grow dry beans the year following Everest GBX application.

0 - May be seeded or reseeded the year of application. No re-cropping restrictions. 1 - Next cropping season after application. 2 - Two cropping seasons after application. NR - Not recommended.

Note: The re-cropping intervals listed may not be sufficient to prevent crop injury during periods of below average rainfall.

Effect of Rainfall on Herbicide Efficacy

Required Interval	Product
15 minutes	Diquat
30 minutes	clodinafop
1 hour	Axial, Barricade II, Broadband, Bromoxynil, Bromoxynil/MCPA ester, clethodim, Equinox, fenoxaprop, flucarbazone Gramoxone, Harmony SG, Infinity, Poast Ultra, quizalofop, thifensulfuron/tribenuron, tralkoxydim, Traxos, Tundra, Velocity m3
2 hours	2,4-D LV Ester, metsulfuron+2,4-D LV Ester, Atrazine (post-emergent applications), fluroxypyr + 2,4-D ester, MCPA Ester
3 hours	dicamba/mecoprop-p/MCPA, Odyssey, Odyssey DLX
4 hours	Accent, metsulfuron + 2,4-D Amine, 2,4-D Amine, Battalion (post-emergent application), Liberty (both), MCPA Amine, Overdrive, Prism, Reflex, Simplicity, Tandem, Ultim
6 hours	Assert FL, Avenge, Blazer, Curtail M, glyphosate/dicamba, imazamethabenz, MCPA-K, MCPA Sodium Salt, Muster, Option, Prestige XC, Metribuzin, Tordon 22K, Triton C
8 hours	Basagran, CleanStart
No specific recommendation*	2,4-DB, Adrenalin, Aim, Altitude FX/FX 2, Amitrol 240, Bromoxynil/2,4-D ester, dicamba, dichlorprop/2,4-D, DyVel, DyVel DSp, Escort, Express Pro, florasulam + glyphosate, florasulam + MCPA, Frontline+2,4-D, glyphosate, Grazon, Harmony K, imazethapyr, Linuron, Lontrel, MCPB/MCPA, mecoprop-p, Momentum, Optica Trio, Pinnacle, Pulsar, Reclaim, Restore, Solo, Spectrum, Stellar, Tensile, tribenuron, Triton K, Trophy, Viper ADV

* The products listed make no specific time recommendation on the label. The required rainfree period could be up to 8 hours. See the product page in the guide or consult the product label.

Note: The term "Rainfastness" refers to the time needed between application and rainfall to avoid significant reduction in efficacy. Rainfall shortly after application of most post-emergent herbicides may reduce weed control. Effect will vary with product, the interval between spraying and rainfall and the intensity and duration of the rainfall. These guidelines are based on label information. Use the longest time interval on the component products when considering tank mixes.

Products Available as Prepackaged Tank Mixes

Product Name (Manufacturer)	Component 1 or A	Component 2 or B	Component 3 or C	Crops	Weeds Controlled	Area Treated per Package
Achieve Liquid Gold (Nufarm)	Achieve Liquid	Buctril M	Turbocharge (adjuvant)	Wheat (spring, durum, winter), barley	See the weeds of the component products plus tame buckwheat	20 acres or 8 ha
Authority Charge (FMC of Canada)	Aim EC	Authority		Chickpea, Field Pea, Flax and Sunflower	See component products	80 to 64 acres
Axial iPak (Syngenta)	Axial	Infinity	Adigor (adjuvant)	Spring wheat (NOT including durum), barley	See component products	40 acres or 16 ha
Black Hawk (Nufarm)	Aim EC	2,4-D 700 ester		Wheat (Spring, durum, winter), barley, rye	See component products	80 acres or 32 ha
Eclipse III (Dow AgroSciences)	Lontrel (Eclipse A)	Vantage Plus Max II (Eclipse III B)		Roundup Ready Canola varieties	See component products	40 acres or 16 ha
Harmony SG (E. I. duPont)	Refine SG	Harmony Grass (clodinafop)	Score (adjuvant)	Spring wheat (including durum)	Weeds controlled by Refine SG plus wild oat, green foxtail	40 acres or 16 ha
Harmony K (E. I. duPont)	Refine SG	Harmony Grass (clodinafop)	dicamba	Spring wheat (NOT including durum)	Weeds controlled by Refine SG plus wild oat, green foxtail & group 2 resistant kochia	40 acres or 16 ha
Ko-Act (Nufarm)	Spike	2,4-D 700 ester		Wheat, barley, rye	See component products	80 acres or 32 ha
Odyssey DLX (BASF)	Odyssey	Equinox	Merge (adjuvant)	Field peas, CLEARFIELD lentil	See component products	40 acres or 16 ha
Refine M (E. I. duPont), Broadside (Loveland Products (formerly Viterro))	Refine SG	MCPA Ester		Spring wheat (including durum), barley, oat, winter wheat	See component products	80 acres or 32 ha
Tersile	Solo	Lontrel Dry		CLEARFIELD canola	See component products	40 acres or 16 ha
Triton K (E. I. duPont)	Express SG	2, 4-D LV Ester	Banvel II	Spring wheat, barley	See component products	40 acres or 16 ha

Note: See the component products listed for information concerning staging, application information, safety precautions, the effect of weather and grazing, re-cropping, harvest interval and storage precautions.
The more stringent recommendation of the two products should be followed. Mix products in the order listed.

Herbicide Product Pages

2,4-D

Herbicide Group
4 - 2,4-D
(Refer to page 38)

Company and Formulation:

	PCP # (Product Name)	
	600 Amine*	700 Ester**
IPCO	17511	27819
Nufarm Agriculture	14726	27820
United Agri-Products	5931	27818 (Salvo)
Loveland Products (formerly Viterra)	29248	29006
Farmers of North America		30460 (MPower 2,4-D)

* formulated as a solution.

** formulated as an emulsifiable concentrate.

Crops, Rates and Staging:

Application rates for individual products may vary from those listed. Refer to the label for product specific use rates. Rates greater than those listed may cause crop injury.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

CROP	MAXIMUM SAFE RATE (L/ACRE)	STAGE
Wheat, barley, spring rye	0.38 (600 g/L) 0.32 (700 g/L)	4 leaf to early flag leaf.
Fall rye, winter wheat*	0.36 (600 g/L) 0.32 (700 g/L)	In spring, apply after winter cereals begin grow but before emergence of the flag leaf. From full tillering to prior to flag leaf stage.
Corn*	0.40 (600 g/L) 0.32 (700 g/L)	Apply as an overall spray before corn is 6 inches (15 cm) tall and before the 6-leaf stage. After 6 inches (15 cm) use a directed spray. Avoid making applications under hot/humid conditions onto corn.
Seedling and established grasses for forage and seed production*	0.38 (600 g/L) 0.32 (700 g/L)	Apply from the 3 leaf stage to emergence of the flag leaf of seedling grasses. For established grasses for seed production, apply in spring up to emergence of the flag leaf.
Established forage grass (not for seed production)*	0.77 (600 g/L) 0.65 (700 g/L)	Apply in spring up to emergence of the flag leaf of established grasses, or in the fall after harvest.
Established grass pastures	1.62 (600 g/L) 1.90 (700 g/L)	No restrictions, apply when weeds are actively growing. For control of brush species, apply at time of rapid growth (usually May to mid-June, and September prior to colour change).
Turf*	0.77 (600 g/L) 0.77 (700 g/L)	Apply in spring and early September on established turf. Do not use on bent grasses, or newly seeded stands.

* Note: Registered for use only with certain brands of 2,4-D; use of non-registered products is at the risk of the user.

Weeds, Rates and Staging:

Apply at lower rates when weeds are small (2 to 4 leaf stage) and actively growing. Higher rates are needed when weeds are larger, in heavy populations, or growing under stressful conditions (excessively cold, hot, dry or wet).

Note: The rates listed differ slightly from product to product. Check individual product labels for exact use rates.

Susceptible Weeds:

0.22 to 0.38 L per acre (600 g per L)

0.19 to 0.32 L per acre (700 g per L)

Bluebur	Prickly lettuce
Burdock	Ragweed (common, false and giant)
Cocklebur	Russian pigweed
False flax	Russian thistle
Flixweed (late fall application or spring seedlings)	Shepherd's-purse**
Goat's-beard	Stinging nettle
Kochia	Stinkweed**
Lamb's-quarters	Sweet clover
Mustards (except dog and tansy mustard)	Thyme-leaved spurge
Narrow-leaved hawk's-beard (fall application to seedlings or spring application at 1-2 leaf stage)	Volunteer canola (including all herbicide tolerant varieties)
Plantain	Wild radish
	Wild sunflower

Harder to kill weeds:

0.40 to 0.61 L per acre (600 g per L)

0.34 to 0.53 L per acre (700 g per L)

Annual sow-thistle	Leafy spurge*
Blue lettuce*	Mustard (dog & tansy)
Burdock (top growth only of bolting plants)	Narrow-leaved hawk's-beard (spring prior to bolting)
Canada thistle***	Oak-leaved goosefoot
Common chickweed	Pineappleweed
Common groundsel**	Prostrate pigweed
Common peppergrass	Purslane
Dandelion*	Redroot pigweed
Flixweed (spring prior to bolting)	Russian thistle
Knotweed	Sheep sorrel
Lady's-thumb	Tumble pigweed

Top Growth Control only (at rates for harder to kill weeds):

Biennial wormwood	Hoary cress
Bull thistle	Horsetail
Buttercup	Mouse-eared chickweed
Curled dock	Perennial sow-thistle
Field bindweed	Russian knapweed
Gumweed	Tartary buckwheat
Hedge bindweed	Volunteer sunflower

* Control of seedlings at rates given above and top growth control only of established plants.

** Spring seedlings. Winter annual weeds - apply in late fall or early spring prior to bolting.

*** Suppression only - Apply when Canada thistle plants are actively growing and have 6 to 8 inches (15 to 20 cm) of new growth. Regrowth will be present the following spring and in-crop treatments will be required.

Formulation Characteristics:

Formulation	Risk of Vapour Drift	Activity on Weeds	Risk of Crop Injury
LV Ester	Medium	Fast	Medium
Amine	Very Low	Medium	Low

Application Information:

Water Volume: Minimum 20 L per acre - ground application. Water rates depend on product and use. Consult label for details.

Higher application volumes (40 L/acre or greater), reduce the risk of crop injury.

Nozzles and Pressure: 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressures designed to deliver proper coverage with ASABE coarse droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Best weed control occurs when temperatures are above 21°C (daytime) or 10°C (nighttime) and humidity is above 70 percent. DO NOT apply if temperature exceeds 27°C.

Tank Mixes:

Mixes listed may not occur on all products labels. Check individual product labels for registered mixes.

Note: Following list is for mixes listed on 2,4-D labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Herbicides:

Wheat and barley:

Banvel (2,4-D amine only)

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: 2,4-D amine: within 4 hours will reduce control.
2,4-D LV ester: within 2 hours will reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours

Grazing: DO NOT permit lactating dairy animals to graze fields within 7 days of application. Do not harvest forage or cut for hay within 30 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Re-cropping: No recropping guidelines are provided on the labels. As a general guideline, there should be no cropping restrictions the year following an in-crop treatment.

Aerial Application: Some formulations may be applied by air. Check the label for detailed instructions.

Storage: 2,4-D LV ester may be frozen. 2,4-D amine requires heated storage.

Buffer Zones:

Crop	Application method	Buffer Zones (metres†) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Field Crops	Ground*	1	0	0
	Fixed wing aircraft	10	0	45
	Helicopter	10	0	40
Fallow, stubble, pastures, range-land	Ground*	1	1	2
	Fixed wing aircraft	15	0	60
	Helicopter	15	0	50
Turf	Ground only*	1	0	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Warning – Poison



Danger Poison – Ester 700 Formulations

For an explanation of the symbols used here see page 11.

2,4-DB

Herbicide Group
4 - 2,4-DB
(Refer to page 38)

Company:

IPCO (Cobutox- PCP# 27911)
Nufarm Agriculture (Embutox- PCP# 27912)
United Agri Products (Caliber- PCP# 27910)

Formulation:

625 g/L 2,4-DB formulated as an emulsifiable concentrate.
Container size - 10 L.

Crops and Staging:

NOTE - When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

CROP	STAGE
Seedling alfalfa, bird's-foot trefoil*	1 to 4 trifoliolate leaf stage
Clover (alsike**, red**, white, Dutch but NOT sweet clover)*	As soon as possible after emergence of the 1st trifoliolate leaf
Wheat, barley or oats	5 leaf to emergence of the flag leaf
Field corn	15 inches (38 cm) to prior to tassling using drop nozzles.
Pastures containing forage legumes	After cutting or grazing and regrowth less than 3 inches (7.5 cm)

* With or without a cereal cover crop.

** Alsike and red clovers may be damaged by 2,4-DB applications.

Seedling Forage Grasses*:

Apply from 2 to 4 leaf stage of:

Bromegrass (smooth)	Timothy
Fescue (creeping red, meadow, tall)	Wheatgrass (crested, intermediate, streambank, tall)
Orchard grass	

* Not for seed production. Not for feeding in the establishment year.

Weeds and Staging:

Weeds controlled at the 0.71 L per acre rate from the 2 to 4 leaf stage at lower recommended rates include:

Ball mustard	Shepherd's-purse
Lamb's-quarters	Stinkweed
Ragweed	Wild mustard
Redroot pigweed	Wormseed mustard

Weeds controlled at higher recommended rates (0.91 to 1.1 L/acre) include:

WEED	STAGE
Bull thistle	Rosette to early bud stage
Canada thistle*	6 in. (15 cm) to early bud
Chicory	Rosette
Curled dock**	Young and actively growing
Dandelion*	Prior to bud
Field bindweed*	Late summer
Green smartweed**	Seedlings
Horsetail*	4 to 5 inches (10 to 13 cm)
Narrow-leaved hawk's-beard	Apply at rosette stage after alfalfa has gone dormant
Oak-leaved goosefoot	Up to 2 leaf stage
Perennial sow-thistle*	Rosette
Lady's-thumb**	Seedlings
Plantain	Prior to flowering
Wild buckwheat	Up to 2 leaf stage
Wild Radish	Up to 2 leaf stage
Yellow rocket	Late September to mid-October

* Top growth control

** Suppression

Refer to individual product labels for details on application rates to use for different weed species.

Rates:

CROPS	RATE (L/ACRE)	ACRES TREATED PER CONTAINER
Cereals, seedling forage legumes and grasses	0.71 to 0.91	14.1 to 11.0
Corn and pastures containing forage legumes	0.71 to 1.11	14.1 to 9.0

Application Information:

Water Volume: 61 to 81 L per acre.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressures designed to deliver proper coverage with a minimum of fine droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Severe damage to legumes can occur if high temperatures (more than 27°C) or high humidity prevail at the time of application. DO NOT apply under dry soil/drought conditions.

Tank Mixes:

Herbicides:

Underseeded Legumes: MCPA amine (28 mL/acre*)

* 500 g/L formulation

This tank mix may increase crop damage (stunting).

Follow all precautions and restrictions on both product labels.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on 2,4-DB labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Not specified on the label. A period of up to 8 hours may be required. Contact the manufacturer for more details.

Grazing: DO NOT graze or cut treated crops or forage until 30 days after application.

Re-cropping: No restrictions the year after application.

Aerial Application: DO NOT apply by air

Storage: May be frozen.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

Caution - Poison

For an explanation of the symbols used here see page 11.

AAtrex Liquid

Herbicide Group

5 - atrazine

(Refer to page 38)

Company:

Syngenta (PCP# 18450)

Formulations:

480 g/L atrazine formulated as a liquid suspension.

Container sizes - Various.

Crops, Rates and Staging:

Corn (silage, field, sweet): 0.85 to 1.25 L per acre* using the following application methods;

Pre-plant incorporated (PPI).

Pre-emergent surface (after planting but before emergence of weeds and crop): Recommended only on irrigated fields. Inconsistent weed control will occur if 0.5 inches (1.25 cm) of water/precipitation does not occur within 7 days of application.

Post-emergence: 1 to 6 leaf stage and when corn is less than 12 inches (30 cm) tall. Add 1.11 to 2.23 L per acre of oil concentrate or 6.88 L per acre crop oil. Crop injury may occur when Aatrex and oil is applied post-emergence during cold weather.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

* Use the low rate on crops grown on sandy soils, and where weed infestations are light.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Weeds and Staging:

For pre-plant incorporated, pre-emergent and post-emergent (when weeds are less than 4 inches or 10 cm tall) control of the following weeds:

Common purslane	Volunteer clover
Lamb's quarters	Wild buckwheat
Ragweed	Wild mustard
Redroot pigweed	Wild oats
Smartweed (including lady's-thumb)	Wormseed mustard

Application Information:

Water Volume: Minimum 61 L per acre.

Nozzles and Pressure: 30 to 45 psi (200 to 300 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift. Use a 50 mesh or coarser screen and filler system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Post-emergent applications made during periods of cold weather may cause crop lightening. Hot, dry weather preceding post-emergent applications may result in reduced weed control. Aatrex will move with soil if eroded.

Tank Mixes:

Herbicides:

Pre Plant Incorporated: Dual II Magnum

Pre-Emergent: Dual II Magnum

Post-Emergent*: Raxcel II (not sweet corn), Pardner, Dual II Magnum and Buctril M**.

* DO NOT use oils or adjuvants with post-emergent tank mixes.

** DO NOT treat after the 6 leaf stage, crop injury may occur.

Fertilizers: For pre-emergence applications, nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier. Aatrex may be impregnated onto dry granular fertilizers. DO NOT impregnate onto nitrate, super-phosphate or limestone.

DO NOT apply Aatrex with nitrogen fertilizer after corn has emerged, as crop injury will occur.

Insecticides: None registered.

Note: The above mixes are those listed on the Aatrex label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. See the general guidelines for mixing pesticides for more information.

Restrictions:

Rainfall: Within 2 hours of post-emergence applications may result in reduced weed control.

Grazing: DO NOT graze or cut for feed before ear emergence.

Preharvest: Leave at least 45 days from application to harvest for sweet corn and 60 days for field corn.

Re-cropping: All crops, except corn and triazine-tolerant canola, may be affected the year following the use of atrazine. Flax, peas and fababeans have some tolerance to atrazine residues and are usually not affected by rates of up to 0.9 L per acre applied the previous year. Other more sensitive crops may be affected 2 or more growing seasons after application.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground only*	10	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

DO NOT mix or load within 30 m of any wells, lakes, streams, ponds, dugouts or sinkholes.

Tank Cleaning:

When finished spraying atrazine, run clean water through the tank, pump and lines. Drain and refill with 1 L of 3% household ammonia solution per 100 L water. Circulate the solution through lines and nozzles. Let solution stand for several hours. Scrub inside surfaces but do not enter tank. Flush sprayer system with water.

DO NOT clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Refer to page 15 for additional information.

Hazard Rating:



Caution - Eye Irritant

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed.

For an explanation of the symbols used here see page 11.

Accent*

Herbicide Group

2 - nicosulfuron

(Refer to page 38)

Company:

E. I. duPont Canada (PCP# 25116)

Formulation:

75% nicosulfuron formulated as a water dispersible granule.

Container size - 133.6 g (4 x 33.4 g water soluble bags per pouch).

Crops and Staging:

Field Corn: 1 to 8 leaf stage (six visible collars), coleoptile (short, blunt leaf) is counted as the first leaf.

Sweet corn **: 1 to 6 leaf stage (4 visible collars).

* NOTE - Since applications to field and sweet corn in western Canada has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to corn is at the risk of the user.

** Note that Accent is registered for use on all sweet corn varieties but tolerance may vary depending on variety. Krispy King, Jubilee and Jubilee Supersweet are the only varieties that have been tested for tolerance in western Canada. Test on small areas planted to other varieties for tolerance prior to widespread use.

Weeds and Staging:

WEEDS	STAGING
Barnyard grass, green foxtail, yellow foxtail*, old witchgrass	1 to 6 leaves (up to 2 tillers)
Quackgrass	3 to 6 leaves (with extended leaf 4 to 8 inches (10 to 20 cm) long)
Wild oats	3 to 6 leaves

* Suppression only

The best control and yield response is achieved by applying at the earlier end of the leaf stage ranges.

Rates:

13.5 g per acre. Add non-ionic surfactant (Citowett Plus, Agsurf or Agral 90) at 0.2 L per 100 L of spray solution.

One water soluble bag will treat 2.5 acres (1 ha).

One pouch will treat 10 acres (4 ha).

Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum 40 L per acre; optimum 56 to 77 L per acre.

Nozzles and Pressure: 25 to 40 psi (175 to 275 kPa) when using conventional flat fan nozzles tilted forward at a 45° angle. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressures designed to deliver proper coverage with *ASABE medium* droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Poor weed control or crop injury may result if at the time of application, plants are under stress from disease, insect or nematode injury, carryover of herbicide from a previous years application, abnormally hot or cold weather, drought, water-soaked soils, hail damage or frost. Delay application until stress passes and both corn and weeds have resumed growth. When corn is injured by frost, wait 48 to 72 hours after normal growing conditions have resumed before applying *Accent*. Stress conditions after application may also result in injury or poor weed control.

Tank Mixes:

Herbicides:

Field corn only:

Banvel II (0.24 L/acre) plus surfactant.

Pardner (0.4 L/acre) plus surfactant.

Fertilizers: Do not mix with fertilizers.

Insecticides: None registered. *Accent* should not be applied to corn that has been treated with Thimet, or other organophosphate insecticides. Leave 7 days between the application of *Accent* and that of a foliar organophosphorous insecticide.

Fungicides: None registered.

Note: The above mixes are those listed on the *Accent* label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: Within 2 to 4 hours of application may result in reduced weed control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated crops or cut for hay.

Preharvest: Leave at least 30 days in field corn and 40 days in sweet corn from application to harvest.

Re-cropping: Winter wheat may be seeded 4 months following *Accent* application. Alfalfa, barley, canola, field corn, red clover, sorghum, soybean, white bean may be seeded 10 months following application. For all other

crops a field bioassay is recommended before planting. For all other crops, a field bioassay is recommended before planting.

Aerial Application: DO NOT apply by air.

Storage: Store product in original containers in a secure, dry area, away from other pesticides, food or feed.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of Terrestrial habitat
Ground only*	2

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Leave a 5 m buffer between the last spray path and woodlots or shelterbelts. Leave a 22 m buffer before wetland areas or water bodies.

Tank Cleaning:

Drain tank and hose down interior surfaces. Flush tank, hoses, boom, and nozzles with clean water for a minimum of 5 minutes. Fill spray tank with a water-ammonia cleaning solution (1 litre of a minimum 3% household ammonia for every 100 L of water). Flush hoses, boom and nozzles with the cleaning solution, then add more water to completely fill the tank. Circulate for 15 minutes, then flush hoses, boom and nozzles with the cleaning solution, and drain the tank. Remove and clean the nozzles and screens separately in a bucket containing the cleaning solution as above. If the spray equipment is to be used to spray crops other than corn, repeat the above process and thoroughly wash the spray mixture from the outside of the spray tank and the boom. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing the water through the hoses and boom. Prior to using the sprayer again, flush the tank, boom and hoses for 5 minutes with fresh water. Do not clean equipment where cleaning solution could flow towards water bodies, ditches, cropland, shelterbelts, or areas where people are likely to frequent or walk.

For additional information, Refer to page 15.

Hazard Rating:

 Caution – Eye Irritant

KEEP OUT OF REACH OF CHILDREN.

Avoid breathing spray mist.

Avoid contact with skin, eyes and clothing.

For an explanation of the symbols used here see page 11.

Achieve Liquid Gold

Herbicide Group

1 - tralkoxydim

4 - MCPA

6 - bromoxynil

(Refer to page 38)

This product is the equivalent of a prepackaged tank mix of *tralkoxydim* (page 279) and *bromoxynil/MCPA ester* (page 113). Information listed is restricted to Crop, Weeds and Rates. For other detailed information on the component products see the product pages listed above.

Company:

Nufarm Agriculture

Formulation:

Achieve Liquid Gold contains 3 components:

Achieve Liquid (PCP# 27011): 400 g/L tralkoxydim formulated as a suspension concentrate.

Container sizes - 4 L; 4 x 20 L.

Mextrol 450 (PCP# 26999): 225 g/L bromoxynil and 225 g/L MCPA ester formulated as an emulsifiable concentrate.

Container size - 10 L; 100 L; 500 L

Turbocharge adjuvant (PCP# 23135):

Container size - 4 L; 2 x 40 L.

Crops and Staging:

Spring wheat (including durum), barley - 2 leaf to early flag leaf stage.

Winter wheat, fall rye - from the time growth begins in the spring to the early flag leaf stage.

Weeds and Staging:

Grass weeds: Same weeds and staging as tralkoxydim plus;

Broadleaf weeds: Same weeds and staging as bromoxynil/MCPA ester

Rates:

One case treats 20 acres or treats 400 acres.

Add *Turbocharge* at a rate of 0.5 L per 100 L spray solution. If water analysis shows bicarbonate levels are 400 ppm or greater, add ammonium sulfate at 0.75 to 1.5 kg per 100 L water.

Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 14.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre. DO NOT apply with air assist sprayers set to apply less than 20 L per acre water volume as mixing problems or unacceptable crop injury could occur.

Aerial: 12 to 18 L per acre.

Nozzles and Pressure: 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Consult with herbicide manufacturer regarding the suitability of low drift nozzles for use with this product.

Use a 50 mesh or coarser screen and filter system.

Effects of Growing Conditions:

Crop Safety: Applications of *Achieve Liquid Gold* to crops prior to tillering and exposed to 4°C or lower 48 hours before or after spraying should be avoided to prevent the possibility of crop injury. Tillered cereals may incur injury if *Achieve Liquid Gold* is sprayed within 48 hours of 0°C or lower.

Tank Mixes:

Herbicides: MCPA ester (0.22 L/acre) (500 g/L formulations). A reduction in green foxtail control may occur with this tank mix.

Insecticides: *Matador* (25 to 34 mL/acre).

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Aerial Application: May be applied by air.

For additional information see the component products.

Adrenalin SC*

Company:

BASF Canada (PCP# 27879)

Formulations:

20 g/L imazamox and 560 g/L 2,4-D ester formulated as a suspension concentrate.

Container size - 2 X 8L.

* NOTE: This product is no longer manufactured but there is still inventories in the distribution system. This page may be removed from future editions of this guide.

Crops and Staging:

CLEARFIELD wheat varieties - 4 leaf to 6 leaf stage.

Apply only to CLEARFIELD wheat varieties; application to any other variety of wheat or any other crop will result in crop death.

Weeds and Staging:

Grasses:

Apply from 1 to 4 main stem leaves, up to a maximum of two tillers.

Barnyard grass	Volunteer tame oat
Foxtail (green and yellow)	Volunteer wheat
Japanese brome*	(non-Clearfield)
Volunteer barley	Wild oat
Volunteer canaryseed	

* Suppression only.

Herbicide Group

2 - imazamox

4 - 2,4-D

(Refer to page 38)

Broadleaves:

The following weeds will be controlled up to the 4-leaf stage under a wide variety of conditions. Application at an earlier weed stage provides the best yield response.

Annual smartweed species (including lady's thumb)	Persian darnel
Annual sow-thistle	Ragweeds
Bluebur	Redroot pigweed
Cocklebur	Russian pigweed
Cow cockle	Russian thistle
Daisy fleabane	Shepherd's-purse
False flax	Stinging nettle
Flixweed	Stinkweed
Goat's-beard	Sweet clover
Kochia	Thyme-leaved spurge
Lamb's-quarters	Volunteer canola (all varieties)
Mustards (except dog and tansy)	Wild buckwheat
Narrow-leaved hawk's-beard	Wild mustard
Plantain	Wild radish
Prickly lettuce	Wild sunflower

The following weeds are suppressed up to the 4-leaf stage with Adrenalin SC when conditions are optimal for growth. Control may be reduced if weeds are found in high population densities, or are growing under stressful conditions (drought, cold, heat).

Biennial wormwood*	Knotweed
Blue lettuce*	Leafy spurge*
Chickweed, common	Mustards (dog and tansy)
Cleavers	Oak-leaved goosefoot
Curled dock*	Peppergrass, common
Dandelion*	Pigweed, prostrate
Groundsel, common	Pigweed, tumble
Gumweed*	Pineappleweed
Hawkweed	Purslane
Heal-all	Russian knapweed*
Hedge bindweed*	Sheep sorrel*
Hemp-nettle	Yellow rocket
Horsetail	

* Only seedlings of biennial and perennial weeds will be controlled. Established plants emerging from roots will not be controlled.

Adrenalin SC will give top growth control of the following weeds. Control may be reduced if weeds are found in high population densities, or are growing under stressful conditions (drought, cold, heat).

Bull thistle	Field bindweed
Burdock	Hoary cress
Buttercup	Perennial sow-thistle
Canada thistle	Tartary buckwheat

Rate:

0.4 L per acre (One case treats 40 acres).

Add a non-ionic surfactant (such as *Agral 90 Ag-Surf* or *Surf92*) at 0.25 L per 100 L of spray solution. Surfactant not included.

Use of an anti-foam agent is suggested.

DO NOT apply *Adrenalin SC* more than once or follow *Adrenalin SC* with any related product in the same year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE* coarse droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury will occur as a result of spray overlap. AVOID SPRAYER OVERLAP.

DO NOT spray if temperatures of +5°C or lower are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze the treated crop within 14 days of application or cut for hay within 42 days of application.

Preharvest Interval: DO NOT apply within 79 days of harvest.

Re-cropping: Winter wheat may be seeded 3 months after application. Barley, canaryseed, canola (all varieties), chick-pea, corn, field peas, flax, lentils, oats, sunflower, wheat (spring and durum) may be grown safely the year following an application. Condiment mustard may be grown the second season following *Adrenalin SC* application. Conduct a field bioassay the year before growing any other crop than those listed above.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Store in a cool, dry place above 5°C.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	11	11	11

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15-16.

Hazard Rating:



Caution – Causes eye and skin irritation.

For an explanation of the symbols used here see page 11.



Herbicide Group 14 - carfentrazone (Refer to page 38)

Company:

FMC of Canada (PCP# 28573)
Distributed by NuFarm Agriculture.

Formulations:

240 g/L carfentrazone formulated as an emulsifiable concentrate.

Container sizes - 0.6 L, 1.2 L, 3.38 L

Crops, Rates and Staging:

Pre-Seeding:

From 14.8 to 29.5 mL per acre prior to the seeding of:

Sorghum

From 14.6 to 46.8 mL per acre prior to the seeding of:

Barley	Mustard
Buckwheat	Oats
Canola (rapeseed)	Rye
Chickpea	Safflower
Corn (field and sweet)	Soybean
Dry bean	Sunflower
Field peas	Triticale
Flax	Wheat (including spring, winter and durum)
Lentil	
Millet (pearl and proso)	

Use Agral 90 or Ag-Surf at 0.25 L per 100 L of spray solution or use Merge at 1 L per 100 L of spray solution.

Harvest aid treatment*:

CROP	RATE (mL per acre)
Barley, oats, wheat, millet, dry bean, chickpea, field pea	29 to 47
Sorghum	29
Potato**	94 to 142

Use Agral 90 or Ag-Surf at 0.25 L per 100 L of spray solution or use Merge at 1 L per 100 L of spray solution.

* DO NOT apply to crops if grown for seed purpose.

** A second application of 94 to 123 mL per acre may be applied in potato.

Weeds, Rates and Staging:

Apply to listed weeds up to ten (10) cm in height unless otherwise indicated:

WEEDS	RATE (mL per acre)
Redroot pigweed	14.6
Above weeds plus: Lamb's-quarters, Round-leaved mallow, Hairy nightshade, Stinkweed, Pigweed (Prostrate, Smooth, Tumble), Purslane, Tansy mustard, Tall waterhemp	23.2
Above weeds plus: Cocklebur, Kochia, Volunteer canola (all varieties), Eastern black nightshade,	29.2
Above weeds plus: Prickly lettuce, Corn spurry	46.8

Application Information:

Water Volume: Use a minimum of 40 L per acre. Higher spray volumes is required for dense weed stands. Weed control improves with the amount of coverage.

Nozzles and Pressure: Maximum 35 psi (210 kPa) if using conventional nozzles. Low drift nozzles may require higher pressure for proper performance. Use nozzles and pressure designed to deliver proper coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect activity. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms may be reduced as weeds hardened off by drought are less susceptible.

Tank Mixes:

Herbicides:

Pre-plant:

*Glyphosate** (0.5 to 1L per acre of 360 g/L formulation)

2,4-D Ester-700 (0.32 L per acre) (fallow only)

Harvest aid treatment:

*Glyphosate** (1L per acre of 360 g/L formulation)

*Reglone*** (0.7 to 0.9 L/acre)

* IPA or K salt only.

** For potatoes only. If this mix is applied, neither the mix, nor the individual products may be applied following the first application of the mix.

Note: The above mixes are those listed on the *Aim* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall within 6 to 8 hours after application may reduce activity. Heavy rainfall shortly after application may reduce activity.

Re-entry: DO NOT enter treated fields for 12 hours.

Preharvest Interval: Leave 7 days between application and harvest for potatoes and 3 days for all other registered crops for harvest aid uses.

Grazing: DO NOT graze the treated crop or cut for feed.

Re-cropping: There are no rotational restrictions on crops registered for pre-seed use. All other crops may be planted 12 months after application.

Aerial application: DO NOT apply by air.

Storage: Store in a cool, dry place and avoid excess heat.

Buffer Zones:

Application method	Buffer Zones (metres) Required for the Protection of Terrestrial Habitat
Ground only*	5

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches above the crop canopy.

DO NOT apply in areas where surface water from the treated area can run off into aquatic habitats.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution – Eye and skin irritant

For an explanation of the symbols used here see page 11.

Altitude FX

Herbicide Group

2 - imazamox

4 - fluroxypyr & MCPA

(Refer to page 38)

Company:

BASF Canada

Formulations:

Altitude FX contains 3 separate components. Each case contains:

AC 299,263 120 AS (PCP# 26705): 120 g/L imazamox formulated as a solution.

Container size - 1.34 L

Starane (PCP# 24815): 180 g/L of fluroxypyr formulated as an emulsifiable concentrate.

Container size - 4.8 L.

MCPA ester 600 (PCP# 27802): 600 g/L of MCPA ester formulated as an emulsifiable concentrate.

Container size - 7.5 L.

Crops and Staging:

CLEARFIELD wheat varieties: 3 leaf (after appearance of first tiller) to 6 leaf stage to ensure optimal crop tolerance. Apply only to CLEARFIELD wheat varieties; application to any other variety of wheat or any other crop will result in crop death.

Weeds and Staging:

Grasses:

Apply from 1 to 4 leaves, up to a maximum of two tillers.

Barnyard grass	Volunteer cereals (barley, canary seed, oat, non-CLEARFIELD spring wheat, durum)
Foxtail (green and yellow)	
Japanese brome (suppression)	
Persian dandel	Wild oat

Broadleaves:

Apply up to 4 leaf stage unless otherwise indicated.

Annual sunflower	Prickly lettuce
Burdock common	Redroot pigweed
Chickweed	Round-leaved mallow*
Cleavers (1 to 4 whorls)	Russian thistle*
Cocklebur	Shepherd's-purse
Common ragweed	Stinkweed
Cow cockle	Stork's-bill (1 to 8 leaf)*
Flixweed	Vetch
Green smartweed	Volunteer canola (all varieties)
Hemp-nettle (2 to 6 leaf)	Volunteer flax (1 to 12 cm)
Kochia	Wild buckwheat
Lamb's-quarters	Wild radish
Mustards (except dog and tansy)	

* Suppression

Rates:

Altitude FX (1 case treats 20 acres)

AC 299,263 120 AS: 67 mL per acre.

Starane: 0.24 L per acre.

MCPA 600 Ester: 0.38 L per acre.

Add a non-ionic surfactant (such as *Agral 90 Ag-Surf* or *Surf 92*) at 0.25 L per 100 L of spray solution. Surfactant not included.

DO NOT apply *Altitude FX* or other products containing the same active ingredients imazamox, fluroxypyr or MCPA more than once per season.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 20 to 40 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of **ASABE** coarse droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury will occur as a result of spray overlap. **AVOID SPRAYER OVERLAP.**

DO NOT spray if temperatures of +5°C are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

None Registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze the treated crop within 14 days of application or cut for hay within 42 days of application.

Preharvest Interval: DO NOT apply within 79 days of harvest.

Re-cropping: Winter wheat may be seeded 3 months after application. Barley, canola (all varieties), field peas, flax, lentils, oats, sunflower, and spring wheat may be grown safely the year following application. Condiment mustard may be grown the second season following *Altitude FX* application. Conduct a field bioassay the year before growing any other crop than those listed above.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Store in a cool, dry place above 5°C. Combustible – DO NOT store near heat or open flame.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	15	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15-16.

Hazard Rating:**Starane**

Danger - Poison



Warning - Eye Irritant



Caution - Skin Irritant

MCPA ester

Warning - Poison

For an explanation of the symbols used here see page 11.

Altitude FX2**Herbicide Group**

2 - imazamox

4 - fluroxypyr

(Refer to page 38)

Company:

BASF Canada

Formulations:

Altitude FX2 contains 2 separate components. Each Case treats 40 acres and contains:

AC 299,263 120 AS (PCP# 26705): 120 g/L imazamox formulated as a solution.

Container size - 2.68 L

Starane (PCP# 24815): 180 g/L of fluroxypyr formulated as an emulsifiable concentrate.

Container size - 9.6 L.

MCPA, 2,4-D, or Curtail M must be added and are purchased separately.

Crops and Staging:

CLEARFIELD wheat varieties: 3 leaf (after appearance of first tiller) to 6 leaf stage to ensure optimal crop tolerance. Apply only to CLEARFIELD wheat varieties; application to any other variety of wheat or any other crop will result in crop death.

Weeds and Staging:**Grasses:**

Apply from 1 to 4 leaves, up to a maximum of two tillers.

Barnyard grass

Foxtail (green and yellow)

Japanese brome (suppression)

Persian dandel

Volunteer cereals (barley, canary seed, oat, non-CLEARFIELD spring wheat, durum)

Wild oat

Broadleaves:

Apply up to 4 leaf stage unless otherwise indicated.

Cleavers (1 to 4 whorls)

Cow cockle

Green smartweed

Kochia

Lamb's-quarters

Redroot pigweed

Round-leaved mallow*

Russian thistle*

Shepherd's-purse

Stinkweed

Stork's-bill (1 to 8 leaf)*

Volunteer canola (except Clearfield varieties)

Volunteer flax (1 to 12 cm)

Wild buckwheat

Wild mustard

* Suppression

Rates:

Altitude FX2 (1 case treats 40 acres)

AC 299,263 120 AS: 67 mL per acre.

Starane: 0.24 L per acre.

Altitude FX2 must be tank mixed with one of the registered tank mix options found under the "tank mix" section below. Add a non-ionic surfactant (such as *Agral 90* *Ag-Surf II* or *Surf 92*) at 0.25 L per 100 L of spray solution. Surfactant not included.

DO NOT apply *Altitude FX2* or other products containing the same active ingredients *imazamox*, or *fluroxypyr* more than once per season.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 20 to 40 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE* coarse droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Initial crop injury may be observed after application but this is outgrown and should not affect yield. Severe crop injury will occur as a result of spray overlap. AVOID SPRAYER OVERLAP.

DO NOT spray if temperatures of +5°C are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

Herbicides:

Altitude FX2 must be mixed with one of the following:

MCPA Ester 600 (0.38 L/acre)

2,4-D Ester 700 (0.32 L/acre)

Curtail M (0.61 to 0.81 L/acre)

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze the treated crop within 14 days of application or cut for hay within 42 days of application.

Preharvest Interval: DO NOT apply within 79 days of harvest.

Re-cropping: Winter wheat may be seeded 3 months after application. Barley, canola (all varieties), field peas, flax, lentils, oats, and spring wheat may be grown safely the year following application. Condiment mustard may be grown the second season following *Altitude FX2* application. Conduct a field bioassay the year before growing any other crop than those listed above.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Store in a cool, dry place above 5°C. Combustible – DO NOT store near heat or open flame.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	15	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15-16.

Hazard Rating:

Starane



Danger – Poison



Warning – Eye Irritant



Caution – Skin Irritant

For an explanation of the symbols used here see page 11.

Amitrol 240

Herbicide Group

11 - amitrole

(Refer to page 38)

Company:

Nufarm Agriculture (PCP#25684)

Formulation:

231 g/L amitrole formulated as a liquid.

Container size - 10 L.

Crops, Rates and Staging:

Fall Stubble: Perennial weed control prior to spring seeding. No planting restrictions for barley, canola*, field corn, field pea, soybean, wheat, or white bean, but leave 8 months between application and the seeding of any other crops.

Alfalfa stand renovation/removal: 4 to 6 inches (10 to 15 cm) high.

Pastures (spot treatment only): For non-selective patch treatment of dandelion, Canada thistle, perennial sow-thistle, hoary cress, milkweed, poison ivy and toadflax apply 0.165 L in 25 L of water to treat a 10 m x 10 m area. For treatment of leafy spurge and horsetail, apply 0.460 L in 25 L of water to treat a 10 m x 10 m area. DO NOT mow treated plants for 3 weeks after application.

Established shelterbelts: up to 11.3 L per acre - Keep spray away from tree foliage or trunks.

Pre-seeding:

CROP	RATE (L per acre)	Delay seeding after application
Barley, wheat, canola	1.7	0 to 1 days
Field pea	1.7	5 to 7 days
Soybean (low rate)	1.7	6 days
Soybean (high rates)	5.1 to 6.7	10 to 14 days
Field corn, white bean, soybean	3.3 to 6.7	10 to 14 days

Adhere to the maximum pre-seeding rates and delays between application and seeding indicated above to avoid the risk of damage to the emerging crop or excess residues in the grain.

Avoid using rates higher than 6.7 L per acre for preplant applications prior to soybeans, dry beans and corn on very light textured soils with low organic matter, as crop damage can occur.

Fallow areas: Apply according to weed stage and rates in the next section.

Weeds, Rates and Staging:

Fall stubble: Canada thistle, perennial sow-thistle - Spray when thistle has 4 to 6 inches (10 to 15 cm) of new growth. DO NOT cultivate for 2 weeks after application. DO NOT apply after October 1. DO NOT replant crops in treated areas within 8 months of application except those registered for pre-seeding uses.

Pre-seeding: Dandelion and annual weeds Apply 1.7 L per acre to actively growing weeds less than 10 cm tall or across. DO NOT cultivate for 10 to 14 days after treatment.

Fallow, Pastures and Shelterbelts:

WEED	RATE (L/ACRE)	WEED STAGE
Canada thistle	5.1 to 6.7	Early bud to bloom stage.
Cattails	15.2 to 18.5	After seed heads have formed.
Dandelion	1.7 to 5.1	Young and actively growing plants.
Hoary cress	7.6 to 11.3	Advanced rosette and bud stage.
Horsetail	5.0 to 6.7	Actively growing plants.
Leafy spurge	15.2 to 18.5	Advanced flowering to early seed set.
Milkweed	7.6 to 11.3	Early summer after majority of shoot emergence.
Perennial sow-thistle	5.1 to 6.7	Early bud to bloom stage.
Poison ivy	3.7	Fully developed green foliage.
Quackgrass	5.1 to 6.7	When plants are 4 to 6 inches (10 to 15 cm) high and actively growing.
Toadflax	7.6 to 11.3	Advanced rosette to prebud.

Application Information:

Water Volume:

Fall stubble: 20 to 81 L per acre.

Pastures, shelterbelts: 40 to 121 L per acre. For poison ivy, apply 202 to 405 L per acre.

Pre-seeding: 20 to 81 L per acre.

Nozzles and Pressure: Maximum 45 psi (less than 300 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressures designed to deliver proper coverage with ASABE coarse droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Less than acceptable results may occur in dry weather.

Tank Mixes:

Nufarm also supports the following mixes that are not on the Amitrol 240 label. Mixes must be applied according to the most restrictive use limitations for either product:

Herbicides: glyphosate

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfall period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT re-enter treated areas for 12 hours.

Grazing: DO NOT graze treated crops or weeds or use for hay or feed.

Re-cropping: DO NOT plant any crop for 8 months following application except those registered for pre-seeding uses.

Aerial Application: DO NOT apply by air.

Storage: DO NOT store where temperatures may exceed 50°C or near open flames. Do not store below 4°C.

Buffer Zones: DO NOT contaminate any body of water. Use cautions to prevent spray, spray mist, or vapours from drifting off target. Spray drift may cause damage to crops or vegetation.

Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15-16.

Hazard Rating:



Caution – Poison



Warning – Eye Irritant

For an explanation of the symbols used here see page 11.

Ares

Herbicide Group
2 - imazamox & imazapyr
(Refer to page 38)

Company:

BASF Canada (PCP# 30188)

Formulations:

33% imazamox and 15% imazapyr formulated as a solution.

Container size - 1x 9.8L jug, Merge 8.1 L jug.

Crops and Staging:

CLEARFIELD canola: 2 to 7 leaf stage.

CLEARFIELD oilseed mustard (*Brassica juncea*): 2 to 7 leaf stage.

Weeds, Rates and Staging:

At 244 mL per acre (40 acres per case), *Ares* will control:
Grasses - From 1 to 6 true leaf stage with up to 2 tillers stage:

Barley grass
Green foxtail
Japanese brome*
Persian darnel

Volunteer cereals (barley, canaryseed, durum, oats and wheat - NOT including CLEARFIELD varieties)

Wild oats
Yellow foxtail

* Spring germinating Japanese brome maximum 4 leaf stage.

Broadleaf Weeds - cotyledon to 4 leaf stage unless otherwise indicated:

Cleavers (up to 4 whorls) ***	Shepherd's-purse
Cow cockle	Stinkweed
Green smartweed	Volunteer canola (not CLEARFIELD varieties)
Hemp-nettle	Volunteer tame mustard (not CLEARFIELD oilseed varieties - B. juncea)
Lamb's-quarters **	Wild buckwheat**
Redroot pigweed	Wild mustard ***
Round-leaved mallow	
Russian thistle	

** Cotyledon to 6 leaf stage.

*** NOT Group 2 resistant biotypes

Merge adjuvant must be used at a rate of 0.5 L per 100 L of spray solution.

DO NOT apply *Ares* or follow *Ares* with other products containing the active ingredient imazamox (e.g. *Solo*, *Odyssey*) in the same year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 20 to 40 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets. Use 50 mesh or coarser filter screens.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT spray if temperatures of +5°C are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

Herbicides:

In CLEARFIELD canola only. DO NOT apply to CLEARFIELD oilseed mustard:

Lontrel 360 (84 mL/acre)

In CLEARFIELD canola and CLEARFIELD oilseed mustard:

Equinox (54 to 81 mL/acre)

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *Ares* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Avoid application when heavy rain is forecast.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Preharvest Interval: DO NOT apply within 60 days of harvest of registered crops.

Re-cropping: Barley, canaryseed, chickpea, CLEARFIELD canola/oilseed B. juncea, field corn, field pea, lentil, oat, and spring wheat may be seeded the first full season after application. Non-Clearfield canola, durum, flax and sunflower may be seeded the second full season after application. The company recommends that a field bio-assay (a test strip grown to maturity) be conducted the year before growing any crops other than those listed above.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place.

Environment: Avoid drift. Leave at least 11 m between the downwind edge of the boom and sensitive areas such as shelterbelts, hedgerows, wetlands, woodlots, vegetated ditch banks, ponds, streams, and sloughs.

Tank Cleaning:

Refer to page 'Method C' in the general tank cleaning section on 15-16.

Hazard Rating:

No specific hazards.

Assert FL

Herbicide Group

2 - imazamethabenz & florasulam

4 - MCPA

(Refer to page 38)

This product is the equivalent of a prepackaged tankmix of *imazamethabenz* (page 198) and *florasulam* + *MCPA* (page 162). Information listed is restricted to Crop, Weeds and Rates. For other detailed information on the component products see the product pages listed above.

Company:

Nufarm Agriculture

Formulation:

The *Assert FL* package has 3 components:

Assert (PCP#21032): 300 g/L imazamethabenz formulated as a suspension concentrate.

Container size - 13.5 L

Frontline A (PCP#27029): 50 g/L florasulam formulated as a suspension concentrate.

Container size - 0.8 L

Frontline B (PCP#27030): 500 g/L MCPA Ester formulated as an emulsifiable concentrate.

Container size - 5.6 L

pH adjuster 3.12 kg

Crops, Rates and Staging:

Spring wheat (including durum) and barley in the 2 to 6 leaf stage.

Weeds and Staging:

Weeds controlled by *imazamethabenz* and florasulam + MCPA at the staging that is common to both component products.

Rates:

Assert: 0.65 L per acre;

Frontline A: 40 mL per acre;

Frontline B: 280 mL per acre.

One case treats 20 acres. Make only one application of this product or other product containing the same ingredients per year.

Restrictions:

Apply using application details that are common for both the component products and adhere to the most stringent restrictions of each.

Attain XC

(this reference text to be removed in the 2015 edition)

See *Fluroxypyr* + *2,4-D* on page 168.

Authority

Company:

FMC of Canada (PCP#29012)

Distributed by NuFarm Agriculture

Formulation:

480 g/L sulfentrazone formulated as a suspension concentrate.

Container sizes - 4 x 3.8L jugs per case.

Crops and Staging:

Chickpeas, Field Pea, Flax and Sunflower: Soil applied in the spring only.

Pre-plant surface: Apply to the soil surface prior to seeding the crop.

Pre-emergent surface: Apply to the soil surface up to 3 days after seeding. Crops emerging or near emerging at application may be injured.

All applications require rainfall for proper activation. (See "Effects of Growing Conditions")

DO NOT use on coarse soils classified as sand which have less than 1% organic matter or where water table is high.

Weeds and Staging:

Controls the following weeds when applied to the soil prior to emergence:

Kochia	Redroot pigweed
Lamb's-quarters	Wild buckwheat

Rates

Apply at 88 to 118 mL per acre (172 to 129 acres per case). Use the higher rates within the rate range for soils with pH less than 7.0 and organic matter greater than 3%.

Herbicide Group

14 - sulfentrazone

(Refer to page 38)

DO NOT APPLY to coarse-textured (sandy) soils with organic matter less than 1%, compacted or heavy clay soils with less than 1.5% organic matter, soils with organic matter content greater than 6%, or to soils with a pH of 7.8 or greater.

Application of *Authority* in back to back years is not recommended.

Application Information:

Water Volume: Minimum 40 L per acre

Nozzles and Pressure: Maximum 30 psi (175 kPa) if using conventional nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver proper coverage with ASABE medium droplets or larger.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

All applications require rainfall for proper activation. If weed growth begins before activation occurs, poor control may result on larger weeds. A moderate rainfall (10 to 20 mm) or equivalent irrigation is required within 10 to 14 days to activate pre-emergent surface treatments. If rain does not occur, a shallow harrowing or use of a rotary hoe may assist activation using existing soil moisture. Dry conditions that persist after any application may reduce weed control. On sandy soils, heavy rainfall following application may cause leaching of *Authority*, resulting in reduced weed control.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Rainfall following application is required for adequate weed control.

Grazing: DO NOT graze treated crops or cut for hay.

Preharvest Interval: Leave 60 days between application and harvest.

Re-entry: DO NOT re-enter treated area within 12 hours.

Re-cropping: Registered crops and soybeans may be planted anytime after application. Alfalfa, barley, field corn and spring, durum and winter wheat may be seeded the season following application (after one winter). Canola, sweet corn and sorghum may be seeded the second season (two winters) after application. Lentils may be seeded the third season following application (3 winters). For all other crops three winters must pass following application and a successful bioassay indicating adequate tolerance before planting. For each year of drought experienced, add one year to the intervals above and conduct a bioassay to confirm tolerance of the rotational crop. Lentils may be particularly sensitive to Authority residue.

Aerial Application: DO NOT apply by air.

Storage: Store above 5°C to keep from freezing. If frozen, and solid crystals are observed, warm to above 15 °C and shake or roll container periodically to dissolve solids.

Buffer Zones: DO NOT fill mix or clean sprayer within 15 m of any water source, unless the well is properly capped or activities take place on impervious pads or properly diked mixing/loading areas. Leave a 1m buffer between the last spray path and water or wetland habitats and 10 meters to sensitive plants and upland habitats when applying by ground. Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

Tank Cleaning:

After spraying and before using sprayer equipment for any other applications, thoroughly clean sprayer using the following procedure:

1. Drain sprayer tank, hoses, and spray boom. Use a high-pressure detergent wash to remove sediment and residues from the inside of the sprayer tank and thoroughly rinse. Then thoroughly flush all sprayer hoses, booms, and nozzles with clean water.
2. Prepare a sprayer cleaning solution by adding three litres of ammonia (containing at least 3% active) per 100 litres of clean water. Prepare sufficient cleaning solution to allow the operation of the spray system for a minimum of 15 minutes to thoroughly flush hoses, spray boom and spray nozzles.
3. Convenient and thorough cleaning of the sprayer can be achieved if the ammonia solution or fresh water is left in the spray tank, hoses, spray booms and spray nozzles overnight or during storage.
4. Drain the sprayer system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and all strainers and screens separately in an ammonia solution.
5. Properly dispose of all cleaning solution and rinsate in accordance with provincial guidelines and regulations.

For additional information, Refer to page 15 - 16.

Hazard Rating:



Caution - Poison.

For an explanation of the symbols used here see page 11.

Authority Charge

Herbicide Group

14 - carfentrazone & sulfentrazone

(Refer to page 38)

This product is a co-pack of *Aim* (page 77) and *Authority* (page 86). Information is restricted to Crop, Weeds, Rates and Tank Mixes. For other detailed information on the component products see the product pages listed above. *Note: this product is based on an unlabeled tank mix supported by the manufacturer.*

Company:

FMC Canada

Formulation:

Aim (PCP# 28573): 240 g/L carfentrazone formulated as an emulsifiable concentrate.

Container size - 1 x 1.2 L jug

Authority (PCP# 29012): 480 g/L sulfentrazone formulated as a suspension concentrate.

Container size - 2 x 3.79 L jug

Crops and Staging:

Chickpea, Field Pea, Flax and Sunflower:

Soil applied in the spring only, tank mixed with glyphosate.

Pre-plant surface: Apply to the soil surface prior to seeding the crop.

Weeds and Staging:

Weeds controlled by component products.

Rates

Aim: 15 to 18.75 mL per acre.

Authority: 88 to 118 mL per acre.

One case treats 80 to 64 acres at the respective rates above.

Tank Mixes:

Authority Charge should be tank mixed with glyphosate at 180 to 360 grams active ingredient per acre (0.5 to 1.0 L/acre of 360 g/L formulation See glyphosate page for equivalent rates for other formulations.) based on the *Aim* component label only. The *Authority* label does not list any registered tank mixes.

For additional information, precautions and restrictions see the individual component at the page numbers shown above.

Avadex Brands

Herbicide Group

8 - triallate

(Refer to page 38)

Company:

Gowan Canada

Formulation:

Extra Strength Avadex BW (PCP# 16759): 480 g/L triallate formulated as an emulsifiable concentrate.

Container size - 2x10 L to 1000 L.

Avadex MicroActiv (PCP# 25112): 10% triallate formulated as a granular. Container size - 22.7 kg, 451.3 kg.

Crops, Rates and Application Timing:

Avadex Liquid Rates – Spring Treatment

CROP	APPLICATION TIMING	RATE (L/acre)		ACRES TREATED PER 100 L CONTAINER	
		Organic Matter		Organic Matter	
		4% or less	Greater than 4%	4% or less	Greater than 4%
Spring and durum wheat	Before Seeding*	1.01	1.17	99	85.4
	After Seeding	1.17	1.41	85.4	70.9
Barley	Before and After Seeding	1.17	1.41	85.4	70.9
Canola, flax†, mustard	Before Seeding	1.41	1.86	70.9	53.7
Peas (dry)	Before Seeding	1.41	1.41	70.9	70.9

* Do not apply this product before seeding wheat in soils with 4 percent or less organic matter (brown, dark brown or grey wooded soils) where discers are to be used for seeding. If an air seeder is to be used, it must be equipped with a depth control device to ensure accurate seed placement, otherwise crop injury may occur.

† Excluding Solin (low linolenic acid flax).

Avadex Granular Rates – Fall Treatment

CROP	RATE (KG/ACRE)			ACRES TREATED PER 22.7 KG CONTAINER		
	Organic Matter			Organic Matter		
	Less than 2%*	2 to 4%	Greater than 4%	Less than 2%*	2 to 4%	Greater than 4%
Spring and durum wheat	4.45	5.67	6.88	5.1	4.0	3.3
Barley, canaryseed	4.45	5.67	6.88	5.1	4.0	3.3
Canola, flax†, mustard	5.67	6.88	8.90	4.0	3.3	2.6

* Fall treatments conducted under minimum tillage are not recommended on soils with less than 2 percent organic matter.

† Excluding Solin (low linolenic acid flax).

Avadex Granular Rates – Spring Treatment

CROP	APPLICATION TIMING**	RATE (KG/ACRE)		ACRES TREATED PER 22.7 KG CONTAINER	
		Organic Matter		Organic Matter	
		4% or less*	Greater than 4%	4% or less*	Greater than 4%
Spring and durum wheat	Before seeding***	4.45	5.67	5.1	4.0
	After seeding	5.67	6.88	4.0	3.3
Barley, canaryseed	Before and after seeding (barley only)	5.67	6.88	4.0	3.3
Canola, flax†, mustard	Before seeding	6.88	8.90	3.3	2.6

* Minimum tillage treatments must be applied to fields with at least 2 percent organic matter.

** Minimum tillage treatments must be applied 10 to 14 days before seeding or incorporating. For minimum tillage treat-

ments on spring and durum wheat, apply 5.67 kg per acre on soils with 4% organic matter or less and 6.88 kg per acre on soils with greater than 4 percent organic matter.

*** DO NOT apply this product before seeding wheat in soils with 4% or less organic matter (brown, dark brown or grey wooded soils) where discers are to be used for seeding. If an air seeder is to be used, it must be equipped with a depth control device to ensure accurate seed placement, otherwise crop injury may occur.

† Excluding Solin (low linolenic acid flax).

Seedling Forage Legumes (under-seeded only):

Apply recommended rates for the companion crop.

Alfalfa

Bird's-foot trefoil

Clover (alsike, red, sweet)

Weeds and Staging:

For pre-emergent control of wild oats.

Application Information:

Water Volume (Liquid formulations only):

45 L per acre.

Pressure: 30 psi (200 kPa), liquid formulation only.

Nozzles: Flat fan, liquid formulation only. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets.

General Information: The liquid formulation must be incorporated into soil that is free of lumps or trash. The liquid formulation is recommended for spring use because soils are left in an erosion prone state if the liquid is fall-applied. The granular formulation may be incorporated into trashy soil and is best suited for fall use.

Fall Applications (Conventional Tillage): Apply *Avadex* granules to fields that are in good working condition, without excessive trash. Heavy trash or lumpy, wet fields may require tillage prior to application. *Avadex* must be applied after October 1 but before soil freeze-up. Application before October 1 may result in reduced weed control. Only one incorporation is required in the fall. The second incorporation may be done in the fall (before soil freeze-up) or in the spring.

Fall Application (Minimum Tillage): Applications of *Avadex* granules should be made to standing stubble, chemical fallow, or summerfallow fields that are not prone to erosion. DO NOT apply to smooth, hard packed soils that may allow granules to drift. If excessive crop residue exists at the time of application, harrowing should be conducted to ensure the granules are in good contact with the soil. Apply when the soil begins to cool (less than 4°C) and within 3 weeks of soil freeze-up. Incorporation can be performed in the spring before seeding or as part of the seeding operation.

Spring Application (Conventional Tillage): Apply *Avadex* (liquid or granules) to fields that are in good working condition, without excessive trash. Heavy trash or lumpy, wet fields may require tillage prior to application. Liquid formulations should be applied to fields with 30 percent or

less trash cover. *Avadex* may be applied before or after seeding of wheat, barley, or canaryseed and before seeding of canola, flax, mustard or peas (liquid only). If wheat is being seeded into soils with an organic matter content of less than 4 percent, *Avadex* should be applied after seeding.

Spring Application (Minimum Tillage): *Avadex* granules should be applied in spring and when the soil temperature is 4°C or less. Apply granules 10 to 14 days before incorporation. DO NOT apply more than 4 weeks before seeding is intended.

Incorporation:

Conventional Tillage: *Avadex* applications require two incorporations, with the second incorporation at right angles to the first. Using a seeder that provides soil disturbance equivalent to a cultivator may replace one of the incorporations. The first incorporation of the granular formulation should be completed within 48 hours of application and the second incorporation should be delayed an additional 48 hours or more. The first incorporation of the liquid formulation should be completed as soon as possible after spraying, while the second incorporation may be done any time prior to crop emergence.

Incorporate to a depth of 2 inches (5 cm) by setting disc or cultivator implements to cut 3 inches (7.5 cm) into the soil. Mixing the product to greater depths will dilute the herbicide, decrease wild oat control, and may cause injury to cereals. Ensure that cereals are seeded below the treated layer (2 to 3 inches or 5 to 7.5 cm). Incorporations performed after seeding should be conducted with harrows or other suitable tillage equipment adjusted so as not to disturb the seed. Harrowing does not provide effective incorporation if compact soil prevents penetration of harrow teeth, if trash accumulates in the harrow sections, or if the harrows bounce.

Minimum Tillage: Incorporation of *Avadex* granules in minimum tillage systems is achieved with one high disturbance incorporation, which can be conducted prior to seeding, or as part of the seeding pass. A high disturbance system is one that disturbs the soil enough so that emerged weeds are controlled by the operation (example - air seeder with cultivator shovels). Harrowing after the incorporation operation is recommended for best results.

For optimum results in minimum tillage systems, incorporate when wild oat growth is noticeable in the field, as this will ensure that the soil is warm enough for activation of *Avadex*.

Under excessively warm or wet conditions between application and crop emergence, control may be reduced. For best results on heavy wild oat infestations, use the conventional tillage guidelines for incorporation.

Summer Fallow: Incorporation can be done by a disc followed by harrowing at right angles, a vibrashank cultivator followed by harrowing at right angles, or double harrowing. The second operation can be delayed until spring. If summerfallow must be ridged to prevent soil erosion the granular formulation should not be used in the fall. Note that fall minimum tillage applied granules do not require incorporation in the fall. If soils must be ridged following application of the liquid formulation, ridging depth should be kept to a minimum as deep ridging may reduce wild oat control and increase crop injury.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Reduced control may result if prolonged cool conditions or dry soil conditions prevail at the time weeds are emerging. If conditions are dry or wild oats germinate from below the treated zone, the weeds may emerge, but will usually be controlled. Thinning of wheat can occur under conditions of heavy rainfall or if cold soil conditions persist as the crop emerges.

Tank Mixes:

Herbicides: *Avadex* liquid may be tank mixed with liquid formulations of trifluralin for control of wild oats, green and yellow foxtail in wheat and barley. Apply after seeding but prior to crop emergence. Consult the recommendations for trifluralin for rates in different soil types.

Insecticides: None registered.

Fertilizer: *Avadex* liquid alone, or tank mixed with liquid formulations of trifluralin, may be tank mixed with liquid fertilizer. Compatibility of the herbicide and liquid fertilizer should be checked. Follow the instructions on the herbicide label prior to adding the herbicide to the spray tank.

Avadex liquid may be sprayed on dry urea fertilizer. A minimum of 150 kg/ha (60 kg/acre) of dry urea fertilizer must be used. Only commercial blending is recommended.

Note: The above mixes are those listed on the *Avadex* labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: At least 0.5 inches (1.5 cm) within 2 weeks of application is required for activation.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze the treated crop or use as hay or feed prior to crop maturity or in year of treatment.

Re-cropping: DO NOT seed tame oats the year after treatment.

Aerial Application: DO NOT apply by air (*Extra Strength Avadex BW*). Granular formulations may be applied by air with attachments designed for applying low volumes of granules (*Avadex MicroActiv*).

Storage: DO NOT freeze liquid formulations. Store granular formulations in a cool, dry place.

Buffer Zones: (Liquid formulations only)

Application method	Buffer Zones (metres †) Required for the Protection of:			
	Aquatic Habitats of Depths			Terrestrial habitat
	Less than 1 m	1 to 3 m	Greater than 3 m	
Ground only*	5	2	1	5

See page 29 for an explanation of the different habitats.


* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.


† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

 Warning – Poison (Liquid formulation)

 Warning – Contains the allergen soy (Liquid and Granular)

Skin and Eye Irritant (Granular formulation)

For an explanation of the symbols used here see page 11.

Avenge* 200 C

Herbicide Group
8 - difenzoquat
(Refer to page 38)

Company:

AmVac Corporation, distributed by Syngenta
 (PCP#12853)

Formulation:

200 g/L difenzoquat formulated as a solution.
 Container size - 2 x 10 L, 113 L.

Note: This formulation is no longer manufactured, but some supplies may still remain in the distribution system. This product may be removed from future editions of this publication.

Crops and Staging:

Note: Import tolerances (Maximum Residue Limits) for Avenge residues in cereal crops have been revoked by the US-EPA. Cereal crops intended for export to the USA that are treated with Avenge may experience difficulty with export approvals to the USA. Producers should check with their commodity buyer before application.

Apply up the 6 leaf stage of cereal and grass crops:

Barley
 Canaryseed
 Fall rye (Cougar, Frontier, Kodiak, Puma, Rymin only)
 Triticale (Carman and Welsh)
 Wheat (AC Barrie, Biggar, Bluesky, CDC Makwa, CDC Teal, Columbus, Conway, Cutler, Fielder, Genesis, Glenlea, Katepwa, Lancer, Leader, Neepawa, Oslo, Pasqua, Selkirk and Wildcat wheat varieties).
 Winter wheat (Norstar only)

Forages* (only when underseeded to wheat and barley varieties listed above):

Alfalfa	Orchardgrass
Bird's-foot trefoil	Red clover
Bromegrass	Reed canarygrass
Creeping red fescue	Russian wildrye
Crested wheatgrass	Sweet clover
Kentucky bluegrass	Timothy
Meadow fescue	

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Seedling grasses* (apply up to the 3 leaf stage of wild oat):

Meadow bromegrass	Streambank wheatgrass
Meadow foxtail	Tall fescue
Pubescent wheatgrass	Tall wheatgrass

* Grasses may show damage in year of application but will recover the following year.

Weeds and Staging:

Wild oats at the 3 to 5 leaf stage.

Rate:

Apply 1.42 to 1.72 L per acre. Use the higher rate if wild oat population exceeds 200 wild oats per square metre.
 One 20 L container treats 14 to 11 acres.

Application Information:

Water Volume:

Ground application: 40 L per acre.

Aerial application: 8 to 20 L per acre

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles directed 45° forward. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop injury is worse on cold or hot, humid days than when temperatures are moderate. Best weed control is attained when temperatures are 20 to 30°C, particularly when these temperatures follow application. Do not spray when crop or weeds are wet with heavy dew or rain. Do not apply when crop is under stress from drought, excessive moisture, or heat.

Tank Mixes:

Herbicides:

In wheat and barley:

2,4-D ester (up to 0.45 L/acre of 500 g/L formulation).

DO NOT use amine formulations.

2,4-DB (see 2,4-DB section for rates). May be used on cereals underseeded to forages.

Ally (2 to 3 g/acre)*

Bromoxynil/MCPA (recommended rates)

Curtail M (0.8 L/acre)

Dichlorprop/2,4-D (0.71 L/acre)

MCPA ester (up to 0.36 L/acre of 600 g/L formulation).

DO NOT use amine formulations.

Pardner (0.4 to 0.48 L/acre)

Thumper (0.4 L/acre)

In canaryseed:

Bromoxynil/MCPA (*Buctril M*, *Brotex*)(label rates)

MCPA ester (0.32 to 0.36 L/acre of 600 g/L formulation)

Pardner (0.4 L/acre)

*No additional adjuvant required.

Insecticides: None registered

Fungicides: None registered.

Fertilizers: None registered.

Allow a 5-day interval between the application of *Avenge* and other pesticides.

Note: The above mixes are those listed on the *Avenge* label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Syngenta also supports the following mixes that are not on the *Avenge* label. Mixes must be applied according to the most restrictive use limitations for either product:

Herbicides: *Attain XC*; *Broadside*; *Frontline XL*; *Infinity*; *OcTtain XL*; *Prestige XC*; *Refine SG*; *Refine SG + MCPA Ester*; *Retain*; *Spectrum*; *Triton C*.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours of application will reduce control.

Re-entry: DO NOT enter treated fields for at least 48 hours.

Grazing: DO NOT graze or feed treated cereal crop for 8 weeks after treatment. Do not graze or feed treated forages in the year of treatment. Straw from treated fields can be fed to livestock 8 weeks after application.

Preharvest Interval: DO NOT apply beyond the recommended crop stage.

Re-cropping: No restrictions the year after treatment.

Aerial Application: May be applied by air.

Storage: May be frozen.

Buffer Zones: DO NOT contaminate water bodies (sloughs, dugouts, lakes, streams, ponds, etc.).

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Warning - Poison

For an explanation of the symbols used here see page 11.

Axial*/Axial BIA

Herbicide Group

1 - pinoxaden

(Refer to page 38)

Company:

Syngenta Canada

Formulation:

Axial (PCP#28150)*: 100 g/L pinoxaden formulated as an emulsifiable concentrate.

Container size - 9.7 L or 77.6 L container.

Adigor Adjuvant (PCP#28151)*: 11.3 L or 90.4 L container.

-or-

Axial BIA (PCP#30341): 50 g/L pinoxaden formulated as an emulsifiable concentrate.

Container size - 2x10 L, 80 L, 400 L.

* Note: This formulation is no longer manufactured, but some supplies may still remain in the distribution system. This product may be removed from future editions of this publication.

Crops and Staging:

Spring wheat (not including durum), and barley. Up to the emergence of the flag leaf.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates, and Staging:

Apply from the 1 to 6 leaf up to the emergence of the 4th tiller.

Axial at 0.24 L/acre* plus Adigor adjuvant at 0.28 L per acre or Axial BIA at 0.48 L per acre (no adjuvant required)* (one case treats 40 acres):

Foxtail (green and yellow)	Volunteer oat
Barnyard grass	Volunteer canary seed
Proso millet	Wild oat

Apply at the 2 to 3 leaf stage for optimum control.

Optimum weed control and yield response occurs when weeds are controlled before tillering.

* Maximum one application per year of this or other products containing the active ingredient pinoxaden. Do not mix with any other adjuvant other than what is provided in the package.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre.

Aerial: 12 L per acre

Nozzles and Pressure: 40 to 45 psi (275 to 310 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury may result.

Weed control may be reduced if Axial is applied under stress conditions such as drought, heat, insufficient fertility, flooding or prolonged cool temperatures.

Tank Mixes:

Herbicides:

Tank-mix Partner*	Product Rates
Buctril M [†]	0.4 L/acre
Curtail M [†]	0.6 to 0.81 L/acre
Frontline XL	0.5 L/acre
Infinity	0.33 L/acre
MCPA ester [†]	0.34 to 0.44 L/acre (500 formulation)
Mextrol 450M [†]	0.5 L/acre
Refine SG**	12 g/acre
Refine SG** + MCPA ester*** [†]	12 g/acre + (0.23 to 0.28 L/acre)
Spectrum*** [†]	20 acres per case
Stellar	40 acres per case
Trophy [†]	20 acres per case

* Always consult the label of the broadleaf herbicide prior to use.

** Addition of surfactants other than those provided with Axial or Axial BLA are not required.

*** Suppression only of green foxtail.

[†] A reduction in barnyard grass control may be observed.

Insecticides: None registered.

Fungicides: Tilt (label rates).

Fertilizers: None registered.

Note: The above mixes are those listed on the Axial label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Syngenta also supports the following mixes that are not on the Axial label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Barricade II; Broadside; Momentum + MCPA ester; Prestige XC; Pulsar; Pulsar+Express SG (up to 6 g/acre); Pulsar (low rate)+MCPA ester; Refine SG + Perimeter; Stellar; Triton C; Triton C + MCPA^{††}

Fungicides: Propel, Quilt

^{††} Only wild oat is controlled with Axial.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of treatment may reduce control.

Re-entry: DO NOT enter treated fields for 12 hours.

Preharvest: Leave at least 60 days between treatment and harvest of grain and straw.

Grazing: DO NOT graze livestock within 7 days or cut for hay within 30 days of application.

Re-cropping: No restrictions the year following treatment. DO NOT seed any crops in the year of treatment following application (emergency re-crop).

Aerial Application: May be applied by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of: Terrestrial habitat
Ground only*	1
Aerial by airplane or helicopter	25

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Buffers are not required for handheld and backpack applications.

Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15-16.

Hazard Rating:



Caution – Poison



Warning – Eye and skin irritant

For an explanation of the symbols used here see page 11.

Axial iPak

This product is a prepackaged tank mix of *Axial* (page 93) and *Infinity* (page 205). Information listed is restricted to Crop, Weeds and Rates. For other detailed information on the component products see the product pages listed above.

Herbicide Group

1 - pinoxaden
6 - bromoxynil
27 - pyrasulfotole
(Refer to page 38)

Company:

Syngenta Canada

Formulation:

The Axial iPak package may contain either:

Axial (PCP# 28150)*: 100 g/L pinoxaden formulated as an emulsifiable concentrate.

Container size - 1 x 4.85 L jug.

-and-

Adigor Adjuvant (PCP# 28151): 1 x 5.65 L jug.

-or-

Axial BIA (PCP# 30341): 50 g/L pinoxaden formulated as an emulsifiable concentrate.

Container Size - 1x10L

-plus-

Both versions contain:

Infinity (PCP# 28738): 37.5 g/L pyrasulfotole and 210 g/L bromoxynil formulated as an emulsifiable concentrate.

Container size - 1 x 6.7 L jugs per case.

* NOTE: this formulation is no longer manufactured, but some supplies may still remain in the distribution system. This product may be removed from future editions of this publication.

Weeds and Staging:

Weeds controlled by the component products.

Rates:

Axial: 0.24 L per acre*

-plus-

Adigor adjuvant: 0.28 L per acre (add last)

-or-

Axial BIA: 0.48 L per acre (no adjuvant required)*

-plus-

Infinity: 0.33 L per acre.

One case treats 20 acres

Note: Ammonium sulphate is required to be added at 202 g per acre (99% dry) or 0.4 L per acre (49% solution) at 4 to 6 whorls for certain weeds controlled by *Infinity*

* Maximum one application of products containing the active ingredients pinoxaden, pyrasulfotole or bromoxynil per year. DO NOT mix with any other adjuvant other than what is provided in the package.

For additional information, precautions and restrictions, see the individual component at the page numbers shown above

Crops and Staging:

Spring wheat (not including durum), and barley. Up to the emergence of the flag leaf.

Axial Xtreme

Herbicide Group

1 - pinoxaden

4 - fluroxypyr

(Refer to page 38)

Company:

Syngenta (PCP#30391)

Formulation:

50 g/L pinoxaden and 87.5 g/L fluroxypyr formulated as an emulsifiable concentrate.

Container size - 2x 10L, 80 L, 400L.

Crops and Staging:

Spring wheat (NOT including durum) and barley:

1 to 6 leaf stage prior to the emergence of the 4th tiller, and before the first node can be felt in the stem.

Weeds and Staging:

Grasses - 1 to 6 leaf prior to the emergence of the 4th tiller

Foxtail (green and yellow)	Volunteer oat
Barnyard grass	Volunteer canary seed
Proso millet	Wild oat

Broadleaf Weeds - stages indicated below:

Cleavers (up to 4 whorls)	Stork's bill (up to 6 leaf)*
Kochia (2 to 8 leaf)	Wild buckwheat (up to 4 leaf)*

* Suppression.

Rates:

0.48 L per acre.
(40 acres per case)

DO NOT apply *Axial Xtreme* more than once or follow it with any other products containing pinoxaden or fluroxypyr in the same year.

Application Information:

Water Volume: 20 to 40 L per acre.

Nozzles and Pressure: Maximum 40 to 45 psi (275 to 310 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper

performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse classification* droplets. Use 50 mesh or coarser filter screens. Thorough coverage of the plants is essential for consistent control.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Tolerance and efficacy is best when applied during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under prolonged stress caused by excessive cool or heat, flooding or drought, or poor fertility, control of some weeds may be reduced and or crops may be injured.

Tank Mixes:

Herbicides:

Bromoxynil/MCPA (label rates) †

Curtail M (0.61 to 0.81 L/acre) †

Frontline XL (0.5 L/acre) ††

Infinity (336 mL/acre) ††

MCPA ester (0.28 to 0.37 L/acre – 600 g/L form) †

Refine SG (12 g/acre)

Refine SG + MCPA ester (rates above) †

† A reduction in barnyard grass control may be observed with this mix.

†† A reduction in green foxtail control may be observed with this mix.

Fungicides: *Tilt* (101 to 202 mL/acre)

Insecticides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Axial Xtreme* label only. To check for other possible mixes see the blue fold out chart in the back inside cover of the guide.

Syngenta also supports the following mixes that are not on the *Axial* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Broadside*

Fungicide: *Quilt*

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Preharvest: DO NOT apply within 60 days of harvest.

Grazing: Must not be grazed within 7 days or cut for live-stock feed within 30 days of treatment.

Re-cropping: Barley, canola, flax, forage grasses, lentils, mustard, oats, peas, rye or wheat may be seeded the first full season after application or fields can be summerfallowed.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. DO NOT freeze.

Environment: Avoid drift. Leave at least 15 m between the downwind edge of the boom and sensitive areas such

as shelterbelts, hedgerows, wetlands, woodlots, vegetated ditch banks, ponds, streams, and sloughs. Buffer zones can be reduced by 70% when using shrouds or by 30% when using cones mounted less than 12 inches from the crop canopy.

Tank Cleaning:

Refer to 'Method B' on page 15. Use 500 g or mL per 100L of rinsate for alkali detergents or 250 g or mL per 100L of rinsate for concentrated laundry detergents. DO NOT use chlorine based cleaners.

Hazard Rating:



Warning – Eye and skin irritant

Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Barricade II

Barricade II is a tank mix of Barricade SG with fluroxypyr (Perimeter II).

Herbicide Group

2 - thifensulfuron & tribenuron

4 - fluroxypyr

(Refer to page 38)

Company:

E. I. duPont Canada

Formulation:

The *Barricade II* package contains two components:

Barricade SG (PCP#29544): 25% thifensulfuron methyl plus 25% tribenuron methyl formulated as a water soluble granule.

Container size - 486 g bottle.

-plus-

Perimeter II (PCP#30094): 333 g a.e./L fluroxypyr formulated as an emulsifiable concentrate.

Container size - 3.4 L

Crops and Staging:

Barley and spring wheat (including durum) from 2 leaf until first node can be felt at the base of the stem.

Weeds and Staging:

Unless otherwise noted below, apply to young and actively growing weeds that are less than 4 inches (10 cm) in height or width.

Weeds Controlled:

Annual smartweed (green, lady's-thumb)	Lamb's-quarters
Canada thistle (less than 6 inches (15 cm) tall or across and prior to budding)**	Night-flowering catchfly
Cleavers (1 to 4 whorls)	Narrow-leaved hawk's-beard
Common chickweed (1 to 6 leaf)	Redroot pigweed
Cow cockle	Russian thistle
Flixweed	Stinkweed
Hemp-nettle	Stork's-bill (1 to 6 leaf)
Kochia	Volunteer canola (not CLEARFIELD varieties)
	Wild buckwheat
	Wild mustard
	Volunteer flax (up to 12 cm)

** Suppression only.

Rate:

Barricade SG: 12 g per acre.

-plus-

Perimeter II: 85 mL per acre.

(one package treats 40 acres)

Maximum of one application of this or other thifensulfuron/tribenuron products or fluroxypyr products per year.

Add Agral 90, Agsurf II, or Citowett Plus at 0.2 L per 100 L of spray solution. Barricade SG may degrade if left in the sprayer for an extended period. Apply within 24 hours of mixing.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum 40 L per acre.

Nozzles and Pressure: Flat fan nozzles are recommended. Sprayers without drift reduction nozzles should use between 30 to 40 psi (210 to 275 kPa). Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to registered crops that are stressed by severe weather conditions (frost, drought or water saturated soil) as crop injury may result. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Under certain conditions (heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) lightening in crop colour and reduction in crop height may occur.

Tank Mixes:

Herbicides: None registered

Fertilizers: None registered.

Note: The above mixes are those on the Barricade II labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

E.I. duPont also supports the following mixes that are not on the Barricade II label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Horizon NG, Axial, Assert, Flucarbazone + 2,4-D Ester, MCPA ester (190 mL/acre), Puma Advance, Simplicity, Traxos, Varro.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Up to 1 inch within 1 hour of application may reduce control.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Preharvest Interval: Leave 60 days between application and harvest.

Grazing: MUST NOT be grazed or fed to livestock for 7 days after treatment.

Re-cropping: Barley, canola, flax, forage grasses, lentils, mustard, oats, peas, rye or wheat or fields can be summer-fallowed the year after treatment.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. Avoid freezing. If frozen, bring to room temperature and agitate before use. This product is COMBUSTIBLE. DO NOT store near heat or open flame.

Buffer Zones:

Application method	Buffer Zones (metres †) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground*	15	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Follow the tank cleaning instructions on the thifensulfuron/tribenuron page. The addition of a wetting agent (detergent) will also aid the cleaning process.

Refer to page 15 to 16 for additional information.

Hazard Rating:



Danger - Poison



Warning - Contains the allergens milk and sulfites

For an explanation of the symbols used here see page 11.

Basagran Brands

**Herbicide Group
6 - bentazon**
(Refer to page 38)

Company:

BASF Canada

Basagran (PCP# 12221)

Basagran Forté (PCP# 22006)

Formulation:

480 g/L bentazon formulated as a solution in both products.
Basagran Forté has a built-in adjuvant.

Container size -

Basagran: 2 x 9 L jugs.

Basagran Forté: 2 x 10 L jugs.

Crops and Staging:

Basagran and *Basagran Forté*:

CROP	STAGE
Soybean	No restrictions
Dry bean ***	After the first trifoliate leaf
Corn	No restrictions
Pea	After 3 leaf pairs but prior to flowering
Fababean	At least 4 inches (10 cm) tall
Flax	After 2 inches (5 cm) in height

Basagran Forté only:

CROP	STAGE
Forage millet and forage sorghum (forage and seed production)*	3 to 6 leaf prior to canopy closure
Established clover (alsike, red) for seed production only*	7.5 to 25 cm prior to canopy closure

Basagran only:

CROP	STAGE
Spring wheat (excluding durum)**	No restrictions (limited to the 4 leaf to flag leaf by 2,4-D staging)
Solin	After 2 inches (5 cm) in height
Forage grasses for seed production*: Bromegrass, creeping red fescue, crested wheatgrass, meadow foxtail, orchardgrass, timothy.	1 to 7 leaf stage
Forage legumes (seedlings) for seed production*: Alfalfa, alsike clover, red clover, sainfoin.	After the third trifoliate leaf
Established alfalfa for seed production.	Prior to flowering
Established clover (Sweet, or Red) and sainfoin for seed production.	3 to 10 inches (7.5 to 25 cm) high

* One application per season.

** *Basagran* only at 0.4 L per acre. Must be tank mixed with 2,4-D (no adjuvant required).

*** Refer to product labels for a list of dry bean types registered for *Basagran*. *Basagran Forté* registered for all dry bean types but not tested for tolerance on all types. Test a small area of a new variety for tolerance before widespread use.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Add Assist or XA oil concentrate at 0.41 to 0.81 L per acre to *Basagran* only. *Basagran Forté* does not require the addition of Assist or XA oil concentrate. If hot, humid conditions prevail (above 28°C and 80% relative humidity), use only the low rate of Assist or XA oil concentrate. *Citowett Plus* may be used on peas at 0.25 L per 100 L spray mixture. Apply the rate listed when weeds in the table are within the recommended height:

ANNUAL WEEDS	0.71 L per acre		0.91 L per acre	
	Inches	Maximum Leaf Stage	Inches	Maximum Leaf Stage
Buttercup			2 to 4	6*
Cleavers			1 to 3 whorl stage	
Cocklebur	3 to 7	6*	7 to 12	10*
Common chickweed			1 to 3 weeks after emergence	
Common groundsel			2 to 4	
Common ragweed			1 to 2	6
Corn spurry			1 to 4	
Flower of an hour	1 to 2	6*	2 to 4	10*
Giant ragweed			2 to 6	4
Hairy galinsoga			2 to 3	6*
Hairy nightshade			0.2 to 0.8	6
Lady's-thumb (smartweed)	1 to 3	6*	3 to 8	10
Lamb's-quarters			0.5 to 1.0	8
Purslane			1 to 2	6
Redroot pigweed (suppression only)			0.5 to 1.5	4
Russian thistle (suppression only*)			1 to 3	4*
Shepherd's-purse	Rosette to 4	6*	4 to 10	6
Stinkweed	Rosette to 2	6*	2 to 6	6
Stork's-bill			1.5 to 4	2 to 6 leaf stage
Volunteer canola	0.75 to 6	8**	0.75 to 6	8
Wild mustard	1 to 5	6*	5 to 10	10
Wild radish			1 to 2	6
PERENNIAL WEEDS	Repeat application 7 to 15 days after first application (if necessary)			
Canada thistle	6 to 8			
Field bindweed	1 to 2.5			
Yellow nutsedge	6 to 8			

*Applies to *Basagran Forté* only.

**Applies to *Basagran* only.

Basagran may be applied in wheat at 0.4 L per acre when tank mixed with 2,4-D amine or ester (500 g per L) at 0.3 to 0.4 L per acre to control the weeds controlled by 2,4-D plus lady's-thumb, redroot pigweed and daisy fleabane. No adjuvant is required for this mix.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 to 160 L per acre. A minimum of 80 L per acre is recommended for optimum control.*

Nozzles and Pressure: Maintain 40 to 60 psi (275 to 425 kPa)* when using conventional flat fan nozzles capable of delivering high water volumes with *ASABE* medium droplets. Low drift nozzles may require higher pressures for proper performance. Contact the herbicide manufacturer regarding the suitability of low drift nozzles for use with this product. Direct nozzles 45° forward to improve contact with vertical targets.

* Higher water volumes and pressures should be used when the weeds are at the upper end of their recommended treatment stage.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Poor results will occur if temperatures are cool. Optimum results are achieved when applied at daytime temperatures between 20 and 28°C. Applications at temperatures greater than 28°C may result in crop injury.

Tank Mixes:

Herbicides:

In soybean:

Pinnacle (2.2 to 3.2 g/acre)

In dry bean:

* *Basagran* only (0.71 L/acre) plus *Reflex* (0.23 L/acre) plus *Agral 90*.

Basagran Forte (0.5L/acre) plus *Solo* (11.7 g/acre + *UAN* liquid 28-0-0 (0.8 L/acre). Failure to add the 28-0-0 could result in weed escapes.

In cranberry dry bean and black dry bean:

Basagran Forté only at 0.7 to 0.91 L/acre can be tank mixed with *Pinnacle* at 3.2 g/acre.

In spring wheat (not including durum):

Basagran only at 0.4 L/acre can be tank mixed with 2,4-D amine or ester (500 g/L) at 0.3 to 0.4 L/acre. This tank mix DOES NOT need any adjuvant.

* For use in the Red River Valley of Manitoba only.

Fertilizers: Use of fertilizer mixes is not recommended for use under western Canadian environmental conditions for most crops. Ammonium sulphate may be added to a *Basagran* spray solution at a rate of 1.5% v/v to improve weed control consistency in dry beans (pinto, great northern, pink and small reds). The risk of bean injury increases with this mixture under hot humid conditions. Use with *Assist Oil Concentrate*. Not for use with *Basagran Forté*.

Insecticides: None registered.

Fungicides: None registered.

When mixing *Basagran Liquid* or *Basagran Forté* refer to the tank mix partner label for any additional restrictions and precautions.

Allow 4 days between application of *Basagran* and other herbicides, fertilizers or insecticides.

Note: The above mixes are those listed on the *Basagran/Basagran Forté* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 to 8 hours will reduce control.

Re-Entry: DO NOT enter treated field for 12 hours.

Grazing: Allow 30 days between treatment with *Basagran Forté* and harvest of forage sorghum and millet for hay. Otherwise DO NOT graze treated crops or cut for feed prior to crop maturity.

Preharvest Interval: 50 days for *Basagran* + 2,4-D in wheat, 84 days for *Basagran* + *Reflex* in Dry beans in Manitoba. Other uses are restricted only by appropriate staging.

Re-cropping: No restrictions the year after application.

Aerial Application: May be applied by air for weed control in dry beans or soybeans only. Use 23 to 45 L/acre water volume. *Assist* or *XA Oil Concentrate* at 0.05 to 0.1 L/acre must be added. DO NOT use *Assist* or *XA Oil Concentrate* in excess of 0.1 L/acre as substantial crop injury could occur. DO NOT apply fertilizer mixes in soybean or 2,4-D tank mix in wheat by air. Crop canopy should NOT cover the weeds.

Storage: May be frozen.

Buffer Zones:

Application method	Crop	Buffer Zones (metres ^{††}) Required for the Protection of: Terrestrial habitat
Ground [†]	Sorghum ^{**} , Forage millet ^{**} , Forage grasses and legumes, Pea [*]	1
	Corn, Dry bean, Fababean, Flax, Pea ^{**} , Soybean, Spring wheat [*]	2
Fixed wing airplane [*]	Dry bean	20
	soybean	35
Helicopter [*]	Dry bean	20
	soybean	30

See page 29 for an explanation of the different habitats.

* *Basagran* only

** *Basagran Forté* only

[†] Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

^{††} Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method B' in the general tank cleaning section on page 15-16.

Hazard Rating:

Basagran

 Caution – Poison

Basagran Forté



Warning – Poison



Danger – Corrosive to Eyes



Warning – contains the allergen soy.

For an explanation of the symbols used here see page 11.

Battalion*

* (For use only on field corn grown in the Red River Valley region of Manitoba)

Herbicide Group

2 - rimsulfuron

4 - dicamba

15 - metolachlor

(Refer to page 38)

Company:

E.I duPont Canada

Formulation:

The *Battalion* package contains the following:

Elim EP (PCP#23518): 25% rimsulfuron formulated as a water dispersible granule.

Container sizes - 480 grams (8 x 60 gram soluble bags).

Dual II Magnum (PCP#25729): 915 g/L s-metolachlor formulated as an emulsifiable concentrate.

Container sizes - 6 L.

Barvel II (PCP#23957): 480 g/L dicamba formulated as a solution of a diglycolamine salt.

Container sizes - 6 L.

Crops and Staging:

May be applied to field corn as a pre-emergent or post-emergent application. For use in the Red River Valley region of Manitoba only. DO NOT use on sweet corn.

Pre-emergent – apply as a broadcast ground treatment after planting but before weeds and corn emerge.

Post-emergent – apply up to the 3-leaf stage (2 visible collars or 20 cm in height – leaf extended). Corn hybrid sensitivity to post-emergent applications of *Battalion* have been observed in the field with varieties rated less than 2500 CHU in regions with less than 2500 CHU. Refer to the Corn Hybrid Sensitivity section of the *Battalion* label for complete details.

* **NOTE** - Since applications to Corn in the Red River Valley region of Manitoba has been registered under the User Requested

Minor Use program, the manufacturer assumes no responsibility for herbicide performance or crop safety. **Application to corn is at the risk of the user.**

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Weeds and Staging:

Pre-emergent applications**:

Common ragweed

Eastern black nightshade

Green foxtail

Lady's-thumb

Lamb's-quarters

Quackgrass (suppression)

Redroot pigweed

Yellow foxtail (suppression)

** Rain is required within 10 days of application or a shallow cultivation or use of a rotary hoe is required.

Post-emergent applications: Apply prior to the 2 leaf stage of the following weeds. *Battalion* must be applied with a non-ionic surfactant (see 'Rates' for details)

Annual smartweeds

(green, lady's-thumb)

Barnyard grass

Cleavers

Common ragweed

Corn spurry

Cow cockle

Eastern black nightshade

False ragweed

Foxtail (Green and Yellow)

Giant ragweed

Hare's-ear mustard

Indian mustard

Lamb's-quarters

Quackgrass (suppression)

Redroot pigweed

Russian pigweed

Tartary buckwheat

Tumble mustard

Velvetleaf

Wild buckwheat

Wild mustard

Wormseed mustard

Rates:

Component	Rate per acre	
	Application timing	
	Pre-emergent	Post-emergent
<i>Elim EP</i>	24 g	20 g
<i>Dual II Magnum</i>	300 mL	250 mL
<i>Banvel II</i>	300 mL	250 mL
Acres per case	20	24

* *Battalion* must be applied with a non-ionic surfactant such as *Citovett Plus*, *Agral 90* or *Agsurf II* at a rate of 0.2 L per 100 L of spray solution (0.2 % v/v) when applied as a post emergent application to corn.

Application Information:

Water Volume: 60 to 70 L per acre.

Nozzles and Pressure: Flat fan nozzles are recommended. Sprayers without drift reduction nozzles should use a maximum of 40 psi (275 kPa). Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE* medium droplets. Use a 50 mesh or coarser screen and filter system

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Rapid fluctuations in temperature (greater than 20°C difference within 24 to 36 hours) will stress the corn crop. For maximum crop safety, allow 48 to 72 hours for the corn to acclimatize before applying *Battalion*.

Apply **ONLY** when the temperature in the 24 hours before AND after application is between 10°C and 25°C. Temperatures beyond this range increase the potential for crop injury. Separate applications of *Battalion* herbicide followed by a broadleaf herbicide (minimum of 12 hours later) will reduce the potential for injury.

WARNING: Crop injury may result if application is made to corn that has been stressed by abnormally hot, humid, or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease, or insect damage. If corn has been injured by frost, wait 48 to 72 hours before applying *Battalion*.

Tank Mixes:

None registered
NOTE: *Battalion* should **NOT** be applied to corn that has been treated with *Thimet*, *Lorsban* or *DiSyston* insecticides. Leave 7 days between the application of *Battalion* and that of a foliar organophosphate insecticide.

Note: The above mixes are those listed on the *Battalion* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. See the general guidelines for mixing pesticides on page 14 for more information.

Restrictions:

Rainfall:

Pre-emergent applications: Rainfall is required within 10 days of application for proper activation of *Battalion*. Otherwise a shallow cultivation or use of a rotary hoe is required.

Post-emergent applications: Within 4 hours of application may result in reduced weed control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated crops or cut for feed.

Preharvest Interval: Leave at least 80 days from application to harvest.

Re-cropping: Corn, winter wheat and barley may be seeded the year following *Battalion* application. For all other crops, a field bioassay is recommended before planting.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†)		
	Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.
DO NOT mix or load within 30 m of any water sources.

Tank Cleaning:

Refer to product label for complete tank cleaning instructions. DO NOT clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

For additional information, Refer to page 15.

Hazard Rating:

Caution – Poison (Banvel II)



Warning – Eye Irritant

Keep out of reach of children

For an explanation of the symbols used here see page 11.

BlackHawk

This product is a prepackaged tank mix of *Aim* (page 77) and 2,4-D (page 66). Information listed is restricted to Crop, Weeds, Rates and Tank mixes. For other detailed restrictions and other general information on the component products see the product pages listed above.

Herbicide Group**4 - 2,4-D****14 - carfentrazone***(Refer to page 38)***Company:**

Nufarm Agriculture

Formulation:

The *BlackHawk* package contains the following components:

Aim EC (PCP# 28573): carfentrazone-ethyl 240 g/L as emulsifiable concentrate.

Container size - 2x 600mL

2,4-D 700 ester (PCP# 27820): 2,4-D 660g/L as emulsifiable concentrate

Container size - 2x 8.69 L.

Crops and Staging:

Pre-seed burndown prior to seeding the following crops:

Spring wheat
Durum wheat
Winter wheat

Barley
Rye
Chemfallow

Weeds and Staging:

Aim EC: 15 mL/acre + *2,4-D 700 ester*: 212 mL/acre.

One case treats 80 acres

Weeds controlled up to 10 cm or 3 leaf rosette or less, unless specified, includes

Kochia*

Mustard

Narrow-leaved

hawk's-beard

Pigweed

* All biotypes.

Prickly lettuce

Russian thistle

Shepherd's-purse

Stinkweed

Volunteer canola*

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

Tank Mixes:**Herbicides:**

Glyphosate (0.5 to 1 L, 360 g ae per acre)

See component products for more information on restrictions application details and handling. Use the most limiting restrictions across all components for the mix.

Blazer

Herbicide Group
14 - acifluorfen
(Refer to page 38)

Company:

United Phosphorus Inc. (PCP# 23315)

Formulation:

240 g/L acifluorfen present as a sodium salt and formulated as a solution.

Container size - 10 L jug.

Crops and Staging:

Soybean from the 1 to 3 trifoliolate leaf stage.

DO NOT apply before the first trifoliolate leaf stage of the soybean.

DO NOT apply to soybeans grown on sand or loamy sand soils.

Weeds and Staging:

Blazer applied at 0.5 L per acre (one jug treats 20 acres) plus *Assist* adjuvant at 0.5 L per 100 L of spray solution will control:

WEED	MAXIMUM LEAF STAGE
Common ragweed	8
Redroot pigweed	4

Blazer applied at 1.0 L per acre** (one jug treats 10 acres) will control the weeds above plus the following weeds at the maximum leaf stages listed:

WEED	MAXIMUM STAGE
Canada thistle*	Pre-bud
Cocklebur	4 leaf
Common milkweed*	-
Field bindweed*	-
Hedge bindweed*	-
Lamb's-quarters	2 leaf
Nightshade (eastern black)	6 leaf
Redroot pigweed	6 leaf
Smartweed (including Lady's-thumb)	8 leaf
Wild Mustard	10 leaf

* Topgrowth control only. The plant will grow back from underground roots

****DO NOT** add *Assist* adjuvant with the 1.0 L per acre rate as crop injury will result.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: No specific water volume is provided on the label but a minimum of 81 L per acre is implied by the adjuvant rates on the label. Good coverage of weed foliage is required for proper control.

Nozzles and Pressure: Use nozzles and pressure designed to deliver thorough, even coverage with *ASABE* medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Soybeans may exhibit speckling, bronzing and/or leaf burn. The trifoliolate leaf emerging at the time of application may be distorted. Soybeans usually outgrow these conditions and continue to grow at a normal rate with no adverse effect on vigour, maturity, or crop yield. It is important to have good spray coverage on the weeds as *Blazer* works mainly by contact action. Failure to follow the suggested application rate and timing may result in unsatisfactory control.

Tank Mixes:

Blazer (0.5 L/acre) plus *Basagran Forté* (0.5 L/acre) or *Blazer* (0.255 L/acre) plus *Basagran** or *Basagran Forté* (0.71 L/acre) depending on predominant weed species present.

See label for details.

*Add *Assist* adjuvant at 0.5 L per 100 L of spray solution for *Basagran* tank mix only.

Fertilizers: None registered. **DO NOT** add fertilizers to the spray mixture.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *Blazer* label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours may reduce weed control.

Re-Entry: DO NOT enter treated fields for 12 hours.

Preharvest Interval: No specific preharvest interval is indicated on the label.

Grazing: DO NOT graze the treated crop or cut for hay.

Re-cropping: The label has no restriction on crops that may be planted the following season.

Aerial application: DO NOT apply by air.

Storage: DO NOT freeze.

Buffer Zones: Leave a buffer of 15 m from the last spray pass and sensitive upland areas such as other crops, pastures, rangeland, woodlots or shelterbelts.

Tank Cleaning:

See page 15 for general information on tank cleaning using Method B. Tanks may require cleaning after several tanks to remove any excessive oil buildup on the inside of the tank.

Hazard Rating:



Warning – Poison



Danger – Corrosive to eyes.



Warning - Causes skin irritation. Avoid contact with skin. Harmful if inhaled.

For an explanation of the symbols used here see page 11.

Broadband

Herbicide Group

1 - pinoxaden

2 - florasulam

(Refer to page 38)

Company:

Syngenta Canada

Formulation:

Broadband (PCP#29138): 92.7 g/L pinoxaden and 7.7 g/L florasulam formulated as an emulsifiable concentrate.

Container size - 10.5 and 84.2 L.

Adigor Adjuvant (PCP#28151): 11.3 and 90.4 L.

Crops and Staging:

Barley, spring wheat (NOT including durum) up to the emergence of the flag leaf.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Grass weeds controlled from 1 to 6 leaves and prior to the emergence of the 4th tiller:

Barnyard grass	Volunteer oat
Foxtail (green and yellow)	Volunteer canary seed
Proso (Crown) millet	Wild oat

Broadleaf weeds controlled at the 1 to 6 leaf stage:

Annual smartweed	Sow-thistle (annual, perennial**) [†]
Common chickweed	Stinkweed
Cleavers	Volunteer canola*
Hemp-nettle [†]	Wild buckwheat
Pigweed, redroot [†]	Wild mustard
Shepherd's-purse	

[†] Suppression only.

* Not Clearfield varieties.

** Applications made at advanced leaf stages will reduce product effectiveness.

Rates:

Broadband: 263 ml per acre.

Add **Adigor** adjuvant at 280 mL per acre.

(Package sizes listed above will treat 40 or 320 acres)

DO NOT apply more than once per season.

Application Information:

Water Volume: 20 to 40 L per acre.

Nozzles and Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets. Low drift nozzles may require higher pressures for proper performance.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops or weeds that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury or reduced weed control may result.

Temporary crop injury may occur with tank-mixes under extreme weather conditions or when the crop is suffering from stress due to inadequate or abnormally high moisture levels or extreme temperatures.

Tank Mixes:

Herbicides:

Curtail M (0.6 L/acre)

MCPA LV500 ester (0.28 L/acre)

Fungicides: Tilt (label rates)

Fertilizers: None registered

Note: The above mixes are those listed on the *Broadband* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Syngenta also supports the following mixes that are not on the *Broadband* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Infinity; Prestige XC (low rate); Trophy.

Fungicides: Propel; Quilt; Tilt (low rate).

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Re-Entry: DO NOT re-enter treated fields within 12 hours.

Preharvest: Leave 60 days between treatment and harvest.

Grazing: DO NOT cut for livestock feed within 30 days or grazed by livestock within 7 days of treating the crop.

Recropping: No restrictions the year following treatment.

Aerial Application: DO NOT apply by air.

Storage: Store in dry, heated storage.

Buffer Zones:

Application method	Buffer Zones (metres †) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground only*	5	30

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15-16.

Hazard Rating:



Warning Eye Irritant

For an explanation of the symbols used here see page 11.

Bromoxynil

Herbicide Group

6 - bromoxynil

(Refer to page 38)

Company:

Bayer CropScience (*Pardner*)

IPCO (*Brotex*)

Nufarm Agriculture (*Koril 235*)

MANA Canada (*Bromotril*)

Formulation:

Pardner (PCP#18001): 280 g/L bromoxynil formulated as an emulsifiable concentrate.

Container size - 2 x 8 L containers per case.

Koril 235 (PCP#25341): 235 g/L bromoxynil formulated as an emulsifiable concentrate.

Container size - 2 x 9.7 L containers per case.

Bromotril II (PCP#30371) & Brotex (PCP#28519): 240 g/L bromoxynil formulated as an emulsifiable concentrate.

Container size - 2 x 9.7 L containers per case.

Crops, Staging and Rates:

Pardner: At 0.40 to 0.48 L per acre one 8 L jug treats 20 to 16.5 acres.

Koril, Brotex & Bromotril: At 0.49 to 0.57 L per acre one 9.71 L jug treats 20 to 17 acres.

See the chart below for registered crops and specific rates and stages. NR = Not Registered.

CROP	STAGE	RATE (L / acre)			
		<i>Pardner</i>	<i>Koril</i>	<i>Bromotril</i>	<i>Brotex</i>
Barley, oats, triticale, wheat (spring and durum)	2 leaf stage to early flag	0.40 to 0.48	0.49 to 0.57	0.49 to 0.57	0.49 to 0.57
Winter wheat	2 to 4 leaf stage (fall application) First growth to early flag leaf (spring application)	0.40 to 0.48	0.49 to 0.57	0.49 to 0.57	0.49 to 0.57
Corn (field or sweet)	4 to 8 leaf	0.40 to 0.48	0.49 to 0.57	0.49 to 0.57	0.49 to 0.57
Corn (field or sweet) with drop pipes	Beyond 8 leaf	0.40 to 0.48	0.49 to 0.57	0.49 to 0.57	0.49 to 0.57
Canaryseed (seed production only)	3 to 5 leaf	0.40	0.49	0.49	0.49
Seedling alfalfa	2 to 6 trifoliate leaf stage	0.40	0.49	0.49	0.49
Established alfalfa (seed production only)	In spring, before the crop begins to shield weeds. Apply no more than twice in one growing season.	0.40 to 0.48	0.49 to 0.57	0.49 to 0.57	0.49 to 0.57
Fall rye	In spring only, from first growth to early flag	0.40 to 0.48	0.49 to 0.57	0.49 to 0.57	0.49 to 0.57
Flax (including Solin)	2 to 4 inches (5 to 10 cm)	0.40	0.49	0.49	0.49
Forage millet and sorghum	4 leaf to 8 inches (20 cm)	0.40	NR	0.49	0.49
Seedling grasses (seed production only): Bromegrass, Fescue (creeping red, meadow), Orchard grass, Reed canary grass, Russian wildrye, Timothy, Wheatgrass (crested, intermediate, slender, tall)	2 to 4 leaf (Establishment year only)	0.40 to 0.48	NR	0.49 to 0.57	0.49 to 0.57
Pearl millet and sorghum (grain)*	4 leaf to 8 inches (20 cm)	0.40	NR	NR	NR
Prior to direct-seeding cereal crops (mixed with glyphosate only)	Apply according to weed stage.	0.40	NR	0.49	0.49

*NOTE: Since application to grain pearl millet and sorghum is registered under User Requested Minor Use Label Expansion program, the manufacturer assumes no responsibility for herbicide performance. Users of this on product grain pearl millet and sorghum do so at their own risk.

Weeds and Staging:

Weeds controlled at the 1 to 4 leaf stage:

American nightshade	Cow cockle*
Annual smartweed (green, pale, lady's-thumb)	Kochia**
Bluebur	Pigweed*†
Cocklebur	Russian thistle**
Common ragweed	Stinkweed*
	Wild mustard*

Weeds controlled at the 1 to 8 leaf stage:

Common groundsel	Tartary buckwheat
Lamb's-quarters	Wild buckwheat
Tame buckwheat	

* Controlled with high rate only.

** Apply before plants are 2 inches high.

† Not controlled in seedling alfalfa.

Application Information:

Water Volume:

Ground:

Corn, Millet & Sorghum: 80 to 120 L per acre.

Seedling grasses: 60 L per acre.

Other crops: 40 L per acre.

Aerial (wheat and barley only): 8 to 16 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Avoid spraying if temperatures are greater than 25°C. Leaf scorching may occur in corn and flax if applied during or after adverse growing conditions, such as cool and wet or hot (greater than 27°C) and humid weather.

Tank Mixes:

Herbicides:

CROP	TANK MIXES
Spring wheat	2,4-D†, Achieve Liquid, Avenge, Avenge + MCPA (ester only)
Winter wheat	2,4-D, Achieve Liquid, MCPA
Barley	2,4-D†, Achieve Liquid, Avenge, Avenge + MCPA (ester only), MCPA†
Oats	MCPA
Fall rye, canaryseed	MCPA*
Flax	MCPA (amine, ester or K salt)
Seedling forage grasses***	MCPA
Corn	Accent◇ + surfactant (field corn only), Atrazine**, Banvel II (field corn only)**

* The ester formulations are preferred but other formulations can be used.

** Do not add oil or surfactant to this mix. Do not use atrazine formulations that contain oil.

*** Bromotril, Brotex and Pardner only.

† May be applied by air.

◇ Since the use of this tank mix on corn is registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Users of this tank mix on corn do so at their own risk.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the bromoxynil labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Bromoxynil manufacturers may also support mixes with pesticides that are not on the bromoxynil labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application will reduce control.

Re-entry: DO NOT enter treated fields for 24 hours.

Grazing: DO NOT graze treated wheat, barley, oats, forage millet, sorghum or seedling alfalfa crops or cut for feed within 30 days of application.

DO NOT graze other treated crops or cut for hay prior to crop maturity.

Re-cropping: No restrictions.

Aerial Application: Registered for aerial application on wheat and barley. The use of low water volumes, 8 to 16 L per acre may result in less effective weed control than seen with ground application.

Storage: May be stored at freezing temperatures.

Will return to original state by warming to room temperature (20 to 22°C) and agitating thoroughly.

Buffer Zones:

Application method	Buffer Zones (metres †) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	1	1
Fixed wing aircraft**	20	5	55
Helicopter**	20	3	45

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy

**Wheat and barley crops only.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

All:



Warning – Poison

Brotex 240, Bromotril 240:



Warning – Eye and Skin Irritant

Koril 235:



Danger – Eye and Skin Irritant

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Bromoxynil/2,4-D Ester

Herbicide Group
4 - 2,4-D
6 - bromoxynil
(Refer to page 38)

Company:

Bayer CropScience (*Thumper*)
IPCO (*Leader*)
MANA Canada (*Thrasher II*)
Nufarm Agriculture (*Approve*)

Formulation:

Thumper (PCP# 22659): 280 g/L bromoxynil and 280 g/L 2,4-D ester formulated as an emulsifiable concentrate.
Container size - 8 L.

Approve (PCP# 28123), *Leader* (PCP# 28853) & *Thrasher II* (PCP# 30372): 225 g/L bromoxynil and 225 g/L 2,4-D ester formulated as an emulsifiable concentrate.

Container size - 10 L, 100 L*, 500 L*

* *Approve* only

Crops and Staging:

Spring wheat (including durum) and barley at the 4 leaf to early flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 1 to 4 leaf stage:

American nightshade	Lady's-thumb
Ball mustard	Night-flowering catchfly
Bluebur	Pale smartweed
Cocklebur	Redroot pigweed
Common ragweed	Shepherd's-purse
Cow cockle	Volunteer canola
Flixweed	Volunteer sunflower
Green smartweed	

Weeds controlled at the 1 to 8 leaf stage:

Common groundsel	Tartary buckwheat
Lamb's-quarters	Wild buckwheat
Stinkweed	Wild mustard
Tame buckwheat (up to 4 leaf stage with <i>Approve</i>)	

Weeds Controlled from 1 to 12 leaf (max. 2 inches tall):

Kochia	Russian thistle
--------	-----------------

Rates:

Thumper: 0.4 L per acre (one 8 L container treats 20 acres).

Approve, *Leader* or *Thrasher*: 0.5 L per acre
(One 10 L container treats 20 acres).

Application Information:

Water Volume:

Ground: 20 to 40 L per acre.

Aerial: 12 to 16 L per acre. Use the higher volume when there is a heavy crop canopy, or when the majority of weeds are cow cockle, smartweed, or pigweed.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional 80° or 110° flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets. All strainer and nozzle screens must be 50 mesh or coarser.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Less than acceptable weed control may be expected if weeds are under stress because of excessive moisture, drought, or cool weather.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Liquid Achieve (0.2 L/acre) plus Turbocharge adjuvant
Avenge (1.4 to 1.7 L per acre)

In spring wheat (including durum) only:

Clodinafop 240L/C (93 to 117 mL/acre)^Δ plus adjuvant

* *Approve*, *Leader* and *Thrasher* only.

^Δ Manufacturers may only support specific mixes. Contact the manufacturer for more information.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Bromoxynil/2,4-D Ester* labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Bromoxynil/2,4-D ester manufacturers may also support mixes with pesticides that are not on the *Bromoxynil/2,4-D ester* labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT enter treated fields for at least 24 hours.

Grazing: DO NOT graze or cut for livestock feed within 30 days of application. Withdraw meat animals 3 days before slaughter.

Preharvest Interval: DO NOT harvest within 30 days of application.

Re-cropping: No restrictions the year after application.

Aerial Application: May be applied by air.

Storage: May be frozen. Shake well before using after being frozen.

Buffer Zones:

Application method	Buffer Zones (metres †) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	1	1
Fixed wing aircraft	20	5	55
Helicopter	20	3	45

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.


† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:


Refer to page 15.

Hazard Rating:


All Products:


 Warning – Poison

Thumper and Approve

 Caution – Skin and Eye Irritant

Leader and Thrasher

 Warning – Skin and Eye Irritant

 Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Bromoxynil/MCPA ester

Herbicide Group

4 - MCPA

6 - bromoxynil

(Refer to page 38)

Company:

Bayer CropScience (*Buctril M*)

IPCO (*Logic M*)

Nufarm Agriculture (*Mextrol 450*)

MANA Canada (*Badge*)

Formulation:

Buctril M (PCP#18022): 280 g/L bromoxynil and 280 g/L of MCPA ester formulated as an emulsifiable concentrate. Container size - 8 L.

Mextrol 450 (PCP#26999), *Badge II* (PCP#30370) & *Logic M* (PCP#28109): 225 g/L bromoxynil and 225 g/L of MCPA ester formulated as an emulsifiable concentrate. Container size - 10 L.

Crops and Staging:

Field Crops:

All Products:

CROP	STAGE
Barley, oats, spring wheat (including durum)	2 leaf to early flag
Winter wheat	2 to 4 leaf stage in the fall or after growth resumes up to early flag leaf.
Fall rye	When growth commences in spring to early flag leaf
Canaryseed	3 to 5 leaf stage
Flax, Solin	2 inches (5 cm) to early bud stage. Best tolerance occurs when flax is 2 to 4 inches (5 to 10 cm) tall.
Corn	4 to 6 leaf stage

Seedling forage grasses†:

2 to 4 leaf stage†.

All Products:

Bromegrass	Russian wild-rye
Fescue (creeping red, meadow)	Timothy
Reed canarygrass	Wheatgrass (crested, intermediate, slender, tall)

Buctril M, Logic M and Badge only:

Fescue (tall)	Orchard grass
Meadow bromegrass	Wheatgrass (streambank)
Meadow foxtail	

Established Forage Grasses:

Timothy (seed† or hay††)** - prior to emergence of the flag leaf.

† Maximum of two treatments per year at least 21 days apart.

†† Maximum of two treatments per year at least 90 days apart.

Perennial Cereal Rye* (Buctril M only):

Established stands: 2 leaf up to early flag leaf stage.

Establishment year: 2 to 4 leaf stage in the fall, or from the time growth commences to early flag leaf stage in the spring.

* Since the use of this tank mix on perennial cereal rye is registered under the User Requested Minor Use Label Expansion program, the manufacturer assumes no responsibility for herbicide performance. Users of this tank mix on perennial cereal rye do so at their own risk.

** Applications onto Timothy for hay production registered with Buctril M, Logic M, and Mextrol 450 only.

Weeds and Staging:

Weeds up to 4 leaf stage:

American nightshade	Kochia**
Annual smartweeds (green, pale, lady's-thumb)	Night-flowering catchfly
Bluebur	Redroot pigweed*
Ball mustard	Russian thistle**
Cocklebur	Scentless chamomile***
Cow cockle	Shepherd's-purse
Flixweed	Volunteer canola
	Volunteer sunflower

* May not be controlled in flax.

** Control before plants are 2 inches tall.

*** Spring seedlings only.

Weeds up to 6 leaf stage:

Wild tomato (Buctril M, Logic M & Badge only)

Weeds up to 8 leaf stage:

Common groundsel	Tartary buckwheat
Common ragweed	Wild buckwheat
Lamb's-quarters	Wild mustard
Stinkweed	Wormseed mustard
Tame buckwheat	

Weeds suppressed in winter wheat from the 2 to 12 leaf stage:

Prickly lettuce (Mextrol 450 only)

Weeds where top growth is controlled:

Canada thistle	Perennial sow-thistle
----------------	-----------------------

Rate:

Buctril M: 0.4 L per acre. One 8 L jug treats 20 acres.

Mextrol 450, Badge & Logic M: 0.5 L per acre. One 10 L jug treats 20 acres.

Application Information:

Water Volume:

Corn: 80 to 120 L per acre.

Flax, Solin: 20 to 40 L per acre.

Cereals: 20 to 40 L per acre.

Seedling forage grasses: 60 L per acre.

Established timothy: 60 L per acre.

Perennial Cereal Rye: Not less than 20 L per acre.

Aerial: 8 to 16 L per acre.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets. Refer to specific labels for recommended water volumes.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Best weed control when humidity is high at the time of spraying and for the following day or two. Prolonged cool conditions may result in reduced weed control. Spraying during early morning may increase the risk of flax injury.

Avoid spraying in temperatures greater than 25°C.

DO NOT apply to flax, canaryseed or corn if daytime temperatures exceed 27°C within 48 hours before or after application.

Tank Mixes:

Herbicide Tank Mix Table:

Products listed below are at label rates for each crop. See labels for details.

CROP & TANK MIXES	Badge	Buctril M	Logic M	Mextrol 450
Flax (including Solin):				
Post Ultra + Merge adjuvant	✓	✓	✓	✓
Clethodim + adjuvant	✓*	✓**	✓	✓*
Spring wheat (including durum) and Barley:				
Liquid Achieve	✓	✓	✓	✓
Ally	✓	✓	✓	✓
Avenge	✓	✓	✓	✓
MCPA (amine, ester & K)	✓	✓	✓	✓
Refine SG (4 g / acre) #			✓	
Spring wheat only (NOT including durum):				
Axial 100EC	✓			
Spring wheat only (NOT including durum) and Barley:				
Axial 100EC		✓	✓	
Winter Wheat:				
Refine SG (4 g / acre) #			✓	

Oats:				
MCPA (amine, ester & K)	✓	✓	✓	✓
Corn:				
Atrazine	✓	✓	✓	✓

* Select only

** Select and Centurion only.

Requires the addition of a surfactant as per *Refine SG*.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *bromoxynil/MCPA ester* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Bromoxynil/MCPA manufacturers may also support mixes with pesticides that are not on the *bromoxynil/MCPA* labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall within 1 hour of application may reduce weed control.

Re-entry: DO NOT enter treated fields for at least 24 hours, or 15 days for corn to be harvested by hand.

Preharvest Interval: DO NOT harvest perennial cereal rye within 30 days of application.

Grazing: DO NOT graze treated grain or established timothy crops or cut for feed within 30 days of application. DO NOT graze meadow foxtail in the year of treatment. DO NOT graze other treated forage grasses within 56 days of treatment.

Preharvest Interval: DO NOT harvest Flax or Solin within 60 days of application.

Re-cropping: No re-cropping restrictions the year after treatment.

Aerial Application: May be applied by air to wheat, barley, and oats only. Use higher water volume (see 'Application Information') when the majority of weeds are cow cockle, smartweed, hemp-nettle, pigweed, and Canada thistle.

Storage: May be frozen. Shake the container well when thawed to reconstitute components before use.

Buffer Zones:

Application method	Crop	Buffer Zones (metres †) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Ground only*	All	1	1	4
Fixed wing aircraft	Oats	15	2	60
	Barley & wheat	20	5	60
	Rye	1	0	60
Helicopter	Oats	15	1	50
	Barley & wheat	20	3	50
	Rye	1	0	50

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.


† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:


Refer to page 15.


Hazard Rating:


Badge, Buctril M, Logic M:

 Warning

Mextrol 450

 Danger - Poison

 Warning - Skin Irritant
Potential Skin sensitizer

 Caution - Eye irritant

For an explanation of the symbols used here see page 11.

Casoron**Herbicide Group**

20 - dichlobenil

(Refer to page 38)

Company:

Chemtura Canada (PCP#12533)

Formulation:

4% dichlobenil formulated as a granular.

Container size - 15 kg, 22.7 kg.

Crops:

Poplar plantations

Shelterbelts consisting of the following species:

Ash	Juniper
Barberry	Lilac
Birch (cutleaf-weeping)	Linden

Boxwood

Caragana

Cedar (White, Eastern Red)

Crabapple

Elm

Euonymus (Burning bush)

Forsythia

Honeysuckle

Locust

Maple

Mock orange

Poplar

Rose

Spirea

Willow

NOTE: DO NOT apply to shelterbelts with mugo pine, firs, hemlock, holly, spruce or other shallow rooted species or injury may result. **DO NOT** apply in or around greenhouses. **DO NOT** use on light sandy soils.

Weeds and Staging:

Apply in early spring or late fall prior to annual weed emergence, or after cultivation has removed existing weeds.

Annual blugrass	Mustard
Artemisia (absinthe,* worm-wood, sage)	Nutsedge*
Bindweed*	Pigweed
Canada thistle*	Plantain
Chickweed	Purslane
Dandelion*	Quack grass*
Foxtail (green and yellow)	Sheep sorel*
Groundsel	Shepherd's-purse
Horsetail	Smartweed
Knotweed	Sow-thistle
Kochia	Spurge
Lamb's-quarters	Vetch*
Loosestrife	Wild buckwheat*

* Controlled with fall applications at the higher rates.

Rates:

45 to 70 kg per acre.

At the low rate, a 15 kg bag will treat a 4 yd by 407 yd (4 m by 340 m) strip of shelterbelt. At the high rate, a 15 kg bag will treat a 4 yd by 256 yd (4 m by 214 m) strip of shelterbelt. If application is followed by 0.5 to 1.0 inches (1.3 to 2.5 cm) of irrigation, the lower rates are recommended.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply during periods of high soil temperatures (more than 15°C).

Tank Mixes:

None registered.

Restrictions:

Rainfall: Does not reduce activity.

Re-entry: DO NOT enter treated areas for at least 24 hours.

Grazing: DO NOT graze in treated area.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. DO NOT freeze.

Buffer Zones: Site characteristics and conditions that may lead to runoff include, but are not limited to, heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g., soils that are compacted, fine textured or low in organic matter). Avoid application of this product when heavy rain is forecast.

Equipment Cleaning:

Refer to page 15.

Hazard Rating:

No specific rating. Keep out of reach of children.

Harmful if swallowed.

Avoid skin or eye contact.

CleanStart

Company:

Nufarm Agriculture

Herbicide Group

9 - glyphosate

14 - carfentrazone

(Refer to page 38)

Formulation:

One case of *CleanStart* contains 2 components:

Credit (PCP# 25866): 356 g/L glyphosate formulated as a solution.

Container size - 2 x 10 or 450 L.

Aim (PCP# 28573): 240 g/L carfentrazone formulated as an emulsifiable concentrate.

Container size - 1 x 600 mL or 4 x 3.38 L.

Crops and Staging:

Prior to the seeding of most crops* including the following:

Barley	Oats
Beans, dry	Peas, field
Buckwheat	Potato*
Canola	Rye
Chickpea	Safflower
Corn	Sorghum
Flax	Soybean
Lentil	Sunflower
Millet (pearl and proso)	Triticale
Mustard	Wheat

*Note – before using any pesticide on potatoes, consult the list of "Agricultural Pesticides Approved for Use", available from Simplot Canada and McCain Foods (Canada).

Harvest Aid Treatment:

CleanStart may be applied to the following crops when seed is at less than 30% moisture to speed the rate of dry-down of the following crops and green weedy material. See glyphosate preharvest uses for additional staging information.

CROP*	RATE		
	Credit (glyphosate) (g ae/acre) [†]	Aim (mL/acre)	Acres per case
Barley, chickpea, dry bean, field pea, oats, wheat,	360 ^{†**}	30	20

* DO NOT apply to crops if grown for seed purpose.

[†] Equivalent to 1L per acre of a 360 g/L formulation.

Weeds, Rates and Staging (Pre-seeding):

Credit 0.5 to 1.0 L per acre plus Aim at 15 to 30 mL per acre (40 to 20 acres per case or 900 to 450 acre bulk): Weeds controlled by glyphosate at the above rates plus rapid burnoff of:

Chickweed	Tansy mustard
Dandelion (spring seedlings only)	Volunteer canola (including all herbicide tolerant vari- eties)*
Shepherd's-purse	

Kochia (including glypho-
sate resistant biotypes**)

Apply to actively growing weed up to 10 cm in height.

* 1 to 3 leaf stage for glyphosate tolerant volunteer canola

** Use highest registered rate to control glyphosate resistant kochia.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

Application Information:

Water Volume: Minimum 40 L per acre. Higher water volumes will give better performance from the carfentrazone active. Use higher volumes when weed populations are dense.

Nozzles & Pressure: Sprayers without drift reduction nozzles should use maximum pressure of 30 psi (210 kPa). Low drift nozzles may require higher pressures for proper performance. Apply using nozzle and pressure combination that deliver an even spray pattern with good coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Most effective control is achieved when grasses are actively growing. Weeds stressed by drought, flooding, hot or prolonged cool temperatures (<15°C) and poor fertility are more difficult to control. Symptoms of carfentrazone activity on weeds may be accelerated by warm moist conditions. Weeds hardened off by drought may be more difficult to control.

Tank Mixes:

Herbicides:

Aim (17 mL/acre)

Fungicides: None registered.

Insecticides: None registered.

Fertilizers: None registered.

Do not use with additives that are used to modify spray pH.

Restrictions:

Rainfall: Rainfall within 6 to 8 hours after application may reduce activity. Avoid application when heavy rainfall is forecast.

Re-Entry: DO NOT re-enter treated fields for at least 12 hours.

Preharvest Interval: Leave a minimum of 3 days between harvest aid treatment and harvest.

Grazing: DO NOT graze the treated crop or cut for feed.

Re-cropping: CleanStart may be applied as a preseed burn-off prior to the seeding of most crops. Check the product label for a complete list. There are no rotational restrictions 12 months after application.

Aerial Application: DO NOT apply by air

Storage: Store in a cool dry location.

Buffer Zones: Leave a buffer of 3 meters from the downwind edge of the boom to sensitive upland habitats. Apply near wetlands only when wind is blowing away from wetlands.

Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

Tank Cleaning:

Sprayers used to spray *CleanStart* should be drained and flushed out immediately after use. The manufacturer recommends that sprayers used to apply this product be flushed with a water/ ammonia rinse (3 L of 3% ammonia per 100 L of water) by circulating for a minimum of 15 minutes through the tank and all hoses, booms and nozzles. All nozzles, screens and filters should be removed (including disassembly of multi-nozzle assemblies) and cleaned after applying this product. If possible leave the ammonia solution in the tank overnight to enhance cleaning. Rinse the tank, booms, hoses, and nozzles with clean water to complete the process.

DO NOT clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Hazard Rating:



Caution – Skin and Eye Irritant

For an explanation of the symbols used here see page 11.

Clethodim

Herbicide Group

1 - clethodim

(Refer to page 38)

Company:

Arysta LifeScience (Select - PCP# 22625)

Bayer CropScience (Centurion - PCP# 27598)

MANA Canada (Arrow - PCP# 28224)

Loveland Products (formerly Viterro) (Shadow RTM - PCP# 29277)

Formulation:

240 g/L clethodim formulated as an emulsifiable concentrate. Container size - 3 L clethodim + 9 L adjuvant.

Crops, Rates and Staging:

Crops are tolerant at all growth stages although maximum rates but Preharvest Intervals must be observed to prevent excess residue in the grain (see 'Restrictions:').

To a maximum rate of 75 mL per acre:

Chickpea*

Dry bean[†] (black, great northern, navy, pink, pinto, red)

Prairie Carnation^{**†}

To a maximum rate of 150 mL per acre:

Alfalfa (seedling only)

Canola

Coriander^{**†}

Fenugreek^{***†}

Field pea

Flax (including Solin)

Lentil

Mustard, condiment (brown, oriental, yellow)

Mustard, oilseed types
(*B. juncea*)^Δ

Potato

Safflower^{Δ†}

Soybean

Sunflower

* Apply up to the 9 node stage (7 inches or 18 cm maximum height)

** Apply in the 2 to 5 leaf stage, one application per year.

*** Apply in the 3 to 5 leaf stage, one application per year.

Registered for all *Phaseolus vulgaris* varieties. Since not all varieties of dry beans have been tested for tolerance to clethodim, first use of clethodim should be limited to a small area of each variety to confirm tolerance. Arrow and Shadow RTM are registered for use on black, great northern, navy, pink, pinto, and red dry bean types

Δ Select, Centurion and Shadow RTM only.

ΔΔ Select and Centurion only.

† NOTE: Since these uses are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Those who apply these uses do so at their own risk.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

DO NOT apply more than a total rate of 150 mL per acre of these products, or other products containing clethodim, to the same field per season.

Adjuvants: *Clethodim* products must be applied with 0.5 L of *Amigo* adjuvant (*Centurion*, *Shadow RTM* or *Select*) or *X-ACT* adjuvant (*Arrow*) per 100 L of spray solution (unless otherwise indicated on the label). For spray water sources high in bicarbonate ions (CO_3) see 'Effects of Growing Conditions' section below.

Refer to the product labels for complete mixing instructions. A general guide to mixing can be found on page 13

WEED	RATE (mL/ ACRE)	ACRES TREATED PER 3 L CONTAINER	STAGE
Barnyard grass, foxtail (green, yellow)*†, proso millet, volunteer cereals (barley*†, canary seed, corn, oat*†, wheat*†), wild oat†	50**	60	Apply at 2 to 6 leaf stage. † Apply at the 2 to 4 leaf stage when treated with the 60 acre rate.
Moderate to heavy infestations of the above grasses, plus Persian dandel	75	40	For best results in either case, apply at the 2 to 3 leaf stage
Quackgrass (suppression only)	75	40	2 to 6 leaf stage when 3 to 6 inches (6 to 15 cm) tall. For best results, apply at the 3 to 5 leaf stage
Quackgrass (season long control)	150***	20	

* Apply to light infestations of these weeds only for the 60 acre per jug rate. The manufacturers do not provide guidelines for weed densities under light infestations. When in doubt as to the level of weed infestation, use the higher rate or contact the manufacturer.

** At this rate, *clethodim* should NOT be tank mixed with any other pesticide and should only be applied under the following growing conditions: good crop stand, within the recommended leaf staging (2 to 3 leaf is optimum timing) prior to tillering, light weed infestations, adequate moisture and fertility, absence of stress, and good growing conditions.

*** Apply with 1 L of adjuvant per 100 L of spray solution (*Amigo* adjuvant with *Centurion*, *Shadow RTM* and *Select*, *X-ACT* adjuvant with *Arrow*).

Refer to the product labels for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre. Use 40 L per acre under dense weed infestations or dense crop canopies.

Aerial: Minimum of 11.3 L/acre.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional 80° flat fan nozzles tilted forward at a 45° angle. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver proper coverage with *ASABE medium* droplets or larger.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Clethodim will be less effective when plants are stressed by lack of moisture, excessive moisture, low temperature

and/or very low relative humidity. Re-growth of tillers may occur if application is made under any of the above stress conditions.

Clethodim activity is reduced by levels of bicarbonate ions in spray water equal to or greater than 500 ppm. The addition of ammonium sulphate at 1.6 L/acre (480 g/L liquid) or 0.8 kg/acre (99% dry), or the addition of 28-0-0 liquid fertilizer at 0.5 L per acre to the tank prior to the addition of *clethodim* has been shown to restore control.

Tank Mixes:

Clethodim may be tank mixed with other pesticides at the 75 to 150 mL per acre rates. Add the recommended amount of adjuvant with all tank mixes unless otherwise indicated.

Herbicides:

In flax (not including solin)
Bromoxynil/MCPA ester (label rates)^Δ
Curtail M (0.6 to 0.8 L/acre)[†]
MCPA ester (rates for flax).
Lontrel at 0.23 to 0.34 L/acre.[†]

In solin (low linolenic flax)

Bromoxynil/MCPA ester (label rates)^Δ

Curtail M (0.6 to 0.8 L/acre)[†]

Lontrel at 0.23 to 0.34 L/acre[†]

In canola:

Lontrel at 0.17 to 0.34 L/acre[†]

Muster at 8 to 12 g/acre[†] (redroot pigweed is controlled at the 8 g/acre rate of *Muster* in this tankmix).

In Clearfield canola only:

Pursuit at 42 to 85 mL/acre[†]

In Liberty Link canola only:

Liberty at 1.1 to 1.35 L/acre plus *clethodim* at 25.5 mL/acre (120 acres/case) plus 0.5 L of the recommended adjuvant per 100L of spray solution. When mixing add adjuvant to the water first, then *Liberty*, then *clethodim*. Consult labels for detailed mixing instructions.

In field peas:

Pursuit (85 mL/acre)[†]

In Glyphosate tolerant soybean:

Glyphosate (540 g ai formulation) at 0.68 to 1.35 L/acre.[†] ♦

* Light infestations only in Clearfield canola. For heavy infestations, use high rate of *Pursuit*.

† Apply with the 75 mL/acre rate of *clethodim* only.

Δ Manufacturers may only support specific mixes. Contact the manufacturer for more information.

♦ Arrow only.

Allow 4 days between application of *clethodim* and any other chemical not recommended as a tank mix combination on the label.

Fertilizer: None registered.

Insecticide: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *clethodim* labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Clethodim manufacturers also support mixes with pesticides that are not on the *clethodim* labels.

Herbicides: *Liberty 150* with *clethodim* rates of 50 ml/acre)

Check with each manufacturer for other products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-Entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT graze or cut treated crops for forage until 60 days after application of *clethodim* to annual crops, and 30 days after application to seedling alfalfa.

Preharvest Interval:

Preharvest Interval (days)	CROP(S)
30	Alfalfa, fenugreek
60	Canola, coriander, dry beans, flax (including Solin), lentils, potatoes, chickpeas (Desi and Kabuli) or mustard (brown, yellow, oriental)
72	Sunflower
75	Soybeans, field peas

Aerial Application: Only *Centurion*, *Shadow RTM* and *Select* may be applied by air in canola, chickpea, dry bean, flax, field pea, lentil, mustard, potato, soybean, sunflower only.

Storage: May be stored at any temperature. Shake well before use.

Other: DO NOT apply more than 0.15 L per acre to the same land area per season.

Buffer Zones:

Application Method	Crops	Buffer Zones (metres †) Required for the Protection of:
		Terrestrial habitat
Ground	All Crops	15
Aerial	Desi and Kabuli chickpeas, dry common beans	15
	Other Registered Crops	20

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method B' in the general Tank Cleaning section on page 15-16.

Hazard Rating:

Select, *Shadow* and *Centurion*:



Warning – Skin and Eye Irritant

Arrow:



Caution – Skin and Eye Irritant

For an explanation of the symbols used here see page 11.

Clodinafop

Herbicide Group

1 - clodinafop

(Refer to page 38)

Company:

AgWest

(*Slam'R* - PCP#30137; *Slam'R* COC Adjuvant - PCP#30138)

Arysta LifeSciences

(*NextStep* NG - PCP#29614, built in adjuvant)

Cheminova

(*Bullwhip* - PCP#30445; *XA Oil Concentrate Adjuvant* - PCP#11769)

Farmers of North America

(*Aurora* - PCP#29711; *Chem-Spray* COC Adjuvant - PCP#29712)

IPCO

(*Legend* - PCP#29526; *Legend B Adjuvant* - PCP#29527)

MANA Canada (*Ladder* - PCP#29495;

*XA Oil Concentrate** - PCP#11769; *MANA Adjuvant 80* - PCP#30419)

Nufarm Agriculture

(*Signal* - PCP#29172; *Signal Adjuvant* - PCP#29173)

(*Nufarm Clodinafop* - PCP#29962; *Enhance Adjuvant* - PCP#29952)

Syngenta Canada

(*Horizon* NG - PCP#29089; built in adjuvant)

Loveland Products (formerly Viterra)

(*Foothills* NG - PCP#30341; built in adjuvant)

* This product is no longer manufactured but some product still remains in the retail system. This product may not be in future editions.

Formulation:

Aurora, *BullWhip*, *Ladder*, *Legend*, *Signal*, *Nufarm Clodinafop*, *Slam'R*: 240 g/L clodinafop-propargyl formulated as an emulsifiable concentrate.

Container size - 3.68 L, 11 L, 18.4 L or 2 x 11.04 L.

Slam'R COC, *Signal*, *Steppe* and *XA Oil Concentrate Adjuvants*: 2 x 6.4 L.

Enhance Adjuvant: 20 L.

MANA Adjuvant 80: 4 L; 12 L

*Horizon NG**, *Foothills NG**, *NextStep NG**: 60 g/L clodinafop-propargyl formulated as an emulsifiable concentrate.

Container size - 2 x 7.57 L, 121.1 L or 424.4 L.

* These products have a built in adjuvant system and does not require the addition of an adjuvant.

Crops and Staging:

Spring wheat (including durum) - prior to the emergence of the 4th tiller.

When tank mixing, check broadleaf product description for additional restrictions.

Weeds, Rates and Staging:

NG Formulations: 376 mL per acre, no additional adjuvant required (packages listed above treats 40, 322 or 1129 acres).

-or-

240EC Formulations: 93 mL per acre plus recommended adjuvant at 0.8 L per 100 L spray solution. For *Nufarm Clodinafop* and *Signal* only add *Enhance* adjuvant, or for *Ladder* add *MANA Adjuvant 80*, at 0.25 L per 100L spray solution).

For control of:

WEED	STAGE
Barnyard grass	1 to 5 leaf prior to tillering
Green and yellow foxtail	1 to 5 leaf stage, prior to emergence of 3rd tiller
Volunteer canaryseed, wild oats	1 to 6 leaf, maximum 3 tillers
Volunteer oats	3 to 6 leaf, maximum 3 tillers

NG Formulations: 474 mL per acre, no additional adjuvant required (package sizes above treats 32, 258 or 903 acres);

-or-

240EC Formulations: 115 mL per acre plus recommended adjuvant at 1.0 L per 100 L spray solution of the recommended adjuvant. For *Nufarm Clodinafop* and *Signal* only add *Enhance* adjuvant or for *Ladder* add *MANA Adjuvant 80* at 0.32 L per 100L spray solution).

For control of:

WEED	STAGE
Persian dandel	1 to 5 leaf prior to tillering

Apply at the 2 to 3 leaf stage for optimum control.

Optimum weed control and yield response occurs when weeds are controlled before tillering.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume:

Ground - 20 L to 40 L per acre.

Aerial - 12 L/acre.

Nozzles and Pressure: 40 to 45 psi (275 to 310 kPa) when using conventional 80° or 110° flat fan stainless steel nozzles tilted forward at an angle of 45°. Low drift nozzles may require higher pressures for proper performance. Consult with herbicide manufacturer regarding the suitability of low drift nozzles for use with this product.

Tank Mixes:

Mixes provide control of wild oat, green foxtail, and weeds/insects controlled by the tank mix partner unless otherwise noted.
Herbicides:

	Aurora	BullWhip	NG Formulations	Ladder	Legend	Nufarm Clodinafop	Signal	Slam'R
2,4-D amine (0.34 to 0.45 L/acre) ^{††}	•	•	•	•	•	•	•	•
Bromoxynil [#]	•	•	•	•	•	•	•	•
Bromoxynil/MCPA ^{***} [#]	•	•	•	•	•	•	•	•
Bromoxynil/2,4-D (label rates)	•	•	•	•	•	•	•	•
Curtail M (0.6 to 0.81 L/acre)	•	•	•	•	•	•	•	•
Dichlorprop/2,4-D (0.71 L/acre) ^{**}	• ^Δ	•	• ^Δ	• ^Δ	• ^Δ	• ^Δ	• ^Δ	• ^Δ
DyVel (0.4 to 0.50 L/acre)	•	•	•	•	•	•	•	•
Lontrel (0.17 to 0.34 L/acre)			•	•	•		•	
Lontrel (0.11 to 0.17 L/acre) + MCPA ester (0.45 L/acre) ^{††}	•	•	•	•	•	•	•	•
MCPA amine or ester ^{††} (0.34 to 0.45 L/acre)	•	•	•	•	•	•	•	•
MCPA Sodium Salt (0.48 to 1.09 L/acre) [*]			•	•	•		•	
Mecoprop-p (2.2 to 2.8 L/acre)			•	•	•		•	
Metsulfuron (3 g/acre) ^{†††} [#]	•		•	•	•	•	•	•
Pulsar (80 acres / case)			•					
Pulsar + MCPA Ester (rates above)			•					
Refine SG (12 g/acre) ^{†††} or (N = Nimble (8 g/acre) ^{†††} [#]		N	•		•		•	
Target (0.4 to 0.6 L/acre) ^{**}			•	•	•		•	
Trophy (20 acres per case)			•	•	•		•	

Refer to the broadleaf herbicide label for crop staging, and other information. When tank mixing *clodinafop* 240 EC, always add the broadleaf herbicide first, followed by *clodinafop*, with the adjuvant added last. Reductions in green foxtail and wild oat control may be observed when tank mixed with 2,4-D amine and MCPA amine.

Insecticides: Matador (25 to 33 mL/acre) [◇]

Fungicides: Tilt (0.1 L[#] to 0.2 L/acre) [◇]

Fertilizers: None registered.

Clodinafop may also be mixed with *Matador* plus *Tilt* at the rates above. [◇]

[#] Check product label for specific tank mix partners and appropriate rates

^Δ NOT for use with *Estaprop XT* or *Dichlorprop DX*.

^{*} Rate above 0.81 L/acre may cause crop injury.

^{**} Barnyard grass also controlled.

^{***} Barnyard grass and Persian dandel also controlled. May be applied by air.

^{††} Rates for 500 g ai/L formulations are given. Other formulation strengths are also permitted.

^{†††} Additional adjuvants are not required.

[◇] All products except *Aurora*

Note: The above mixes are those listed on the *clodinafop* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Clodinafop manufacturers may also support mixes with pesticides that are not on the *clodinafop* labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

For optimum results, apply to actively growing weeds. Do not apply to crops or weeds that are stressed by hot or cool conditions, frost, drought, low fertility, water-saturated soil, disease or insect damage as crop injury and poor weed control may result.

Restrictions:

Rainfall: Within 30 minutes may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze or harvest treated crops for forage within 3 days of application.

Preharvest: Leave at least 60 days from application to harvest.

Re-cropping: No restrictions in the year following treatment.

Storage: May be frozen.

Aerial Application: May be applied by air.

Buffer Zones:

Application method	Buffer Zones (metres †) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground *	15	0
Aerial	72	76

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to page 15 to 16.

Hazard Rating:

240EC formulations:



Caution – Poison



Warning – Eye and Skin Irritant

NG formulations:



Caution – Eye and Skin Irritant

All except Ladder:



Warning – contains the allergen soy.

For an explanation of the symbols used here see page 11.

Curtail M

Herbicide Group
4 - clopyralid & MCPA
(Refer to page 38)

Company:

Nufarm Agriculture (PCP# 22764)

Formulation:

50 g/L clopyralid and 280 g/L MCPA ester formulated as an emulsifiable concentrate.

Container size - 8 L, 112 L, 960 L.

Crops and Staging:

Apply at the 3 leaf to just before the flag leaf stage of the following crops:

Barley	Timothy (established for seed, and hay or forage production)*
Canaryseed*	
Oat	Wheat (spring & durum)

Flax, solin (low linolenic acid flax) at 2 to 6 inches (5 to 15 cm) height.

**NOTE: Since these uses are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Those who apply these uses do so at their own risk.*

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

The following weeds are controlled at the 1 to 4 leaf stage unless specified:

At 0.61 L per acre (8 L jug treats 13 acres) for light infestations of:

Burdock	Ragweed
Canada thistle***	Shepherd's-purse**
Cocklebur	Stinkweed**
Field horsetail†	Sunflower annual
Flixweed**	Volunteer sunflower
Lamb's-quarters	Wild mustard
Plantain†	Wild radish
Prickly lettuce	Vetch

At 0.81 L per acre (8 L jug treats 10 acres) for medium to heavy infestations of the above weeds and:

Annual sow-thistle	Russian pigweed
Canada thistle***	Scentless chamomile**
Common groundsel	Smartweed
Dandelion*	Tartary buckwheat
Kochia (suppression only)**	Volunteer canola
Perennial sow-thistle†	Wild buckwheat
Redroot pigweed	

* Spring rosettes only.

** 2 to 4 leaf stage, (spring seedlings only for winter annuals).

*** Season long control, some regrowth may occur in the fall. Apply from the 4 inch (10 cm) to prebud stage.

† Top growth control only.

Application Information:

Water Volume:

Canary seed and timothy: 40 to 80 L per acre

All other crops: 40 to 60 L per acre.

Nozzles & Pressure: Use 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Select the nozzle and pressure combination that produces of ASABE coarse droplets while maintaining good coverage of foliage. Flat fan tips tilted forward at a 45° angle are recommended in flax.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

When weeds are stressed because of drought, flooding, hot or cool (less than 15°C) temperatures, weeds are not actively growing and control may be reduced. DO NOT apply to weeds stressed longer than 20 days from lack of moisture as poor control can result.

Tank Mixes:

Curtail M at 0.81 L per acre should be used in all tank mixes unless otherwise indicated. See labels for adjuvant rates.

Herbicides:

In spring wheat (including durum) and barley:

Achieve Liquid (0.20 L/acre) plus Turbocharge adjuvant

Assert (0.52 to 0.64 L/acre) plus water pH adjuster *Refine*
In spring wheat (not including durum) and barley:
Axial 100TC (0.24 L/acre) plus *Adigor* adjuvant.

Check product labels for additional crop staging restrictions.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *Curtail M* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours will reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated fields or cut for hay within 7 days of application.

Preharvest Interval: Leave 60 days between application and harvest.

Re-cropping: Wheat, barley, oats, rye, corn, flax, canola, forage grasses and mustard may be planted the year after application. DO NOT under-seed crops to forage legumes the year after treatment.

DO NOT seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Nufarm Agriculture Inc. representative or retailer for more information before seeding field peas following drought conditions in the previous year.

DO NOT sow *any other* crops until the second year after application. Apply manure bedded with straw from treated crops only to the crops listed above.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool (above 5°C), dry area. If product is frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	4

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Spray when winds are under 16 km/hr, but not dead calm.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution - Poison



Caution - Eye Irritant

For an explanation of the symbols used here see page 11.

Dicamba

Herbicide Group

4 - dicamba

(Refer to page 38)

Company:

BASF Canada (*Banvel II*) (*Banvel VM* - distributed by Engage Agro)

Gharda Chemicals Ltd. (*Oracle*, *VMD*) 480 *Dicamba*, *Hawkeye Power* - Distributed by Adjuvants Plus)

Formulation:

Banvel II (PCP# 23957), *Banvel VM* (PCP# 29249): 480 g/L dicamba formulated as a solution of a diglycolamine salt.

Oracle (PCP# 26722), *VMD 480 Dicamba* (PCP# 29251), *Hawkeye Power* (PCP# 29223): 480 g/L dicamba formulated as a solution of a dimethylamine salt.

Container sizes - 2 x 10 L jugs.

Crops, Rates and Staging:

Barvel II, Oracle, Hawkeye Power and VMD 480 Dicamba are registered for the all of the uses below. *Barvel VM* is only registered for pasture and rangeland uses below.

CROP	STAGE	RATE	
		(mL per acre)	Acres per 10 L
Spring wheat*	2 to 5 leaf.	93 to 117	107.5 to 85
Barley*	2 to 5 leaf.	93 to 117	107.5
Oats*	2 to 5 leaf.	93 to 117	107.5 to 85
Canaryseed*	3 to 5 leaf.	117	85
Winter wheat*	In spring 6 to 10 inches (15 to 25 cm) - prior to flag leaf	93 to 117	107.5 to 85
Spring rye*	2 to 3 leaf	93 to 117	107.5 to 85
Corn, field	Broadcast up to 8 inches (20 cm). When higher, use drop-nozzles.	243 to 505	41 to 20
Corn, field + 2,4-D	Apply no later than 2 weeks prior to tassle emergence and prior to 20 inches (50 cm).	117	85
Red fescue (for seed production)	Seedling: 2 inches (5 cm) tall. Established: up to the flag leaf stage.	243	41
Pastures	Established and actively growing	850 to 1,480 (0.85 to 1.48 L)	11.7 to 6.8
Seedling grasses (for seed and forage production): Fescue (creeping red, meadow, tall), Meadow foxtail, Orchardgrass, Smooth brome grass, Timothy, Wheatgrass (crested, intermediate, pubescent, slender, streambank, tall)	2 to 4 leaf	93 to 117	107.5 to 85
Fall stubble	Apply according to weed stage.	1000 (1.0 L)	10
Fall stubble + glyphosate	Apply according to weed stage.	500	20
Pre-seeding cereals	Apply according to weed stage.	127	79
Chemfallow + 2,4-D	Apply according to weed stage.	93 to 117	107.5 to 85
Chemfallow + glyphosate	Apply according to weed stage.	117 to 243	85 to 41

* Should be mixed with a tank mix partner for broad spectrum control

Weeds, Rates and Staging:

Apply to annual broadleaf weeds at the 2 to 3 leaf stage and to winter annual rosettes up to 2 in. (5 cm) across.

Dicamba applied alone at 93 to 117 mL per acre will control:

Cleavers (high rate only)	Perennial sow-thistle*
Cow cockle	Smartweed (green)
Corn spurry	Tartary buckwheat
Canada thistle*	Wild buckwheat
Lady's-thumb	

Dicamba at 0.25 to 0.5 L per acre will control:

Weeds listed above plus:

Canada thistle**	Perennial sow-thistle**
Canada fleabane	Pigweed (redroot, Russian)
Field bindweed**	Ragweed (common, false, giant)
Lamb's-quarters	
Mustard (hare's-ear, Indian, tumble, wild, wormseed)	

Dicamba at 0.85 L per acre in rangeland or 1.0 L per acre in summerfallow will control:

Weeds listed above plus:

Curled dock*	Goldenrod
English daisy	Tansy ragwort

Dicamba at 1.86 L per acre will control:

Weeds listed above plus:

Diffuse knapweed	Povertyweed
Goat's-beard	Sheep sorrel
Ground cherry	Thyme-leaved spurge
Pasture sage	

* Top growth only.

** Three consecutive years of treatment are required for complete control.

The following chart indicates weed and brush controlled by dicamba + 2,4-D mixes at the listed rates.

WEEDS	RATE (L/acre) [†]	
	Dicamba	2,4-D (600 g/L forms)
Poison ivy	0.67	0.76
Wild carrot	0.85	0.76
Aspen poplar	1.32	1.52
Prickly rose, western snowberry ^{†††}	1.48	1.52
	RATE (L/1000 L of water) ^{††}	
Alder, aspen poplar, cherry, western snowberry, wolf willow, wild rose	0.85	1.33

[†] Applied by broadcast sprayer.

^{††} Apply to the foliage and stems to the point of run-off using high volume equipment.

^{†††} Ester formulations of 2,4-D only.

Canada thistle, Perennial sow-thistle in summerfallow: Apply prior to the bud stage. Must be applied to thistle plants with 6 to 10 inches (15 to 25 cm) of new growth.

Canada thistle control in fall stubble: When thistles exhibit new growth and at least 2 weeks prior to a killing frost.

Refer to label for full lists of weeds controlled by dicamba plus tank mixes in cereals, pastures, summerfallow and other situations.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Brush control in pastures: When brush is actively growing and is 6 feet (2 m) in height or less (in spring or early summer). Growth greater than 2 meters may be cut and allowed to regrow prior to treatment.

Application Information:

Water Volume:

Preseeding burnoff: 20 to 45 L per acre.

Annual crops: at least 45 L per acre.

Pastures, summerfallow and stubble: 45 to 90 L per acre.

Corn: 90 to 140 L per acre.

Brush: high volumes to the point of run-off.

Nozzles and Pressure:

Broadcast application: Maximum 40 psi (275 kPa) with conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver even coverage of ASABE coarse droplets.

Brush Control: Use high volume spray equipment producing large droplets including, but not limited to, hand-wand, boomless nozzle and Radi-Arc technologies.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop damage (stunting, reduced seed set) can occur if the chemical is applied at any time other than the recommended stage. DO NOT apply to crop under stress from adverse environmental conditions, such as excess moisture, drought and disease. Apply when air temperature is between 10 and 25°C.

DO NOT apply:

- when there is a risk of severe temperature fall in the night;
- under high humidity, temperatures above 30°C, or fog conditions, to prevent drift to sensitive crops;
- when wind is blowing toward a nearby sensitive crop;
- when winds are gusty up to 5 mph (8 km/hr).

Tank Mixes:

Herbicides:

	Spring wheat	Winter wheat	Barley	Oats	Seedling grasses
2,4-D Amine (500g ai/L) (0.34 L/acre)	✓	✓	✓		✓
MCPA Amine (0.34 L/acre)	✓	✓	✓	✓	✓
MCPA K (0.44 L/acre)	✓	✓	✓	✓	✓
Sencor (0.11 to 0.17 L/acre)	✓		✓		
Ally (2 g/acre)	✓		✓		

In Canaryseed: MCPA amine (0.34 L/acre - 500 g ai/L formulation)

In Corn, Spring rye: 2,4-D amine (0.34 L/acre - 500 g ai/L formulation)

In Corn (Banvel II only):

Accent (13.5 g/acre) plus non-ionic surfactant

Option 351D (40 g/acre) plus Hasten adjuvant plus liquid 28-0-0 (Banvel II at 0.12 L/acre) (Manitoba only).

Option 2.25 OD (0.63 L/acre) plus liquid 28-0-0 (Banvel II at 0.12 L/acre) (Manitoba only).

In Chemical fallow, stubble: 2,4-D, glyphosate products.

In Red fescue: 2,4-D amine (0.61 L/acre - 500 g ai/L formulation)

In Preseeding burnoff: Glyphosate (0.38 L/acre - 360 g ai/L formulation)

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the dicamba labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for 12 hours.

Grazing and Harvest Intervals:

Canaryseed: Use only as birdseed.

Corn: DO NOT graze cattle or harvest for silage until 7 days after treatment of dicamba alone or for at least 12 weeks following dicamba tank mixes with other herbicides.

Cereals, seedling grasses, pasture: DO NOT harvest for silage for or graze lactating dairy cattle until 7 days after treatment. If treated vegetation has been consumed by dry dairy animals or meat animals within 30 days of dicamba application, feed the animal with untreated diet for 30 days before slaughter. Meat animals or dry dairy animals may graze or feed on treated pasture 3 days after dicamba application without restrictions on slaughter. Feed untreated forage within 3 days of slaughter.

Re-cropping: Grow only cereals, corn, soybeans or white beans the year after treatment with the 1.0 L per acre rate. Grow only cereals, corn, field beans, soybeans or canola the year after applications of 0.5 L per acre. If applications are made after September 1, or if dry weather persists after application, crop injury may occur the following spring.

Aerial Application: May be applied by air on cereals only. Use a minimum water volume of 8 L per acre.

Storage: May be stored at freezing temperatures.

Buffer Zones:

Application method	CROP	Buffer Zones (metres†) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Ground*	Barley, oats, rye, wheat, canary seed, seedling forage grasses	0	0	1
	Corn, established forage grasses, red fescue	1	1	4
	Stubble, fallow	1	1	5
	Pasture and rangeland	1	1	10
Winged airplane	Barley, oats, rye, wheat,	0	0	50
Helicopter		0	0	45

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Buffers are not required for handheld and backpack applications.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution - Poison



Warning - Eye Irritant

For an explanation of the symbols used here see page 11.

Dicamba/Mecoprop/ MCPA

Herbicide Group
4 - dicamba, mecoprop-p
& MCPA
(Refer to page 38)

Company:

Syngenta (*Target* - PCP# 28028)
United Agri Products (*Sword* - PCP# 27892)
IPCO (*Tracker XP* - PCP# 27790)

Formulation:

275 g/L MCPA + 62.5 g/L mecoprop-p + 62.5 g/L dicamba formulated as a solution.

Container size - 2x10 L and 160 L (*Target*), 500 L (*Sword*), 1000 L (*Sword*).

Crops and Staging:

All Products:

Cereals:

CROP	STAGE
Barley	2 to 4 leaf (3 leaf for best crop safety)
Canaryseed, Oats, Spring wheat (including durum)	2 to 5 leaf (3 to 4 leaf for best crop safety)
Winter wheat	Spring application only; up to 12 inches (30 cm) high (top leaf extended)
Summerfallow	Fall stubble

Target and Sword only:

Seedling grasses grown for forage only (NOT for seed production):*

Apply at the 2 to 4 leaf stage.

Creeping red fescue	Orchardgrass
Crested wheatgrass	Smooth brome grass
Intermediate wheatgrass	Timothy
Meadow foxtail	

Established grasses for forage only (NOT for seed production):*

Apply up to flag leaf stage.

Brome grass (meadow, smooth)	Orchardgrass
Fescue (creeping red, meadow, tall)	Timothy
Kentucky bluegrass	Wheatgrass (crested, intermediate, pubescent, slender, streambank, tall, western)
Meadow foxtail	

* NOTE: Use only one application per year by ground. Since applications to forage grasses in western Canada has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to forage grasses are at the risk of the user.

Weeds and Staging:

Weeds controlled at the 2 to 3 leaf stage unless otherwise indicated:

Annual sow-thistle	Night-flowering catchfly
Ball mustard	Perennial sow-thistle*
Canada thistle (6 to 8 inches (15 to 20 cm) and actively growing)*	Prostrate pigweed
Cleavers (1 to 2 whorls)	Redroot pigweed
Common ragweed	Russian thistle (less than 2 inches-5 cm)
Corn spurry	Shepherd's-purse
Cow cockle	Stinkweed
Field bindweed*	Tall mustard
Flixweed	Tartary buckwheat
Green smartweed	Volunteer buckwheat
(including lady's-thumb)	Volunteer canola
Hedge bindweed*	Volunteer sunflowers
Hemp-nettle (less than 2 pairs of true leaves)	Wild buckwheat
Knotweed	Wild mustard
Kochia	Wormseed mustard
Lamb's-quarters	Yellow mustard

* Top Growth Control only

Rates:

0.4 to 0.6 L per acre (10 L treats 25 to 16.7 acres).

Use the higher rate under adverse weather conditions, when weed density is high, for cleavers control, winter annual control and for suppression of Canada thistle and perennial sow-thistle.

Although dicamba/mecoprop-p/MCPA is registered up to the 5 leaf stage of the crop for the rates listed here, the low rate should be used when the crop is at the 5 leaf stage for optimum crop safety.

For Canada thistle, post-harvest or summerfallow application, use 0.81 L per acre (one 10 L container treats 12.4 acres).

Application Information:

Water Volume:

Ground: Minimum 40 L per acre.

Aerial: Minimum of 12 L per acre

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE coarse* droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Hot and dry or cold and wet weather prior to spraying may result in reduced weed control and increased crop injury. Do not apply within 2 weeks of a killing frost.

Tank Mixes:

Herbicides:

Spring wheat (including durum)

Horizon NG * (label rates - no adjuvant required)

Wheat and Barley:

Sencor or *linuron* for chickweed control.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

* Target only.

Note: The above mixes are those listed on the *dicamba / mecoprop-p/MCPA* labels only. To check for other possible mixes see the blue chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Activity may be reduced if rainfall occurs within 3 hours of application. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT graze or harvest for livestock feed within 7 days of application.

Preharvest: Leave at least 80 days from application to harvest.

Re-cropping: No restrictions the year after application.

Aerial Application: All may be applied by air.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Crops	Buffer Zones (metres [†]) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Ground *	Standing Crops	1	1	5
	Fallow and stubble	1	1	5
Fixed wing airplane	Cereals	1	0	60
	Canaryseed	1	0	75
	Forage	1	0	75
	Fallow and stubble	5	1	100
Helicopter	Cereals	1	0	50
	Canaryseed	1	0	60
	Forage	1	0	60
	Fallow and stubble	4	1	80

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Buffers are not required for handheld and backpack applications.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution - Poison



Warning - Eye Irritant

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Dichlorprop/2,4-D

Herbicide Group
4 - dichlorprop & 2,4-D
 (Refer to page 38)

Company/Products:

United Agri Products (*Turboprop*)
 Nufarm Agriculture (*Estaprop XT*)
 IPCO, Loveland Products (formerly Viterra) (*Dichlorprop-D*)
 IPCO (*Dichlorprop-DX*)

Formulation:

Turboprop (PCP# 27967); *Dichlorprop-D* (PCP# 27966): 300 g/L of dichlorprop and 282 g/L of 2,4-D ester formulated as an emulsifiable concentrate.

Estaprop XT (PCP# 29660); *Dichlorprop-DX* (PCP# 29664): 210 g/L of dichlorprop-P* and 400 g/L of 2,4-D ester formulated as an emulsifiable concentrate.

* NOTE: dichlorprop-P is a more active version of dichlorprop.

Container sizes:

Turboprop, *Dichlorprop-D* - 2 x 10 L, 115 L

Estaprop XT - 2 x 9.7 L, 97.1 L, 466.1 L.

Dichlorprop-DX - 2 x 10L, 115 L

Crops and Staging:

Wheat (spring, durum) and barley - 4 leaf until prior to the emergence of the flag leaf.

Winter wheat - in spring after the initiation of tillering but prior to the emergence of the flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Rates:

Turboprop; *Dichlorprop-D*: 0.71 L per acre

One 10 L container treats 14 acres

One 10.6 L container treats 15 acres

Estaprop XT (PCP# 29660); *Dichlorprop-DX* (PCP# 29664): 486 mL per acre.

One jug treats 20 acres

How it Works:

Refer to Table 2 on page 40.

Weeds and Staging:

Treat weeds when young and actively growing and before they are shielded by the crop. Additional stage restrictions are indicated.

Annual sow-thistle	Prickly lettuce (2 to 12 leaf)†
Ball mustard	Ragweed
Bluebur	Redroot pigweed
Burdock	Round-leaved mallow
Canada thistle*	Russian pigweed
Cocklebur	Russian thistle♦♦♦
Curled dock*	Shepherd's-purse
Dandelion ***	Smartweed♦♦
Dog mustard	Stinkweed
Flixweed	Stork's-bill
Hare's-ear mustard	Tartary buckwheat♦♦
Indian mustard	Toadflax**
Kochia♦♦♦	Tumble mustard
Lady's-thumb♦♦	Volunteer canola♦♦
Lamb's-quarters	Volunteer sunflower♦♦
Night-flowering catchfly♦	Wild buckwheat♦♦
Oak-leaved goosefoot	Wild mustard
Perennial sow-thistle*	Wormseed mustard

* Top growth control only

** Suppression only. Treat before the majority reach 6 inches (15 cm).

*** *Estaprop XT* and *Dichlorprop-D* only - season long control in winter wheat.

♦ Spring annuals only

♦♦ Treat prior to the 4 leaf stage

♦♦♦ Treat up to 2 inches (5 cm)

† *Estaprop XT* and *Dichlorprop-D* only in winter wheat.

Application Information:

Water Volume:

Ground: 20 to 97 L per acre*. Use a minimum of 40 L of water per acre to reduce the risk of drift.

Aerial: Minimum 12 L per acre.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver proper coverage with ASABE coarse droplets or larger.

* May vary by product. Check label closely.

Effects of Growing Conditions:

Applications made under dry conditions may result in reduced control. Crops under stress from adverse environmental conditions, such as excess moisture, frost or drought, may be injured. Best weed control when adequate soil moisture is present and warm temperatures prevail. DO NOT apply when daytime temperatures exceed 27°C.

Tank Mixes:

Herbicides: *Estaprop XT, Dichlorprop DX and Turboprop only*

Tank Mix Partner (Mixed at label rates unless oth- erwise indicated)	CROPS			
	Spring wheat	Durum	Winter wheat	Barley
<i>Avenge</i> *	•	•	•	•
<i>Imazamethabenz</i> ^Δ	•	•		•
<i>Clodinafop</i> ^Δ	•	•		
<i>Fenoxaprop</i> ^Δ	•	•		•†
<i>Thifensulfuron/ tribenuron</i> ^{Δ††}	•	•	•	•
<i>Tralkoxydim</i> ^Δ	•	•	•	•

* Refer to *Avenge* label for a list of tolerant wheat varieties.

† *Cordon, Vigil WB and WildCat* only.

†† *Estaprop XT* only.

^Δ Manufacturers may only support mixes with specific products. Contact the manufacturer for more information.

Note: Always refer to the label or the page for the tank mix partner in this guide for additional restrictions on staging and varieties.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on *dichlorprop+2,4-D/dichlorprop-P+2,4-D* labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: Leave 12 hours before entering treated fields.

Grazing: DO NOT graze the treated crop or harvest for hay or feed within 40 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Preharvest Interval: Leave 40 days from spraying until harvest of winter wheat and 60 days for other crops.

Re-cropping: No restrictions the year after application. Fields treated with *Estaprop XT* may be replanted after a minimum of 30 days.

Aerial Application: May be applied by air. Refer to specific product labels for full details for application by air.

Storage: May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat**
	Less than 1 m	Greater than 1 m	
Ground	1	1	1
Fixed wing aircraft	10 (5††)	1	45 (30††)
Helicopter	10 (3††)	1	40 (30††)

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

** Handheld or backpack sprayers do not require a buffer zone.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

†† *Dichlorprop - DX and Estaprop XT*.

Tank Cleaning:

Manufacturers of this product warn that even after thorough cleaning, the use of a sprayer that has come in contact with this product may cause damage to susceptible crops.

DO NOT use spray equipment to apply other pesticides to crops sensitive to these products. To clean sprayer, rinse all parts several times with water, then fill sprayer with a water/ammonia solution (1 L of a 3% household ammonia solution per 100 L of water) and let stand for 24 hours. Rinse several times after with clean water.

DO NOT clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk.

Hazard Rating:



Warning – Poison



Warning – Skin Irritant (*Turboprop*)



Caution – Skin Irritant (*Dichlorprop-D*)

For an explanation of the symbols used here see page 11.

Diquat

Herbicide Group

22 - diquat

(Refer to page 38)

Company:

Syngenta Canada

Formulation:

Reglone (PCP# 26396) and **Desica (PCP# 30488)** 240 g/L diquat formulated as a solution:

Container size - 2x10 and 115 L.

Reglone Ion (PCP# 31058): 200 g/L diquat formulated as a solution. Comes with a built-in-adjuvant

Container size - 2x10 L, 115 L, 450 L.

Crops and Staging:

Diquat is used to dry immature green material at top of indeterminate crops and green weeds to facilitate harvest. *Diquat* will not speed maturity of green crops. Treatment before the recommended stage can result in reduced yield and quality. For *Reglone* and *Desica* only add 0.1 L of *Agral 90* or *Agsurf II* per 100 L of spray solution for all applications except those indicated for *Reglone Ion*

CROP	STAGE	RATE (L/ACRE)	
		Ground	Aerial
Dry beans, soybeans	Crop has lost 80 to 90 percent of leaves and 80 percent of pods are yellow.	0.5 to 0.7 or 0.6 to 0.8†	0.7 to 0.9 or 0.8-1.0†
Faba beans	Most plants are ripe and dry. Pods fully filled, bottom pods are tan or black in colour.	(use high rate for dense crop, heavy weed infestations)	(use high rate for dense crop, heavy weed infestations)
Flax, solin	75 percent of bolls brown		
Lentils	Lowest pods are light brown and rattle when shaken.		
Mustard (condiment type only)	75 percent of seed has turned brown		
Canola*	80-90 percent of seed has turned brown.		
Peas	Bottom pods are ripe and dry, seeds detached from pods.		
Chickpeas [◇]	Plants yellow, pods mature, seeds changed colour and detached from pods.		
Sunflowers	Backs of sunflower heads and bracts are turning yellow and seed moisture is 20 to 50 percent.		
Potatoes†† (some top growth and / or some weeds)	Two weeks prior to harvest	0.7 to 0.9**	2 applications required. First application - 0.7 to 0.9 L/acre** Second application (4 to 5 days later) at 0.5 L/acre
Potatoes†† (dense crop, heavy weed infestations)		1.4**	
Alfalfa, bird's-foot trefoil, red and white clover (for seed production only)***	Pods are ripe but before shattering. Harvest within 7 days.	0.7 to 1.1 or 0.8 to 1.3† (use high rates for dense crops, heavy weed infestations)	0.7 to 1.1 or 0.8 to 1.3† (use high rates for dense crops, heavy weed infestations)

* This product can cause shattering losses in canola. It should only be used on *B. napus* canola to assist in the harvest of a severely lodged crop.

- ** Use high rate for dense or immature vines.
 *** DO NOT use on forage legumes that have been treated with a residual herbicide in the previous 12 months.
 ◇ Aerial applications to chickpea - 0.7 L per acre only.
 † For *Reglone Ion* formulation
 †† DO NOT use *Reglone Ion* on potatoes.

Application Information:

Water Volumes:

Ground applications: 91 to 222 L per acre. Use 222 to 445 L per acre on potatoes.

Aerial applications: 18 L per acre.

Use the highest water volumes when crop canopy is heavy or if weed growth is dense.

Nozzles and Pressure: 30 to 40 psi (210 to 275 kPa) when using conventional Flat fan nozzle tips are recommended for proper coverage. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Best results under cloudy conditions or in evening. Shattering losses can increase if heavy winds, rain or hail occur after the crop has dried down.

Tank Mixes:

Herbicides: None registered.

Insecticides: None registered.

Fungicides: Fungicides may be added when applying *diquat* to potatoes for vine killing.

Fertilizers: None registered.

Restrictions:

Rainfall: Within 15 minutes may reduce effectiveness.

Re-entry: Leave 24 hours before entering treated fields.

Grazing: Crop residues remaining after harvest may be fed to livestock.

Preharvest Interval: Wait 4 to 7 days before harvesting faba bean and lentil. DO NOT exceed 7 days to harvest when treating forage legumes. Wait at least 7 days before harvesting lupins. Wait 7 to 10 days before combining canola and mustard, but do not wait longer than 14 days. Wait 15 to 20 days for sunflowers. Harvest flax and peas when sample tests dry.

Re-cropping: No restrictions the year after application.

Aerial Application: May be applied by air in a minimum of 18 L per acre water volume.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Crops	Buffer Zones (metres [†]) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Ground	Potatoes	10	5	5
	Other crops under Crops: section	5	3	3
Winged aircraft	Potatoes	200	100	100
	Beans, Legume forage seed	150	80	90
Helicopter	Potatoes	125	65	80
	Beans, Legume forage seed	100	55	70

See page 29 for an explanation of the different habitats.

* Buffer zones for ground applications can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

When finished spraying *diquat*, rinse the sprayer out with clean water. Run through pump, lines and nozzles. Drain tank by spraying out on an untreated portion of a crop on which the product is registered, or by spraying on uncropped land. Refill sprayer with water and *Agral 90*, or *Agsurf II* at 0.6 L per 1,000 L spray solution. Run the solution through lines and boom. Spray out, then refill with clean water. Leave equipment standing overnight, then drain water out.

Refer to page 15 to 16 for additional information.

Hazard Rating:



Warning - Poison



Caution - May cause eye damage

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Distinct

Herbicide Group

4 - dicamba

19 - diflufenopyr

(Refer to page 38)

Company:

BASF Canada (PCP#25811)

Formulation:

One case of *Distinct* contains:

Distinct (PCP#): 20% diflufenopyr and 50% dicamba, sodium salt formulated as a water dispersible granule. Container size - 2 x 2.3 kg.

Crops and Staging:

Chemfallow and Post Harvest applications

Weeds, Rates and Staging:

Distinct must be tank-mixed with glyphosate and *Merge* adjuvant must be added at 200mL per acre.

Distinct at 58 g per acre (40 acres per jug) controls:

Kochia

Dandelion*

Distinct at 114g per acre (20 acres per jug) controls:

Weeds listed above plus:

Canada thistle*

Redroot pigweed

Common cocklebur

Tall waterhemp

Common ragweed

Velvetleaf

Lady's-thumb

Wild buckwheat

Lamb's-quarters

*Top growth control only.

A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 20 to 40 L per acre. High water volumes are required for adequate coverage, particularly when weed densities are high or weed staging is large.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of a *ASABE coarse* droplets. Use 50 mesh (or coarser) filter screens.

Effects of Growing Conditions:

DO NOT spray if temperatures of +5°C or less are forecast within 3 days of application. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

Glyphosate (0.5 to 1.0L per acre of 360 g/L formulation.)

Restrictions:

Rainfall: Rain within 3 hours may reduce control.

Re-entry: Leave 12 hours before entering treated fields.

Grazing: DO NOT graze or cut for 75 days.

Re-cropping: A plant back interval of 30 days is required for the planting of rotational crops. Consult BASF for further information on rotational cropping.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place above 5°C.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only	15	15	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Distinct can cause injury to sensitive crops at very low concentrations. Sprayers used to spray this product should be flushed out immediately after use.

Use 'Method B' on page 15-16 to clean sprayers after using *Distinct*.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Dual II Magnum

Herbicide Group
15 - metolachlor
 (Refer to page 38)

Company:

Syngenta Canada (PCP#25729)

Formulation:

915 g/L s-metolachlor formulated as an emulsifiable concentrate. Container size - 12 L.

Crops and Staging:

Pre-plant incorporated. In areas with good rainfall or under irrigation, *Dual II Magnum* may be applied as a pre-emergence surface treatment. When applied to the soil surface, after planting but prior to emergence, at least 0.5 inches of water (1.25 cm) is required within 10 days of application for proper activity. Refer to product label for more specific information on timing and rates of applications for each crop type.

Corn (field, sweet, silage)	Potatoes
Dry beans (white, kidney, and pinto)*	Soybeans
	Sweet white lupins

* Beans should be planted at least 4 cm deep to avoid crop injury. Dry bean varieties vary in their tolerance to *Dual II Magnum*. Test a limited acreage on all new varieties first.

Weeds and Staging:

Pre-emergent and Pre-Plant Incorporated Treatments:
 Apply prior to weed emergence.

American nightshade	Old witch grass
Barnyard grass	Redroot pigweed*
Eastern black nightshade	Yellow foxtail
Green foxtail	Yellow nutsedge**

* Suppression only.

** Pre-plant incorporated treatment only.

Rates:

0.47 to 0.7 L per acre (12 L treats 24 to 17 acres).

Use higher rates on heavy textured soils or when high populations of weeds are expected.

DO NOT apply to soils with less than 1% or more than 10% organic matter.

Make only one application per season.

Refer to product label for more specific information on timing and rates of applications for each crop type.

Application Information:

Water Volume: A minimum of 60 L per acre.

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Use flat fan nozzles, 50 mesh screens.

Incorporation:

Apply to a firm seed bed free of large clods or lumps. If using tandem disks, set disks to work the soil at a depth of 4 inches (10 cm) and operate at a speed of 6 km/hr (4 mph). If using an S-tine cultivator, set the implement to work the soil to a depth of 4 inches (10 cm) and operate at a speed of 10 km/hr (6 mph). Incorporation equipment should include rolling or western harrows.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

A moderate rainfall or equivalent irrigation is required within 10 days to activate pre-emergent surface treatments. If rain does not occur, a shallow cultivation or use of a rotary hoe is necessary. Drought conditions that persist after any application may reduce annual grass control. On sandy soils, heavy rainfall following application may cause leaching of *Dual II Magnum*, resulting in reduced weed control.

Tank Mixes:

Herbicides:

In Corn: *AAtrex* in both PPI and pre-emergent applications.

In Soybeans: *Sencor*, and *glyphosate*, in both PPI and pre-emergent applications.

Fertilizers: May be applied with liquid fertilizer. May be impregnated onto dry bulk fertilizers (except nitrate fertilizers, superphosphate fertilizers or limestone).

Insecticides: None registered.

Note: The above mixes are those listed on the *Dual II Magnum* label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: When applying as a pre-emergent surface treatment, 0.5 inches (1.25 cm) of rain or irrigation is required after application for proper activity.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze the treated immature crop or cut for hay. In corn, immature means before ear emergence.

Preharvest Interval: DO NOT harvest corn within 80 days of post-emergent application.

Re-cropping: In the year of treatment, seed only corn, soybeans, white beans, potatoes, snap beans, lima beans, processing peas, sweet white lupins, or (a minimum of 4.5 months after application) winter cereals. If *Dual II Magnum* has been applied in a tank mix with another product, consult those products' labels for additional re-cropping restrictions.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones: Leave a buffer zone of 29 meters between last spray swath and the edge of important wildlife habitats such as wetlands, sloughs and water bodies.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Warning - Eye Irritant



Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

DyVel

Herbicide Group
4 - dicamba & MCPA
(Refer to page 38)

Company:

BASF Canada (PCP#16545)

Formulation:

84 g/L of dicamba and 336 g/L of MCPA K+ formulated as a solution.

Container size - 10 L, 55 L, 110 L, 1000 L

Crops and Staging:

Spring wheat (including durum), barley or oats - 2 to 5 leaf stage.

Winter wheat - apply in spring when crop is 6 to 10 inches (15 to 25 cm) tall but before shot blade stage.

Note: Crop damage can occur if applications are made at other than the recommended crop stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled in the 2 to 4 leaf stage unless otherwise stated:

Ball mustard	Pigweed (prostrate, redroot, Russian)
Burdock	
Cleavers (suppression only)	Ragweed (common, false, giant)
Cocklebur	Russian thistle
Corn spurry (2 to 3 leaf)	Shepherd's-purse
Cow cockle (2 to 3 leaf)	Smartweed (green, Lady's-thumb)
Flixweed	Stinkweed
Hemp-nettle (2 to 3 leaf)	Tartary buckwheat
Kochia	Wild buckwheat
Lamb's-quarters	Wild radish
Mustards (Hare's ear, Indian, Tumble, Wild, Wormseed)	Volunteer canola (2 to 4 leaf)
	Volunteer sunflowers

Top growth control:

Canada thistle	Perennial sow-thistle
----------------	-----------------------

Rate:

0.51 L per acre (one 10 L jug treats 19.7 acres).

Application Information:

Water Volume:

Ground: 40 L per acre.

Aerial: Minimum 8 L per acre

Nozzles and Pressure: Maximum 40 to 45 psi (275 to 310 kPa) when using conventional flat fan nozzles. To reduce the risk of drift damage to sensitive non-target crops when using conventional nozzles, 20 to 30 psi (150 to 200 kPa) as well as higher water volumes are recommended. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

For best weed control, apply when temperature is between 10 and 25°C. DO NOT treat crops under stress from excessive moisture or drought. To avoid crop injury, DO NOT apply when temperature is expected to exceed 30°C, or when there is a risk of a severe drop in overnight temperature.

Tank Mixes:

None registered.

Note: No mixes are listed on the DyVel label with currently marketed products. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Avoid applying this product when heavy rain is forecast. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated crop or cut for hay within 30 days of application.

Preharvest Interval: Leave 60 days between application and harvest.

Re-cropping: No restrictions the year after treatment.

Aerial Application: May be applied by air.

Storage: May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	4
Fixed wing aircraft	1	0	60
Helicopter	1	0	50

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to "Method A" in the general tank cleaning section on page 15 to 16. If mixed with other pesticides, the cleaning method above should be combined with the method recommended for the tank mix partner if different from above.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

DyVel DSp

Herbicide Group
4 - dicamba, 2,4-D & mecoprop
(Refer to page 38)

Company:

BASF Canada (PCP# 27856)

Formulation:

110 g/L dicamba, 295 g/L 2,4-D amine and 80 g/L mecoprop-p formulated as a solution.

Container size - 10 L, 55L, 100 L.

Crops, Rates and Staging:

CROP	STAGE	RATE	
		(L PER ACRE)	ACRES PER 10L JUG
Spring wheat (including durum)	3 to 5 leaf	0.34 to 0.45	29 to 22
Barley	2 to 3 leaf	0.34	29
Winter wheat	Before crop is 12 inches (30 cm) tall in spring	0.34 to 0.45	29 to 22
Corn (field)** (DO NOT apply to sweet corn)	Before corn reaches 6 inches (15 cm) in height with the top leaf extended or by directed spray with drop nozzles once over 12 inches (30 cm).	0.34 to 0.45	29 to 22
Native range and permanent grass pasture*	Established	1.3	7.7
Fall stubble, summerfallow	Stage according to weed	0.45 to 0.71	22 to 14

* Legumes will be severely injured by this application.

** NOTE: Under environmental stress corn will become brittle for 2 weeks after application. In-field mechanical processes and strong winds may cause stalk lodging during that time.

Applications outside the recommended stage may result in crop injury.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

The low registered rate for each crop will control the following weeds at the 2 to 3 leaf stage unless otherwise indicated:

Annual smartweed (including lady's-thumb)	Mustards (wild, ball, tall, wormseed, yellow)
Annual sow-thistle	Pigweed (prostrate, redroot)
Cocklebur	Russian thistle
Common ragweed	Stinkweed*
Corn spurry	Volunteer canola (2 to 4 leaf, prior to bolting)
Hedge bindweed	Volunteer tame buckwheat
Knotweed	Wild buckwheat
Kochia	
Lamb's-quarters	

Use the high registered rate for each crop to control the following weeds:

Canada thistle (top growth only)***	Flixweed*
Cleavers (1 to 2 whorls)**	Jerusalem artichoke
Cow cockle	Round-leaved mallow**
Field bindweed†	Shepherd's-purse*
Velvetleaf	Tartary buckwheat

† Apply when actively flowering.

* Rosette stage in winter wheat.

** Suppression only.

*** Canada thistle should be treated when 6 to 8 inches (15 to 20 cm) of new growth is present in Fall Stubble and in the early bud stage in Summerfallow.

Rates for Native Range and Pasture will control:

Alder	Poison ivy
Bull thistle	Ragwort
Chicory	Sheep laurel
Goat's-beard	White cockle

The high rate for each crop should be used for all weeds under adverse growing conditions, when weeds are at an advanced stage of growth or when weed densities are high. Guidelines are not provided for weed densities under light or heavy infestations. When in doubt as to the infestation level, use the high rate or contact the manufacturer.

NOTE: It is possible that poisonous plants such as ragworts, hemlocks and death camas could be more palatable to livestock after treatment with DyVel DSp. Suitable precautions should be taken to avoid livestock access when such plants are present.

Application Information:

Water Volume:

Cereals: Minimum 40 L per acre.

Corn: 81 to 142 L per acre.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE *coarse* classification or larger droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crops under stress from excess moisture, drought or disease may suffer a setback when this herbicide is applied. DO NOT apply when temperature exceeds 27°C or when relative humidity is high. Stubble treatments for thistle control in fall should be made at least 2 weeks prior to killing frost.

DO NOT apply *DyVel DSp* at wind speed greater than 5 mph (8 km/hr).

Tank Mixes:

Herbicides:

In corn:

AAtrex Liquid (0.91 L/acre).

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *DyVel DSp* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for opti-

mum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information. DO NOT irrigate for 24 hours after application.

Re-entry: DO NOT enter fields for at least 12 hours for field corn and leave 14-days from application to hand harvest sweet corn.

Grazing: DO NOT harvest for livestock feed within 30 days of application. DO NOT permit lactating dairy animals to graze fields within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Pre-harvest Interval: Leave 30 days between application and harvest.

Re-cropping: No restrictions the year after treatment.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones: If there are sensitive plants within 400 m, apply only when there is a light breeze away from the sensitive area. DO NOT contaminate wetlands or water used for domestic or livestock consumption, irrigation or natural habitat.

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Cropland	1	1	5
Range and Pasture	1	1	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

No cleaning information provided on the label. Refer to 'Method B' in general tank cleaning section on page 15 to 16.

Hazard Rating:



Warning – Poison



Warning – contains the allergen soy

For an explanation of the symbols used here see page 11.

Eclipse III

This product is a prepackaged tank mix of *Eclipse III A* (Lontrel page 216) and *Eclipse III B* (Vantage Plus Max II pages 178). Information listed is restricted to Crop, Weeds and Rates. For other detailed information on the component products see the product pages listed above.

Herbicide Group
4 - clopyralid
9 - glyphosate
(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

Eclipse III A (PCP# 29032): 360 g/L clopyralid formulated as a solution.

Eclipse III B (PCP# 29033): 480 g/L glyphosate present as a dimethylamine (DMA) salt and formulated as a solution.

Container size -

Eclipse III A: 4.45 L jug

Eclipse III B: 2 x 7.5 L jugs

Crops and Staging:

Glyphosate tolerant canola varieties only in the 2 to 6 leaf stage. Some yellowing may occur when applied at the 4 to 6 leaf stage. This effect is temporary and will not influence crop growth, maturity or yield.

Weeds and Staging:

No staging is specified on the label.

The weeds controlled by glyphosate at 180 g ae per acre plus:

Annual broadleaf weeds:

Chickweed	Shepherd's-purse
Corn spurry	Smartweed
Cow cockle	Wild tomato
Kochia	Volunteer canola*
Night-flowering catchfly	

Perennial weeds (season long control):

Canada thistle	Perennial sow-thistle**
Dandelion less than 15 cm diameter**	Quackgrass
Dandelion greater than 15 cm diameter***	

* Not including glyphosate tolerant (Roundup Ready) varieties.

** Top growth only.

*** Suppression only.

Rates:

Eclipse III A: at 112 mL per acre

Eclipse III B: at 375 mL per acre

One case treats 40 acres.

To prepare spray solution, add *Eclipse III A* to the spray tank. Once it is half filled with water, add *Eclipse III B* as the remaining water is added to the tank.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift. DO NOT use with galvanized sprayer tanks since explosive hydrogen gas can be produced.

Restrictions:

Re-cropping: Wheat, oat, barley, rye (not underseeded to legumes such as alfalfa and clover), forage grasses, flax, canola, mustard and field pea* can be grown the year after application. Apply manure bedded with straw from treated crops only to the crops listed above excluding field pea.

*DO NOT seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field pea following drought conditions in the previous year.

Aerial application: DO NOT apply by air.

Edge Granular

Herbicide Group
3 - ethalfluralin
(Refer to page 38)

Company:

Dow AgroSciences (PCP# 20980)

Formulation:

5% ethalfluralin formulated as a granular.
 Container size - 25 kg or 544 kg.

Crops and Staging:

Edge can be applied prior to seeding the following crops:

Seedling alfalfa (seed production only)	Lentils (fall application only) [†]
Canola	Mustard (yellow only)
Caraway	Peas
Coriander	Safflower
Dry beans (white or kidney)	Soybeans
Fababeans	Sunflowers

[†] See special instructions for lentils section below

Weeds and Staging:

For pre-emergent control of the following weeds:

Grasses:

Barnyard grass	Volunteer barley*
Crabgrass	Volunteer spring wheat*
Foxtail (green & yellow)	Wild oat*

Broadleaf Weeds:

Cleavers*	Lamb's-quarters
Chickweed	Nightshade*
Corn spurry	Prostrate pigweed
Cow cockle	Purslane
Hemp-nettle*	Redroot pigweed
Kochia	Russian thistle*
Lady's-thumb*	Wild buckwheat

* Suppression only.

Rates:

TIME OF APPLICATION	RATE (KG PER ACRE)				
	LIGHT TEXTURED SOILS		MEDIUM TO HEAVY TEXTURED SOILS		
	2-6% Organic Matter / Dark Brown-Black	6-15% Organic Matter Deep Black	2-4% Organic Matter Dark Brown	4-6% Organic Matter Black	6-15% Organic Matter Deep Black
Spring	6.9	8.9	6.9	8.9	8.9 - 11.3
Fall *	8.9	11.3	8.9	11.3	11.3

* To conserve crop residue, one incorporation may be completed in the fall and the second incorporation completed in the spring prior to planting.

DO NOT apply to peat soils, mineral soils containing less than 2 percent organic matter or soils containing greater than 15 percent organic matter. Application to eroded knolls or grey-wooded soils with highly variable texture or organic matter may result in reduced crop stand, delayed development or reduced yields in either the treated or rotational crop.

To reduce the possibility of injury to the treated crop, use good quality certified seed. Seed shallow into a warm,

moist, firm seedbed using recommended agronomic practices that will promote rapid and even crop germination and emergence.

[†] Special instructions for lentils:

Registered for use on lentils for fall application only. One incorporation must be completed in the fall. Seeding depth is critical - DO NOT seed more than 1.5 inches (4 cm) deep. Avoid loose seedbeds and planting into cold soils.

Application Information:

Equipment: Apply using a calibrated granular applicator.

Incorporation: Two incorporations are required at right angles for thorough mixing. The first incorporation must be completed within 24 hours of application. Delay the second incorporation for at least three days after the first. When applying *Edge* in the fall, it is preferred that both incorporations be done in the fall. The second incorporation may be delayed until spring to conserve trash; however, both incorporations must be done to the same depth.

Incorporate with a tandem disc, discer or field cultivator (Vibrashank type). Cultivators should have 3 to 4 rows of sweeps spaced 8 inches apart and staggered so that no soil is left unturned. Set equipment to work at a depth of 3 to 4 inches (8 to 10 cm). Operate disc implements at 4 to 6 mph (7 to 10 km/hr), cultivators at 6 to 8 mph (10 to 13 km/hr).

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crops stressed by cold weather, excessive moisture or drought may be injured by *Edge*. Dry soil conditions between application and emergence may result in decreased weed control.

Tank Mixes:

Not applicable.

Restrictions:

Rainfall: No effect once incorporated.

Grazing: DO NOT graze or cut treated crops for livestock feed prior to crop maturity.

Re-cropping: DO NOT grow sugar beets, oats, and small-seeded annual grasses such as timothy, canaryseed and creeping red fescue in rotation following a crop treated with *Edge*. DO NOT seed wheat as a rotational crop onto land that has been treated with trifluralin and/or ethalfluralin at oilseed/special crop/barley rates for two consecutive crop years. Thinning of crop may occur in areas that have received abnormally low amounts of precipitation or in crops that are emerging slowly.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen. DO NOT expose to prolonged sunlight or heat.

Buffer Zones: Toxic to fish and other aquatic organisms. DO NOT contaminate water bodies or wetland areas.

Tank Cleaning:

Not applicable.

Enforcer D

Herbicide Group
4 - fluroxypyr and 2,4-D
6 - bromoxynil
(Refer to page 38)

Company:

Nufarm Agriculture (PCP#30690)

Formulation:

80 g/L fluroxypyr, 190 g/L bromoxynil and 240 g/L 2,4-D ester formulated as an emulsifiable concentrate.

Container size: 2 x 10 L, 120 L or 480 L.

Crops and Staging:

Spring wheat (including durum) and barley at the 4 leaf to early flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Apply to emerged weeds up to the 6 leaf stage unless otherwise indicated.

Weeds controlled at the 0.24 L per acre rate:

Kochia (up to 5 cm tall)	Volunteer canola
Lamb's-quarters	Wild mustard

Weeds controlled at the 0.48 L per acre rate:

Weeds listed above plus:

Redroot pigweed	Wild buckwheat
Russian thistle	

Application Information:

Water Volume: Minimum 20 to 40 L per acre. Use the higher volume when there is a heavy crop canopy or weeds are at an advanced stage.

Nozzles and Pressure: Use 40 psi (275 kPa) if applying without drift reduction nozzles. Drift reduction nozzles may require higher pressures for proper performance. Select the nozzle and pressure combination that produces of ASABE *coarse* droplets while maintaining good coverage of foliage.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Optimum activity is experienced between 12 to 24 °C when weeds are actively growing. Weeds may not be actively growing and as a result reduced activity will occur when temperatures are below 8 °C or above 27 °C. Frost 3 days before or after an application may reduce crop tolerance and weed control efficacy.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Liquid Achieve (0.2 L/acre) plus *Turbocharge* adjuvant

Cordon or *Puma Advance* (0.15 to 0.31 L/acre)

In spring wheat (including durum) only:

Signal or *Nufarm Clodinafop* (93 to 117 mL per acre) plus recommended adjuvant

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Enforcer D* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry Interval: DO NOT enter treated fields for 24 hours.

Grazing: DO NOT graze or cut for livestock feed within 30 days of application. Withdraw meat animals from treated feed 3 days before slaughter.

Pre-harvest Interval: DO NOT harvest within 60 days of application.

Re-cropping: Barley, canola, flax, forage grasses, lentil, mustard, oat, pea, rye and wheat can be seeded the following year or fields can be fallowed.

Aerial Application: DO NOT apply by air.

Storage: Store in a ventilated room above freezing. If frozen, allow container to warm and shake well before using.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Spray when winds are under 16 km/hr, but not dead calm.

Tank Cleaning:

The manufacturer provides no recommendations on how to clean equipment used to apply this product. As a petroleum based emulsifiable concentrate, 'Method B' in the general section on tank cleaning on page 15-16 may be the most effective.

Hazard Rating:



Caution - Poison



Caution - Skin and Eye Irritant

For an explanation of the symbols used here see page 11.

Enforcer M

Herbicide Group
4 - fluroxypyr and MCPA
6 - bromoxynil
(Refer to page 38)

Company:

Nufarm Agriculture (PCP#30691)

Formulation:

80 g/L fluroxypyr, 200 g/L bromoxynil and 200 g/L MCPA ester formulated as an emulsifiable concentrate.

Container size: 2 x 10 L, 120 L or 450 L.

Crops and Staging:

Wheat (durum, spring) and barley - 4 leaf to early flag leaf stage.

Winter wheat - in spring from fully tillered to the flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Apply to emerged weeds up to the 6 leaf stage unless otherwise indicated.

Weeds controlled at the 0.25 L per acre (10 L treats 40 acres) rate:

Kochia (up to 5 cm tall)	Wild buckwheat *
Lamb's-quarters	Wild mustard

Weeds controlled at the 0.51 L per acre (10 L treats 20 acres) rate:

Weeds listed above plus:

Cleavers (up to 6 whorls)	Redroot pigweed*
Chickweed	Wild buckwheat

* Suppression only.

Application Information:

Water Volume: Minimum 20 to 40 L per acre. Use the higher volume when there is a heavy crop canopy or weeds are at an advanced stage.

Nozzles and Pressure: Use 40 psi (275 kPa) if applying without drift reduction nozzles. Drift reduction nozzles may require higher pressures for proper performance. Select the nozzle and pressure combination that produces of ASABE coarse droplets while maintaining good coverage of foliage.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Optimum activity is experienced between 12 to 24 °C when weeds are actively growing. Weeds may not be actively growing and as a result reduced activity will occur when temperatures are below 8 °C or above 27 °C. Frost 3 days before or after an application may reduce crop tolerance and weed control efficacy.

Tank Mixes:

Herbicides:

In wheat (durum, spring, winter) and barley:

Liquid Achieve (0.2 L/acre) plus Turbocharge adjuvant

Cordon or Puma Advance (0.15 to 0.31 L/acre)

In spring wheat (including durum) only:

Signal or Nufarm Clodinafop (93 to 117 mL per acre) plus recommended adjuvant

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Enforcer M* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry Interval: DO NOT enter treated fields for 24 hours.

Grazing: DO NOT graze or cut for livestock feed within 30 days of application. Withdraw meat animals from treated feed 3 days before slaughter.

Pre-harvest Interval: DO NOT harvest within 60 days of application.

Re-cropping: Barley, canola, flax, forage grasses, lentil, mustard, oat, pea, rye and wheat can be seeded the following year or fields can be fallowed.

Aerial Application: DO NOT apply by air.

Storage: Store in a ventilated room above freezing. If frozen, allow container to warm and shake well before using.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Spray when winds are under 16 km/hr, but not dead calm.

Tank Cleaning:

The manufacturer provides no recommendations on how to clean equipment used to apply this product. As a petroleum based emulsifiable concentrate, 'Method B' in the general section on tank cleaning on page 15-16 may be the most effective.

Hazard Rating:



Warning – Poison



Caution – Skin and Eye Irritant

Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Eptam 8-E

Herbicide Group

8 - EPTC

(Refer to page 38)

Company:

Gowan Canada (PCP# 11284)

Formulation:

800 g/L of EPTC formulated as an emulsifiable concentrate.
Container size - 10 L, 1000 L

Caution:

The level of weed control may be reduced where *Eptam 8-E* is used on soils that have been treated with *Eptam 8-E* the previous growing season. It is expected that the reduction in control will be greater where *Eptam 8-E* have been used repeatedly for 2 or more years.

Crops, Rates and Staging:

Eptam 8-E is applied as a preplant incorporated treatment prior to seeding the following crops:

CROP	RATE (L/ACRE)	ACRES TREATED PER 10 L CONTAINER
Dry and snap beans	1.72 to 2.23	5.8 to 4.5
Alfalfa, Bird's-foot trefoil Cicer milk-vetch** Sweet clover** Sunflowers*	1.72	5.8
Potatoes	1.72 to 3.44	5.8 to 2.9
Flax*	1.42 to 1.72	7.0 to 5.8

* May also be applied in late fall prior to freeze-up

** Seed production only

NOTE: The use of *Eptam 8-E* on flax is not recommended in Saskatchewan because of the risk of crop injury.

Where a rate range appears, use the lower rate on light tex-

tured soils and the higher rate on heavy textured soils. DO NOT use on soils with less than 3 percent organic matter or more than 15 percent organic matter.

Weeds and Staging:

Must be applied prior to the emergence of the following weeds. Emerged weeds will not be controlled.

Barnyard grass	Pigweed (prostrate, redroot, tumble)*
Chickweed*	Purslane*
Corn spurry*	Quackgrass (suppression)**
Green foxtail	Volunteer cereals (wheat, barley, oats)
Hairy nightshade*	Wild oat
Henbit *	Yellow foxtail
Lamb's-quarters*	Yellow nutsedge**
Nettleleaf goosefoot*	

* Will be controlled only if treatment is made when conditions are favourable for germination and growth.

** Roots of perennial weeds must be thoroughly chopped prior to application.

Application Information:

Carrier Volume: Minimum of 40 L per acre of water or liquid fertilizer (see label for liquid fertilizer compatibility).

Pressure: 30 to 40 psi (200 to 275 kPa).

Equipment and Nozzles: Since *Eptam 8-E* is highly volatile, the product must be incorporated immediately. This is best accomplished by mounting spraying equipment directly onto the incorporation equipment (tandem disks, field cultivators on light soil).

May also be applied to cleanly cultivated soil for potatoes, by metering into the irrigation water to achieve the recommended rate per acre ("herbigation" or "chemigation"). See label for detailed instructions.

Incorporation: All growth and stubble should be thoroughly worked into the soil before treatment. Apply to a dry soil surface. Incorporate immediately after application preferably during the spraying operation as *Eptam 8-E* is volatile. Set disc and cultivator implements to cut to a depth of 4 to 6 inches (10 to 15 cm). A second operation at a right angle to the first is required. The disc or cultivator must be followed with a harrow or other levelling device that extends beyond the width of the implement. Speeds in excess of 5 mph (8 km/h) will result in excessive pulverization and trash destruction leaving the field susceptible to erosion. The maximum recommended tillage depth is 4 inches (10 cm).

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop injury can occur if stressful environmental conditions (cold, wet soils, drought or excessive heat) prevail after

seeding. To minimize crop injury, delay seeding 10 days if these conditions prevail at the time of application, or select an alternative product. Very cold or dry soil conditions during weed emergence will reduce control.

Tank Mixes:

Herbicides:

Dry beans (white and red kidney only): Liquid formulations of *Treflan* and *Rival*.

Insecticides: DO NOT tank mix with insecticides.

Fungicides: None registered.

Fertilizers: May be mixed with liquid fertilizer.

Compatibility test should be conducted according to instructions on the herbicide label.

Dry bulk fertilizers, except nitrate fertilizers, may be impregnated or coated with *Eptam 8-E*. The impregnated fertilizer should be spread uniformly onto the field using a double overlap pattern immediately after impregnation. The impregnated fertilizer must be applied to the field when the soil surface is dry to at least 1/2 inch (1.5 cm) depth. The first incorporation must be done immediately after application.

Note: The above mixes are those listed on the *Eptam 8-E* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No effect once incorporated. DO NOT apply prior to pre-irrigation.

Re-entry: DO NOT enter treated fields for 24 hours.

Grazing: DO NOT graze or feed treated crops to livestock in the year of application.

Re-cropping: Will not injure crops the year after spring application.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones: DO NOT apply within 15 m of fish bearing waters or wildlife habitat.

Soil Type: DO NOT use on soils with less than 3 percent organic matter as crop injury will result.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution - Poison

For an explanation of the symbols used here see page 11.

Equinox*

Herbicide Group
1 - tepraloxymid
(Refer to page 38)

Company:

BASF Canada (PCP# 27603)

Formulation:

200 g/L tepraloxymid formulated as an emulsifiable concentrate

Container size -

Equinox: 3.24 L jug;

Merge: 8.1 L

* NOTE: This product is no longer manufactured, but some supplies may still remain in the distribution system. This product may be removed from future editions of this publication.

Crops, Rates and Staging:

Rates are dictated by weed type and stage below.

CROP	RATE (mL/acre)	MAXIMUM STAGE
Canola ^{CL}	54 to 81	Prior to bud and flower formation
Chickpeas	54 to 101	9 node
CLEARFIELD Oilseed quality mustard (<i>Brassica juncea</i>)	54 to 81	Prior to bud and flower formation
Dry beans	54 to 101	7 trifoliate leaf
Flax, Solin (low linolenic acid flax)	54 to 101	35 cm
Field pea, Lentil ^{CL}	54 to 101	9 leaf
Mustard (brown, oriental, yellow),	54 to 81	6 leaf
Sunflower	54 to 81	10 leaf

^{CL} Including CLEARFIELD varieties.

Weed, Rates and Staging:

Apply at the 1 to 6 leaf stage of the annual weeds listed below unless otherwise indicated.

Optimum yield response occurs when weeds are controlled early.

Equinox will not control weeds emerging after application.

WEEDS AND STAGES	RATE	
	mL/ACRE	ACRES PER 3.24 L JUG
Green foxtail, Wild oat, volunteer cereals (barley, oat, and wheat including CLEARFIELD varieties)	54 to 81*†	60 to 40*†
Quack grass control (3 to 6-leaf stage**)	101	32

Equinox must always be used with *Merge* adjuvant at a rate of 0.5 L of *Merge* per 100 L of total spray solution.

* Use the 40 acre/case rates when weed densities exceed 45-weeds per square meter.

** For optimum control of quackgrass apply at the 2 to 4 leaf stage prior to tillering (6 to 15 cm in height).

† Use higher rate in range when weed densities and overlapping are high, when staging is late, or when weeds are under stress and not growing as actively due to moisture stress or temperature stress.

Application Information:

Water Volume: 20 to 40 L per acre.

Ground: 20 to 40 L per acre.

Aerial: 10 to 20 L per acre.

Nozzles and Pressure: Use a minimum pressure of 35 psi (240 kPa) when using conventional nozzles. For high weed pressures or if crop or weed growth is dense use a pressure 40 to 62 psi (275 to 425 kPa). Low drift nozzles may require higher pressures for proper performance. Use 80° or 110° flat fan nozzles designed to deliver thorough, even coverage of ASABE medium droplets tilted forward at an angle of 45° are recommended for optimum coverage. The use of flood jet or hollow cone nozzles is not recommended. Consult with herbicide manufacturer regarding the suitability of low drift nozzles for use with this product.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Most effective control is achieved when grasses are actively growing. Weeds stressed by drought, flooding, hot or prolonged cool temperatures (<15°C) and poor fertility are more difficult to control. Use the higher of the recommended rates for grasses under stress. To avoid reduced control do not apply to grasses stressed because of lack of moisture, flooding, hail, herbicide damage, mechanical injury and widely fluctuating temperatures. To maximize control apply *Equinox* in warm weather as control may be reduced if temperatures are below 15°C.

Tank Mixes:

Tank mixes are not to be applied by air.

Herbicides:

In Flax and Solin (low linolenic acid flax):

Buctril M (0.4 L/acre) (Including Solin varieties)

In Field peas, CLEARFIELD canola, and CLEARFIELD lentil only:

Odyssey (17 grams/acre)

In LIBERTY LINK canola only:

Liberty 150SN (1.08 to 1.35 L/acre)

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Allow 4 days between the application of *Equinox* and any chemical not recommended as a tank mix.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-Entry: DO NOT enter treated fields for 12 hours.

Grazing: Field peas may be used for grazing following the 60-day Preharvest interval. DO NOT graze other treated crops or cut for feed prior to crop maturity.

Preharvest Interval: Lentil, field peas, canola & flax – 60 days.

Recropping: Leave at least 14 days between application and replanting of cereal or grass crops. Leave at least 40 days between application and the seeding of other crops. The company recommends that treated area be cultivated to a depth of 10 cm at least 7 days prior to seeding.

Aerial Application: May be applied by air.

Storage: Store in a cool, dry, well-ventilated area without floor drain away from food, feed, seed or fertilizers.

Buffer Zones: Leave a buffer of 1 meter from the down wind edge of the boom and sensitive terrestrial habitat by ground and 25 m by airplane or helicopter. Corn is particularly sensitive.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution – Poison



Warning – Eye and Skin Irritant

For an explanation of the symbols used here see page 11.

Escort

Herbicide Group
2 - metsulfuron
(Refer to page 38)

Company:

E. I. duPont Canada (PCP#23005)

Formulation:

60% metsulfuron methyl formulated as a water dispersible granule.

Container size - 0.25 kg.

Crops and Staging:

Pasture, rough turf, and rangeland - No stage restrictions.

Weeds, Rates and Staging:

For seedling weeds apply to young plants up to 4 inches (10 cm) tall or wide. For established non-woody plants (biennial or perennial) apply up to the early bud stage. For western snowberry, wild rose and other woody species, apply between mid-June and mid-August after the brush has leafed out, but before the leaves begin to turn their fall colours.

RATE	WEEDS CONTROLLED
8 g/acre	Canada thistle* Common tansy Dandelion* Scentless chamomile Russian thistle Sweet clover Sow-thistle*
10 g/acre	<i>Above weeds plus:</i> Western snowberry
12 g/acre	<i>Above weeds plus:</i> Wild rose Dandelion
40 g/acre**	Balsam poplar Willow
60 g/acre**	Cherry Trembling aspen

At all rates add *Agral 90*, *Agsurf II*, or *Citovett* at 0.2 L per 100 L of spray solution.

* Suppression only.

** Rangeland only. See label for detailed application instructions.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information

Water Volume: 40 to 91 L per acre for weedy growth and up to 809 L per acre applied to the point of run-off for woody species. See the label for details.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply during periods of intense rainfall or to soil saturated with water. Warm, moist conditions following treatment enhance the activity of *Escort*, while cold, dry conditions may reduce or delay activity. Brush hardened off by cold weather and drought stress may not be controlled.

Tank Mixes:

Herbicides:

2,4-D amine or ester at 0.79 L/acre (500 g/L formulations) plus surfactant.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT re-enter treated fields for 12 hours.

Grazing: May be grazed by cattle on the day of treatment.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Rate (g/acre)	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat**
	Less than 1 m	Greater than 1 m	
8 to 12	1	1	10
40	2	1	35
60	3	1	45

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

** Terrestrial buffers are not required for transport and utility rights of way.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

Escort can cause severe injury to sensitive crops at very low concentrations. Use 'Method A' on pages 15 to 16 to clean sprayers immediately after using *Escort*.

Hazard Rating:

▽ Caution – Eye Irritant

For an explanation of the symbols used here see page 11.

Everest GBX

Herbicide Group
2 - flucarbazone
4 - fluroxypyr
(Refer to page 38)

Company:

Arysta LifeScience Canada

Formulation:

Everest 2.0 (PCP# 30342): 397 g/L flucarbazone formulated as a suspension concentrate.

Container size - 1.937 L container

GBX (PCP# 29958): 333 g a.e./L of fluroxypyr formulated as an emulsifiable concentrate.

Container size - 5 L

Crops and Staging:

Spring wheat (including durum) with 2 leaf to a maximum of 4 main stem leaves plus 2 tillers (6 total leaves).

Tank mix options are listed in the tank mix section.

Note: Some of the tank mix partners may have more limiting staging than Everest. When tank mixing use the most restrictive application stage or injury may result.

Weeds, Rates and Staging:

RATE (Acres per case)	WEEDS
100	Weeds controlled by Everest 2.0 at 19.4 mL per acre plus: Cleavers from the 1 to 4 whorl stage Kochia (2 to 8 leaf stage) (2,4-D ester mix only) Wild buckwheat (1 to 4 leaf stage)
80	Weeds controlled by Everest 2.0 at 24.3 mL per acre plus: Kochia (2 to 8 leaf stage) Volunteer flax (1 to 12 cm) Stork's-bill (Suppression only) (1 to 8 leaf)

As well as any of the weeds controlled by the tank mix partner.

This product **MUST ALSO BE MIXED** with one of either 2,4-D Ester, MCPA Ester or Curtail M.

Add non-ionic surfactant (such as Agral 90, Aqsurf) at 0.25 L per 100 L of spray solution.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Tank Mixes:

Herbicides:

Note: All mixes must be applied with a registered surfactant unless otherwise indicated. Only one registered surfactant is required.

In spring wheat (including durum):

2,4-D Amine or Ester at recommended rates up to 0.34 L/acre (500 g/L formulation)

In spring wheat (NOT including durum):

2,4-D Amine or Ester at recommended rates up to 0.45 L/acre (500 g/L formulation)

Curtail M (0.6 to 0.8 L/acre)

MCPA Amine or Ester at recommended rates up to 0.45 L/acre (500 g/L formulation)

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the Everest GBX label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Grazing: DO NOT graze treated fields. Mature grain or straw may be fed to livestock.

Preharvest: Leave at least 80 days from application to harvest.

Re-cropping Interval: Follow the chart below for Everest GBX components only. If mixing with Curtail M as well

check for any additional recropping restrictions on the Curtail M page:

Soil Zones and Rotational Crops		
Grey-Wooded	Black and dark brown	Brown
Spring Wheat Barley Canola (all varieties) Field Pea*	Wheat (Spring & durum) Barley Canola (all varieties) Field Pea* Flax	Spring Wheat

* Field pea may be grown the year following Everest GBX application in fields where precipitation has been equal to or above the 10 year average during the growing season, and where organic matter content is above 4%, and pH is below 7.5. The company suggests a minimum of 100 mm (4 inches) of rain is needed in the 60 days following application for adequate breakdown to take place.

NOTE: Other rotational crops may also be affected if rainfall is less than the 10 year average for the area. Soils in the grey wooded, black and dark brown soil zones with a combination of low organic matter (less than 2%), light textured soils or high pH (greater than 7.5) (i.e. eroded knolls, sandy soils) may result in delayed growth and development in rotational crops.

Aerial Application: DO NOT apply by air.

Storage: Store in closed original container in a cool, dry area away from fertilizers, food or feed. DO NOT freeze. If frozen bring Everest GBX component to room temperature and agitate (shake) before use. The GBX component is combustible – DO NOT store near heat or open flame.

Buffer Zones: Leave at least 20 m from the downwind edge of the spray swath to sensitive upland plants like shelterbelts and woodlots and at least 35 m to water sources or wetland habitats. Avoid drift onto sensitive crops like canola and tame oat. DO NOT mix or load within 10 m of water sources or wetland habitats.

Tank Cleaning:


Everest GBX residues in the spray tank can cause severe injury to sensitive crops at very low concentrations. Sprayers should be cleaned out immediately before using another product. Follow 'Method A' on page 15 to 16 to clean sprayers immediately after Everest GBX application. Since the GBX component is petroleum based, it is recommended that detergent as indicated in 'Method B' also be included in the cleaning process.

Hazard Rating:


Everest Component:


 Warning – Contains the Allergen Milk

GBX component:

 Danger – Poison

 Warning – Eye and Skin Irritant

 Caution – Skin Irritant

 Caution – Flammable

For an explanation of the symbols used here see page 11.

Express Pro

Herbicide Group
2 - tribenuron & metsulfuron
 (Refer to page 38)

Company:

E. I. duPont Canada (PCP# 29212)

Formulation:

42.9% tribenuron methyl, and 8.6 % metsulfuron methyl formulated as a water soluble granule.

Container size - 560 g container.

Express Pro is purchased alone but must be used in a mix with glyphosate before use.

Crops and Staging:

Allow at least one day (24 hours) prior to seeding barley, winter wheat, and wheat (durum, spring, winter) and summer-fallow.

Allow 10 days between summer-fallow treatment and tillage.

DO NOT use *Express Pro* on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits

Weeds, Rates and Staging:

Express Pro at 7 g per acre (one container treats 80 acres) plus glyphosate at a rate equivalent to 180 g ae per acre (equivalent of 0.5 L per acre of a 360 g per L formulation):

Weeds controlled by glyphosate products at these rates plus up to 3 inches (8 cm) unless otherwise indicated:

Canada thistle rosettes**	Scentless chamomile
Cleavers†	White cockle (rosettes)
Cow cockle*	Volunteer canola
Dandelion ***†	(including glyphosate
Narrow-leaved hawk's-beard†	tolerant varieties) ***†
Night-flowering catchfly**	

† Residual control

* Up to the 3 leaf stage

** Suppression only

*** Up to 6 inches

Express Pro may degrade if left in the sprayer for an extended period. Apply within 24 hours of mixing.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 22 to 45 L per acre.

Nozzles and Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets. Low drift nozzles may require higher pressures for proper performance.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Registered crops seeded following *Express Pro* application become stressed by drought, low fertility, saline soils, waterlogged soils (soils at or near field capacity), disease or insect damage may be injured. This injury may be worse on light or low organic matter soils. Weeds hardened off by environmental stress such as those above may not be adequately controlled.

Tank Mixes:

Herbicides: Must be mixed with glyphosate.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Express Pro* label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check product labels for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: Wait 12 hours before re-entering treated fields.

Re-cropping: Barley and wheat (spring, winter and durum) may be seeded 24 hours after application. Canola, flax and peas may be planted 10 months following application.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat*
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	4

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Express Pro can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray this product should be flushed out immediately after use.

1. Drain tank and flush tank, boom and hoses with clean water.
2. Visually inspect tank to ensure removal of all visible herbicide residues. If necessary, repeat Step 1.
3. Fill the tank with clean water, and then add 1 litre household ammonia (containing minimum of 3% ammonia) per 100 litres of water, or equivalent amount of a sprayer tank cleaner containing ammonia,

4. Flush solution through boom and hoses, and then add more water to completely fill tank. Allow to sit for 15 minutes with agitation.

5. Drain the tank.

6. Remove the nozzles, screens, and boom end caps and clean separately in a bucket containing ammonia cleaning agent and water.

7. Thoroughly rinse the tank with clean water for a minimum of 5 minutes, flushing water through the hoses and boom.

Do not clean equipment upslope of water bodies or ditches, near cropland or shelterbelts. Clean your sprayer away from areas where family members or others are likely to frequent or walk. See product label for more detailed cleaning instructions. For additional information on tank cleaning see page 15.

Hazard Rating:

Caution – Poison



Warning – Eye Irritant



Potential Skin Sensitizer.

Contains the allergens sulphites and milk.

For an explanation of the symbols used here see page 11.

Fenoxaprop**Herbicide Group**

1 - fenoxaprop

(Refer to page 38)

Company:

Bayer CropScience (*Puma Advance*)

Farmers of North America (*HellCat*)

IPCO (*Vigil WB*)

MANA Canada (*Bengal WB*)

Nufarm Agriculture (*Cordon*)

Loveland Products (formerly Viterra) (*WildCat*)

Cheminova (*Cougar*)

Formulation:

Bengal WB (PCP#30843), *Cordon* (PCP#29494), *Cougar* (PCP#30473), *HellCat* (PCP#30055), *Vigil WB* (PCP#30844), *WildCat* (PCP#29151):

120 g/L fenoxaprop-p-ethyl formulated as an emulsifiable concentrate.

Container size* - 6.2 L, 12.4L, 18.6 L, 99.3L, 312L.

Puma Advance (PCP#29615): 90 g/L fenoxaprop-p-ethyl. Container size - 8.25L, 123.75L, 412.5L.

* Check with individual suppliers for the container sizes they have available.

Crops and Staging:

Application beyond the maximum rates provided below may result in crop injury.

CROP	STAGE
Spring wheat (including durum), Barley [†]	1 to 6 leaves on the main stem plus 3 tillers
Barley ^{††}	1 to 5 leaves on the main stem plus 2 tillers
Perennial ryegrass seedlings (seed production only)*	2 to 4 leaves
Meadow brome grass (seedling or established) (forage or seed production)**	

[†] *Puma Advance* only. Late application of other products could result in injury to barley

^{††} *Bengal WB, Cordon, HellCat, Vigil WB, or WildCat* only. Apply to barley only when tank mixed with a registered broadleaf product. NOTE: Application of other fenoxaprop products to barley can result in crop injury.

* Perennial ryegrass with *Bengal WB, Cordon, Cougar, Vigil WB* or *WildCat* by ground only.

** Meadow brome grass with or *Puma Advance* by ground only.

NOTE: Since the uses on forage grasses were registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Those who apply this use do so at their own risk.

Durum wheat, forage grasses and barley may experience some initial, temporary stunting and yellowing that rarely results in yield loss. Injury is more likely under stress conditions (see "Effect of Growing Conditions" section).

Treatment at the 3 to 4 leaf stage of cereal crops and weeds will maximize crop tolerance and weed control. Temporary crop injury such as shortening or discolouration may be observed after application. Such injury is more likely to occur in barley and also when *fenoxaprop* is applied outside recommended stages.

Application Information:

Water Volume:

Ground application: 23 to 45 L per acre. Use higher water volumes for dense canopies.

Aerial application: A minimum of 14 L per acre.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional 80° or 110° flat fan nozzles. Angle nozzles forward at 45° to improve contact with vertical leaf surfaces.

Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver proper coverage with *ASABE medium* droplets or larger.

DO NOT use flood jet nozzles, controlled droplet application equipment or Spraying equipment.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply *fenoxaprop* 2 to 3 days prior to, or following, temperatures of 3°C or lower as crop injury may occur. Under stressful conditions (hot/dry, water logging, disease or insect damage) or heavy crop canopy, early application will improve weed control.

DO NOT apply by air when both the temperature is greater than 25°C and the relative humidity is less than 30%.

Weeds, Rates and Staging:

Apply from the 1 to 6 leaf stage up to emergence of 3rd tiller of the weeds below. Apply at the 3 to 4 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering. DO NOT apply *fenoxaprop* or products containing *fenoxaprop* to a crop more than once per year.

WEEDS	RATE (mL/ACRE)		RATE (ACRES PER PACKAGE [†])
	<i>Puma Advance</i>	120 g/L forms	
Green foxtail only	206	156	80
Low wild oat infestations*	360	271	46
Moderate-heavy wild oat infestations, barnyard grass, green and yellow foxtail	413	312	40

[†] Based on 12.4 L for 120g/L formulations and 16.5 L for *Puma Advance*.

* Low wild oat rate for use on WHEAT AND DURUM ONLY, and when applied alone and NOT in a tank-mix. NOT for use with perennial ryegrass or meadow brome grass.

Tank Mixes:

Herbicides:

DO NOT apply *Bengal WB*, *Cordon*, *Vigil WB* or *WildCat* in barley without a broadleaf herbicide mix. ALWAYS tank mix with a registered broadleaf herbicide.

2,4-D Ester LV 600 (0.28 L/acre), LV 700 (0.24 L/acre)[†]
Ally (2 to 3 g/acre)[†]
Attain XC (label rates)^{††}
 Bromoxynil/2,4-D ester (0.4 L/acre)^Δ
 Bromoxynil/MCPA ester (0.4 L/acre)^Δ
Curtail M (0.6 to 0.8 L/acre)
 Dichlorprop/2,4-D (label rate)^Δ
Estoprop XT (label rates)^{†††}
DyVel (0.5 L/acre)^{**}
DyVel DSp (0.45 L/acre)^{*** Δ Δ}
Infinity (0.33 L/acre)^{ΔΔΔ}
*Lontrel 360**** (0.17 L/acre)
*Lontrel 360**** (0.17 L/acre) + MCPA 500 Ester (0.34 L/acre)^{*}
*Lontrel 360**** (0.112 L/acre) + MCPA 500 Ester (0.34 to 0.45 L/acre)^{*}
 MCPA Amine or Ester (0.34 L/acre) (500 g ai/L formulation)
 Mecoprop-p^{***} (2.2 to 2.8 L/acre)^{ΔΔ}
Prestige XC (label rates)^{††}
Refine SG (12 g/acre)^{††}
Refine SG (12 g/acre) + MCPA (rates above)^{††}
Refine SG (4 g/acre) + Buctril M (0.4 L/acre)^{††}
Spectrum (20 acres/case rate)^{***}
Triton C (label rate)^{ΔΔΔ}
Trophy (20 acres per case)^{***}

[†] All products except *HellCat*.

^{††} *Puma Advance* and *Cougar* *** only.

^{†††} *Cordon* and *Cougar* only.

^Δ Manufacturers may only support mixes with specific brands. Contact the manufacturers for more information.

^{ΔΔ} *Puma Advance*, *Cordon* and *Cougar* only.

^{ΔΔΔ} *Puma Advance* only.

^{*} Use only at the high rate of fenoxaprop.

^{**} Use only at the green foxtail rate of fenoxaprop.

^{***} Use in wheat only

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *fenoxaprop* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Fenoxaprop manufacturers may also support mixes with pesticides that are not on the *fenoxaprop* labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Leave an interval of 7 days prior to application or 4 days after application of *fenoxaprop*, when applying any pesticide that is not registered as a tank mix.

Adding ingredients in the correct order is critical for opti-

mum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-Entry: DO NOT re-enter treated fields for 12 hours.

Grazing: DO NOT graze or cut cereal crops or meadow brome grass for hay, within 25 days of application. DO NOT graze or cut perennial ryegrass crop for hay within 65 days of application.

Preharvest Interval: DO NOT harvest within 65 days of application.

Re-cropping: No restrictions in the year after application. Only one application may be made per year.

Aerial Application: May be applied by air.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground*	3	10
Aerial	3	33

See page 29 for an explanation of the different habitats.

* These distances can be reduced by 30% using cones on individual nozzles and by 70% using a full shield (shroud, curtain) that extends to the crop canopy.

[†] Distance is measured from the downwind edge of the boom to sensitive areas.


Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15.

Hazard Rating:

 Caution - Poison.


Cougar, *HellCat* and *WildCat*:

 Warning - Eye and Skin Irritant.

Bengal:

 Danger - Eye and Skin Irritant

All:

 Warning - Contains the allergen soy

For an explanation of the symbols used here see page 11.

Flexstar GT*

*(For use only in the Red River Valley of Manitoba)

Company:

Syngenta Canada (PCP# 30412)

Formulation:

67 g/L fomesafen and 271 g/L glyphosate formulated as a solution.

Container size: 2x10 L and 450 L

Crops and Staging:

Flexstar GT may be applied as a pre-seed burn down or as pre-emergent to the crop of soybeans or as early post-emergent on 1 to 2 trifoliate leaf stage of glyphosate tolerant soybeans only.

For use in the Red River Valley of Manitoba only.

Weeds and Staging:

Control of the following weeds at the cotyledon to 3 or 4 true leaf stage.

Grasses

Barnyard grass	Proso millet
Bromegrass (smooth)	Quackgrass
Cattail (common)	Rye, tame
Crabgrass (large, smooth)	Volunteer barley
Downy brome	Volunteer corn
Foxtail barley	Volunteer wheat
Foxtail (green, yellow)	Wild oats
Persian dandelion	Yellow nutsedge

Broadleaf Weeds

Absinthe
Canada thistle
Chickweed, common
Cleavers
Clover, white
Cocklebur
Cow cockle
Curled dock
Dandelion, common
Field bindweed
Fleabane (Canada)
Flixweed
Hemp-nettle
Horsetail
Knotweed (Japanese, prostrate)
Kochia
Lamb's-quarters
Milkweed (common)
Narrow-leaved hawk's-beard
Night-flowering catchfly

Herbicide Group

9 – glyphosate

14 – fomesafen

(Refer to page 38)

Nightshade, Eastern black
Volunteer canola (NOT glyphosate tolerant varieties)
Pigweed (redroot, smooth)
Prickly lettuce
Ragweed (common)
Round-leaved mallow
Russian thistle
Shepherd's-purse
Smartweed (green, Lady's-thumb)
Sow-thistle (annual, perennial)
Stinkweed
Stork's-bill
Volunteer alfalfa
Volunteer flax
Wild buckwheat
Wild mustard
Wild tomato

Rates:

840 mL per acre.

DO NOT apply *FlexStar GT* or other products containing the ingredient fomesafen more than once in two consecutive years.

Add *Turbocharge* adjuvant at 0.25 L per 100 L spray solution only when weeds are under stress conditions and for larger weeds.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum of 60 to 80 L clean, clear water per acre. Higher spray volume is required for dense weed stands.

Pressure: 210 kPa (35 psi). Increase pressure to 420 kPa (70 psi) for fields with heavy weed densities or with weeds at the upper limit of their recommended stage.

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Moisture is necessary to activate the herbicide for residual weed control. Dry weather following application of the herbicide may reduce effectiveness. Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect activity.

Tank Mixes:

Herbicides:

Touchdown Total (0.28 to 1.0 L per acre)

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Pre-harvest Interval: Leave at least 90 days from application to harvest.

Grazing: DO NOT graze treated crop or cut for hay. There is insufficient data to support such use.

Re-cropping: Winter wheat may be sown 4 months after application. Spring wheat, dry beans, soybeans and field corn may be grown the year following an application. DO

NOT apply *Flexstar GT* to any field more often than once every 2 years. These re-cropping restrictions refer only to the Red River Valley of Manitoba. Use outside this region is not registered as re-cropping options have not been determined.

Aerial Application: DO NOT apply by air.

Storage: Store above -10°C, in a dry place in original container, away from food or feed.

Buffer Zones: Leave a buffer zone of at least 15 m between the last spray swath and the edge of sensitive terrestrial areas such as shelterbelts, hedgerows and shrublands as well as aquatic areas such as ponds, streams, rivers, prairie potholes and sloughs. Do not apply when winds are greater than 15 km/hr.

Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Warning - Eye Irritant.

For an explanation of the symbols used here see page 11.

Florasulam + glyphosate

Herbicide Group

2 - florasulam

9 - glyphosate

(Refer to page 38)

Company:

Dow AgroSciences (*PrePass XC*)

MANA Canada (*Priority*)

Formulation:

The *PrePass XC* package contains 2 components:

PrePass XC A (PCP# 29651): 50 g/L florasulam formulated as a suspension concentrate.

PrePass XC B (PCP# 29652): 480 g/L glyphosate DMA salt formulated as a solution.

Container sizes:

PrePass XC A - 1.6 L (40 acre); 4 x 12 L (1200 acre).

PrePass XC B - 2 x 7.5 L (40 acre); 4 x 112.5L or 450 L (1200 acre).

-or-

Priority (PCP# 30831): 50 g/L florasulam formulated as a suspension concentrate.

Container size - 2 x 6.4L (320 acre)

Priority DOES NOT come packaged with glyphosate.

Glyphosate in a DMA or IPA salt form must be purchased separately.

Crops and Staging:

Florasulam + glyphosate can be applied either in the fall or in the spring prior to seeding spring wheat (including durum), barley or oats or as an initial treatment in summerfallow.

Weeds and Staging:

PrePass or *Priority* plus glyphosate IPA or DMA salt will control the following weeds:

Grasses at the 2 to 4 leaf stage:

Downy brome

Green foxtail

Persian darnel

Volunteer barley

Volunteer wheat

Wild oat

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Canada fleabane**	Redroot pigweed
Common chickweed	Russian thistle
Cleavers	Scentless chamomile
Dandelion (up to 30 cm across)	Shepherd's-purse
Flixweed	Stinkweed
Hemp-nettle	Smartweed (including Lady's-thumb)
Kochia	Volunteer canola (all varieties)
Lamb's-quarters	Volunteer flax
Narrow-leaved hawk's-beard**	Wild buckwheat*
Ragweed (common)**	Wild mustard

Broadleaf weeds suppressed:

Annual sow-thistle	Perennial sow-thistle***
--------------------	--------------------------

* Up to 5 leaf stage.

** Up to 8 cm in height.

*** Earlier applications provide better results.

Rate:**PrePass XC A:** 40 mL per acre.

-plus-

PrePass XC B: 375 mL per acre.

-or-

Priority: 40 mL/acre.

-plus-

Glyphosate DMA or IPA salt (not supplied): 180 gae/acre (see glyphosate page for product rates).

(See "Formulations:" section for package rates.)

Note: Maximum one application of this product or other products containing florasulam in back to back years.

Refer to the product labels for complete mixing instructions for these products. A general guide to mixing can be found on page 14.

Application Information:**Water Volume:** 20 to 40 L per acre.**Nozzles and Pressure:** Use 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets.**How it Works:**

Refer to Table 2 on page 40.

Effects of Growing Conditions:**PrePass XC A or Priority:** Warm, moist growing conditions promote active weed growth and enhance activity. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur.**PrePass XC B or glyphosate:** Best results are achieved when temperatures are near 20°C and when weeds are actively growing. Frost that kills more than 40% of above ground tissue will reduce control. Heavy dust layer on leaves will also reduce control.**Tank Mixes:****PrePass:** None registered.**Priority:** must be mixed with glyphosate IPA or DMA salt** at 180 gae/L.**Dow AgroSciences supports the following mixes that are not on the Pre-Pass label. Apply mixes according to the most restrictive use limitations for either product:****Herbicides:** 2,4-D ester, MCPA ester, Vantage Plus Max II**

** NOTE: Mixing with other salts of glyphosate may have a negative reaction with the florasulam component.

Restrictions:**Rainfall:** Heavy rainfall immediately after application may wash the chemical off the foliage. DO NOT apply if rainfall is forecast for the time of application. Contact manufacturer for more information. DO NOT apply excessive irrigation following application as product may leach.**Re-entry:** DO NOT enter treated fields for at least 12 hours.**Grazing:** DO NOT graze treated crop or cut for feed within 7 days of application.**Re-cropping:** Spring wheat (including durum), barley and oat, may be seeded, or the field may be fallowed, after applications made in the spring prior to seeding, or the previous year after August 1. Barley, canola, field peas, oat and wheat, may be grown following applications made prior to August 1 of the previous season.**Aerial Application:** DO NOT apply by air.**Storage:** Store in dry, heated area. DO NOT freeze.**Buffer Zones:**

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	5 ^{††}	5 ^{††}	30 ^{††}

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

†† Buffers apply to PrePass XC A and Priority only. Refer to buffers on PrePass XC B (= Maverick or Vantage Plus Max II) or the specific glyphosate products (see glyphosate page) mixed with Priority and use the most restrictive buffer of the two products.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Caution – Poison



Caution – Irritant, may cause eye irritation.

For an explanation of the symbols used here see page 11.

Florasulam + MCPA Ester

Herbicide Group

2 - florasulam

4 - MCPA

(Refer to page 38)

Company:

Dow AgroSciences (*Frontline XL*)

MANA Canada (*Topline*)

Formulation:

Frontline XL (PCP#28804): 4 g/L florasulam and 280 g/L MCPA ester formulated as an emulsifiable concentrate.
Container size - 2 x 10 L.

The *Topline* package contains two components:

Florasulam SC (PCP#30814): 50 g/L florasulam formulated as a suspension concentrate.

Checkmate MCPA Ester 600 (PCP#27804): 600 g/L MCPA Ester formulated as an emulsifiable concentrate.
Container size - *Florasulam SC*: 1.6 L; *MCPA Ester*: 9.33

Crops and Staging:

Spring wheat (including durum), barley and oats in the 2 to 6 leaf stage.

Frontline XL only:

Seedling and established timothy for forage and seed production.*

Seedlings: from the 2 leaf fully expanded stage up to the flag leaf stage.

Established: no stage restrictions.

* NOTE - Since applications to timothy has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to timothy is at the risk of the user.

When tank-mixing, always check the tankmix partner recommendations for additional staging restrictions.

Weeds and Staging:

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Ball mustard	Redroot pigweed [◇]
Burdock**	Russian pigweed**
Chickweed	Shepherd's-purse
Cleavers	Smartweed
Cow cockle††	Stinkweed
Flixweed**	Sunflower (annual)**
Hemp-nettle [◇]	Volunteer canola*
Lamb's-quarters	Wild mustard
Prickly lettuce**	Wild buckwheat
Ragweed (common)	

Broadleaf weeds suppressed:

Canada thistle [◇]	Stork's-bill
Dandelion*** [◇]	Sow-thistle (annual)
Plantain [†]	Sow-thistle (perennial) [†]

* including all herbicide-tolerant canola varieties

** up to the 4 leaf stage of development

*** seedlings and overwintered rosettes less than 15 cm (6 inches)

[◇] for improved control of this weed add an additional 47.5 ml per acre of MCPA LV600.

[†] top growth control

†† *Frontline XL* only.

Rate:

Frontline XL: 0.5 L per acre

-or-

Topline:

Florasulam SC: 40 mL per acre

-plus-

Checkmate MCPA 600 Ester: 0.23 L per acre

One case of either product treats 40 acres.

Note: Maximum one application of these products or other products containing florasulam in back to back years.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: A minimum of 40 L per acre.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) with conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity of *florasulam + MCPA ester*. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. Under conditions of low crop and high weed density, control may be reduced. Extreme growing conditions such as drought or near freezing temperature prior to, at or following time of application may increase the risk of crop injury at all stages of growth.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley only:

Assert (0.65 L/acre) plus *Acidulate*.

In spring wheat (including durum) only:

Ladder (93 or 117 mL/acre) plus adjuvant.

Simplicity at 0.2 L/acre (no adjuvant required).†

In spring wheat (NOT including durum) and barley only:

*Axial 100EC**† (0.24 L/acre) plus *Adigor* adjuvant.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

* This tank mix may result in reduced levels of green foxtail control.

† *Frontline XL* only.

Note: The above mixes are those listed on the *florasulam + MCPA ester* labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Dow AgroSciences also supports the following mixes that are not on the *Frontline XL* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Avenge*, *clodinafop* (*Signal* and *NextStep*)

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information. DO NOT apply excessive irrigation following application as product has the potential to leach.

Re-Entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT allow lactating dairy animals to graze treated crops or cut for feed or hay within 7 days of application. Withdraw meat animals from treated feed 3 days prior to marketing. DO NOT graze timothy or cut for forage within 7 days of treatment.

Preharvest Interval: DO NOT apply within 60 days of harvest.

Re-cropping: Wheat, barley, oats, canola and peas may be grown the year following an application.

Aerial Application: DO NOT apply by air.

Storage: Store in dry, heated area. If frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	5	5	30

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on pages 15 to 16. When mixing with other pesticides, combine the method above with the method recommended for the tank mix partner if different from above for thorough cleaning.

Hazard Rating:

Frontline XL:



Warning – Potential skin sensitizer.

Florasulam SC:



Caution – Poison.

MCPA 600 Ester:



Warning – Poison

Both may cause eye and skin irritation.

For an explanation of the symbols used here see page 11.

Flucarbazone

Herbicide Group
2 - flucarbazone
(Refer to page 38)

Company:

Arysta LifeScience Canada (*Everest 2.0*)

Syngenta (*Sierra 2.0*)

Formulation:

Everest 2.0 (PCP#30342), *Sierra 2.0* (PCP# 30430): 397 g/L flucarbazone formulated as a suspension concentrate.

Container size - 1.94 L

Crops and Staging:

Spring wheat (including durum) with 1 leaf to a maximum of 4 main stem leaves plus 2 tillers (6 total leaves). Tank mixes containing 2,4-D provide improved crop safety. Tank mix options are listed in the tank mix section.

Note: Several of the tank mix partners have more limiting staging than flucarbazone. When tank mixing use the most restrictive application state or injury may result.

Weeds, Rates and Staging:

Grass weeds: Maximum of 4 main stem leaves and 2 tillers

Broadleaf weeds: 2 to 6 leaf stage

WEED	RATE	
	mL PER ACRE	ACRES PER 1.937 L
Green foxtail*, wild oat*, volunteer canola*	14.6	134
Weeds listed above plus: Wild oat* (light infestations) (< 100 plants/m ²), volunteer oat, green smartweed, redroot pigweed*, shepherd's-purse*, volunteer canola*, wild mustard*, stinkweed* (2 to 9 leaf stage)	19.4	100
Weeds listed above plus: Wild oat* (heavy populations) (> 100 plants/m ²) under ideal conditions for spraying.	24.3	80
Weeds listed above plus: Wild oat* (heavy populations) (> 100 plants/m ²) under poor environmental conditions or when mixing with <i>DyVel</i> or <i>Target</i> .	29.1	67

† *Everest* only

Requires the addition of a non-ionic surfactant (*Agral 90*, *Agsurf II*, *Surf 92*, *Super Spreader*, *LI700*) at 0.25 L per 100 L of spray solution.

* Will not control imidazolinone tolerant (CLEARFIELD) canola volunteers or Group 2 resistant weed biotypes. Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume:

Spring wheat: 20 to 40 L per acre.

Durum wheat and green foxtail control rate: 40 L per acre.

Nozzles and Pressure: Use 30 to 50 psi (200 to 345 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift.

Use of flood-jet or controlled droplet application equipment is not recommended due to poor coverage.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop tolerance and weed control may be reduced if applications are made to plants growing under stress. Stress includes saturated or water-logged soil, drought, extreme temperatures, low fertility or visible disease symptoms at application. Adopting practices to increase crop vigour will improve crop tolerance.

Tank Mixes:

Herbicides:

Note: All mixes must be applied with a registered surfactant unless otherwise indicated. Only one registered surfactant is required. Flucarbazone at all rates except 24.3 mL/acre can be used in the tank-mixes listed below, unless otherwise indicated.

In spring wheat (including durum):

2,4-D Amine or Ester at recommended rates up to

0.34 L/acre (500 g/L formulation)

Spectrum (20 acres per case)

In spring wheat (NOT including durum):

2,4-D Amine or Ester at recommended rates up to

0.45 L/acre (500 g/L formulation)

Ally at 2 to 3 g/acre + 2,4-D Amine or Ester up to rates recommended above[◇]

Buctril M (0.4 L/acre)

Curtail M (0.6 to 0.8 L/acre)[◇]

DyVel (0.5 L/acre)[◇]

Estaprop/Dichloroprop D (label rates)

Frontline 2,4-D (label rate)

MCPA Amine or Ester at recommended rates up to

0.45 L/acre (500 g/L formulation)

Pardner (0.4 L/acre)

Target (0.4 to 0.6 L/acre) [◇] ^{††}

Thifensulfuron/tribenuron 75DF forms

Thumper (0.4 L/acre)

Trophy (label rates)

[◇] Apply in 40 L/acre of water only.

^{††} Wild oat control may be reduced with this mix.

[#] Tank-mix only with the high wild oat rate of *flucarbazone* (29.1 mL/acre).

Fertilizers: None.

Insecticides: None

Note: The above mixes are those listed on the *flucarbazone* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Flucarbazone manufacturers may also support mixes with pesticides that are not on the *flucarbazone* labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall within 1 hour of application may reduce control.

Re-entry: DO NOT re-enter treated area within 12 hours.

Grazing: DO NOT graze treated fields. Mature grain or straw may be fed to livestock.

Preharvest Interval: Leave at least 80 days from application to harvest

Re-cropping Interval: Follow the chart below:

Soil Zones and Rotational Crops			
Grey-Wooded	Black	Dark Brown	Brown
Spring Wheat	Spring Wheat	Spring Wheat	Spring Wheat
Barley	Barley	Barley	
Canola (all varieties)	Canola (all varieties)	Canola (all varieties)	
Field Pea*	Durum Wheat	Durum Wheat	
	Field Pea*	Field Pea*	
	Flax	Flax	
	Field Bean		

* Field pea may be grown the year following *flucarbazone* application in fields where precipitation has been equal to or above the 10 year average during the growing season, and where organic matter content is above 4%, and pH is below 7.5. The company suggests a minimum of 100 mm (4 inches) of rain is needed in the 60 days following application for adequate breakdown to take place.

NOTE: Other rotational crops may also be affected if rainfall is less than the 10 year average for the area. Soils in the grey wooded, black and dark brown soil zones with a combination of low organic matter (less than 2%), light textured soils or high pfl (greater than 7.5) (i.e. eroded knolls, sandy soils) may result in delayed growth and development in rotational crops.

Aerial Application: DO NOT apply by air.

Storage: Store in closed original container in a cool, dry area away from fertilizers, food or feed. DO NOT freeze.

Buffer Zones: Leave at least 20 m from the downwind edge of the spray swath to sensitive upland plants like shelterbelts and woodlots and at least 35 m to water sources or wetland habitats. Avoid drift onto sensitive crops like canola and tame oat. DO NOT mix or load within 10 m of water sources or wetland habitats.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16. When mixing with other pesticides, combine the method above with cleanout methods for the tank mix partner.

Hazard Rating:



Warning – Contains the Allergen Milk

For an explanation of the symbols used here see page 11.

Flumioxazin (Chateau/Valtera)

Herbicide Group
14 - flumioxazin
(Refer to page 38)

Company:

Valent Canada, Inc.

Formulation:

51.1% flumioxazin formulated as a water dispersible granule.

Container size -

Chateau (PCP# 29231): 1.13 kg

Valtera (PCP# 29230): 2.27 kg

Crops, Rates, and Staging:

Valtera:

Soybeans: Prior to seeding or up to 3 days after seeding but prior to emergence.

If weeds are emerged apply Valtera in a mix with glyphosate (see tank mix section).

Dry Bean Desiccation:

Apply 42.5 g per acre when beans are mature to dry green weed material. Add metholated seed oil at 1 L per acre.

Chateau:

Potatoes: Apply after hilling. A minimum of 2 inches (5 cm) of soil must cover the vegetative portion of the potato or crop injury may result.

Weeds, Rates, and Staging:

Apply prior to crop and weed emergence.

WEEDS	Soil Type (all must have less than 5% organic matter)*	RATE (g/acre)
Chateau - In potatoes only.		
Suppression of: Pigweed (green, redroot) Common ragweed Lamb's-quarters Hairy nightshade Eastern black nightshade	Coarse and medium textured	42.5
Valtera - For soybeans only.		
Control of the weeds suppressed by Chateau above plus; Dandelion Green foxtail (suppression)	Coarse textured	56.7
	Medium textured	85

DO NOT apply on soils with > 5% organic matter, or fine soils. Soils such as clay, clay loam, silty clay or silty clay loam are considered fine textured soils. DO NOT apply to soils composed of more than 90% sand and gravel.

Spray within 6 hours of mixing.

Application Information:

Water Volume: Minimum application volume is not indicated on the label. Use appropriate water volumes to ensure good spray coverage.

Nozzles & Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

DO NOT perform any tillage operations after application otherwise weed control will be reduced.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Rainfall is required to activate *flumioxazin* in the soil. Crop injury may occur when soils are wet and cool following application or soils are poorly drained. Severe injury may occur with flooded soils. Newly emerging foliage can be temporarily injured by heavy rain splashing treated soil on leaves.

Tank Mixes:

Herbicides:

Soybeans only:

Glyphosate (IPA or K salts) (1.35 L/acre - 360 g/L formulations - see page 163 for rates for other glyphosate formulations).

Dry Bean Desiccation only:

Glyphosate (IPA or K salts) at preharvest rates.

Fertilizers: None registered.

Fungicides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rain or irrigation shortly after application is required for activation. If rainfall does not occur, irrigation with at least 5 mm of water is recommended before ground crack occurs.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Grazing: DO NOT graze or cut crops for livestock feed from treated fields.

Preharvest:

Desiccation: Leave 5 days between application and harvest. Leave 7 days to harvest if mixing with glyphosate.

Re-cropping: Soybeans may be seeded immediately after treatment. Winter wheat may be seeded in the fall following spring application. Alfalfa, barley, canola, field corn, sorghum, dry edible beans**, sunflower and spring wheat may be seeded the season after spring application. All other crops require a minimum of 12 months and a successful bioassay prior to indicate safe seeding.

** Note: Not all varieties of dry beans have been tested for recrop tolerance. Test new varieties of dry beans on a small area before attempting large acreages.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Crops	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Potato	2	1	5
Soybean	3	2	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16. See product label for further information.

Hazard Rating:



Caution - Poison

For an explanation of the symbols used here see page 11.

Fluroxypyr + 2,4-D

Herbicide Group
4 - fluroxypyr & 2,4-D
 (Refer to page 38)

Company:

Dow AgroSciences (*Attain* XC, *OcTtain* XL)
 Nufarm Agriculture (*Flurox* 2,4)
 MANA Canada (*Rush* 24)

Formulation:

The *Attain* XC package has 2 components:
Attain XC A (PCP# 29463): 333 g/L fluroxypyr
Attain XC B (PCP# 29264): 660 g/L 2,4-D LV ester
 Container sizes -
Attain XC A: 5L, or 8 x 15L
Attain XC B: 2 x 6.8 L, or 4 x 82L

-or-

The *Flurox* 2,4 package has 2 components:
Nufarm Fluroxypyr 180 (PCP# 30028): 180 g/L fluroxypyr.
2,4-D Ester 700 (PCP# 27820): 660 g/L 2,4-D LV ester.
 Container sizes -
Nufarm Fluroxypyr 180: 7.28 L
2,4-D Ester 700: 10.3 L

-or-

The *Rush* 24 package has 2 components:
Fluroxypyr 180 EC (PCP# 30815): 180 g/L fluroxypyr.
Salvo 2,4-D Ester 700 (PCP# 27818): 660 g/L 2,4-D LV ester.
 Container sizes -
Fluroxypyr 180 EC: 9.6 L
Salvo 2,4-D Ester 700: 2 x 6.8 L

-or-

OcTtain XL (PCP# 30077): 90 g/L fluroxypyr plus 360 g/L
 2,4-D LV ester.
 Container size - 2 x 9 L, 108L, 576 L

All products above are formulated as emulsifiable concentrates.

Crops and Staging:

Spring wheat (including durum), barley:
 4 leaf up to the emergence of the flag leaf.

Winter wheat: Apply to winter wheat in the spring from the 3 tiller stage to just before the flag leaf stage. (*Attain* XC and *OcTtain* XL only)

Forage Grasses for seed production only*:

Seedling and established grasses at the 4 leaf up to the emergence of the flag leaf.

Bromegrass	Timothy
(meadow, smooth)	Wheatgrass
Fescue (creeping red, tall)	(crested, intermediate)

* NOTE: Since these uses are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Those who apply these uses do so at their own risk.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

The following weeds are controlled at the 2 to 4 leaf stage, unless otherwise specified:

Attain XC A 95 mL/acre plus *Attain* XC B 260 mL/acre or; *Nufarm* or *MANA Fluroxypyr* 180 at 180 mL/acre plus *2,4-D Ester* 700 at 260 mL/acre (40 acres per case) controls the following weeds:

Bluebur	Plantain
Burdock	Prickly lettuce
Cleavers*	Ragweed
Clover (sweet)	Shepherd's purse
Cocklebur	Stinkweed
Field horsetail**	Sunflower (annual)
Flixweed†	Vetch
Goat's-beard	Volunteer canola
Hoary cress**	Wild radish
Kochia	Wild mustard
Lamb's-quarters	Wild buckwheat (1-4 leaf
Mustards (except dog and	with <i>Flurox</i> 2,4 and <i>Rush</i> 24,
tansy)	1-6 leaf with <i>Attain</i> XC and
	<i>OcTtain</i> only)

* 1 to 4 whorls with *Flurox* 2,4 and *Rush* 24; 1 to 6 whorls with *Attain* XC only.

OcTtain XL at 0.45 L per acre (2 x 9 L treats 40 acres, 108 L treats 250 acres, 576 L treats 1280 acres) controls:

All weeds listed above plus:

Common chickweed (up to 8 cm or 3 inches) [†]	Round-leaved mallow (1 to 6 leaf)
Cleavers (1 to 8 leaf)	Sow-thistle (perennial) ^{†**}
Hemp-nettle (2 to 6 leaf)	Stork's-bill (1 to 8 leaf)
Redroot pigweed [†]	Volunteer flax (1 to 12 cm)

Attain XC A at 130 mL per acre plus **Attain XC B** at 340 mL per acre (40 acres per case) or;
Fluroxypyr 180 at 240 mL per acre plus **2,4-D Ester 700** at 340 mL per acre (30 acres per case) or;

OcTtain XL at 0.45 L per acre (one 2 x 9 L case treats 40 acres, 108 L treats 250 acres and 576 L treats 1280 acres) plus **2,4-D ester (LV700)** at 81 mL/acre or **LV600** at 95 mL/acre) controls:
All weeds listed above plus:

Annual sow-thistle [†]	Leafy spurge ^{**}
Blue lettuce ^{**}	Oak-leaved goosefoot
Canada thistle ^{**†}	Perennial sow-thistle ^{**†}
Cleavers (1-8 whorls) [†]	Redroot pigweed
Dandelion ^{***}	Russian thistle
Docks	Smartweed (including Lady's-thumb)
Dog mustard	Tansy mustard
Field bindweed ^{**}	Tartary buckwheat
Field peppergrass	Wild buckwheat (1 to 4 leaf, 1 to 8 leaf with Attain XC , Rush 24 , and OcTtain XL only)
Gumweed	
Hairy galinsoga	
Hedge bindweed	
Hemp-nettle (2 to 6 leaf stage) ^{††}	

^A **Attain XC** and **OcTtain XL** only.

[†] Suppression only

^{††} Control with **OcTtain XL**, suppression with other products.

^{**} Top growth only

^{***} Spring rosettes only.

Make only one application per year of any of these products or other products containing the same active ingredients.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume:

Ground: **Attain XC** and **OcTtain XL** use 20 to 40 L per acre. All other uses minimum 40 L per acre.

Aerial: **Attain XC** and **OcTtain XL** only use 12 to 20 L per acre.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper

performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of **ASABE coarse** droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

The activity these products are influenced by weather conditions. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions (drought, heat or cold stress) or if extremely heavy infestations exist.

Tank Mixes:

Herbicides:

The following mixes may be used with each of the combinations above unless noted otherwise.

In spring wheat (including durum) and barley:

Tralkoxydim[†] (0.2 L/acre) plus adjuvant^{*}

Assert (0.53 to 0.65 L/acre)

In spring wheat (including durum) only:

Clodinafop 240EC[†] (93 mL/acre) plus **Score** adjuvant[†]

Simplicity (0.15 to 0.20 L per acre)^{††}

Insecticides: None registered.

Fungicides: None registered.

Fertilizer: None registered.

^{*} Temporary crop injury or reduced wild oat control may occur with this tank mix.

[†] See product labels for specific brands registered.

^{††} Low rate of **Attain XC** or **OcTtain XL** without additional **2,4-D** ester only.

Note: The above mixes are those listed on the **fluroxypyr + 2,4-D** product labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Manufacturers may also support mixes between their products and other pesticides that are not on their labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall:

Attain XC and Flurox 2,4: Within 2 hours will reduce control.

OcTTain XL: Wet foliage at application will reduce control.

Re-entry: DO NOT re-enter treated area within 12 hours.

Grazing: DO NOT permit lactating dairy animals to graze cereal fields within 7 days of application. DO NOT harvest cereal crops for forage or cut hay within 30 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

DO NOT feed or cut forage grasses for hay

Preharvest Interval: Leave 60 days between application and harvest.

Re-cropping: Barley, canola, flax, forage grasses, lentils, mustard, oats, peas, rye, and wheat, may be grown the year after application. Alfalfa, corn, dry beans, potatoes, soybeans, and sunflowers may also be seeded the year following *Attain XC* and *OcTTain XL* only. There are no re-cropping restrictions the second year after application.

Aerial Application: *Attain XC* and *OcTTain XL* may be applied by air.

Storage: Avoid freezing. If frozen, bring to room temperature and agitate before use. These products are combustible. DO NOT store near heat or open flame.

Buffer Zones:

Flurox 2,4, Rush 24: Leave a buffer of 15 meters from water bodies, wetland areas and plants that may be injured.

Attain XC:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	3
Fixed wing aircraft	6	0	100
Helicopter	6	0	80

OcTTain XL only:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Freshwater Habitat of Depths:		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Field sprayer	1	0	1
Fixed wing aircraft	1	0	55
Helicopter	1	0	45

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

Attain XC, Flurox 2,4 and Rush 24:



Danger – Poison.

OcTTain XL:



Caution – Poison

All products:



Warning – Eye Irritant.



Caution – Skin Irritant.

For an explanation of the symbols used here see page 11.

Focus

Herbicide Group
14 – carfentrazone
15 – pyroxasulfone
(Refer to page 38)

Company:

FMC of Canada

Formulation:

The *Focus* package contains:

Aim EC Herbicide (PCP# 28573): 240 g/L carfentrazone formulated as an emulsifiable concentrate.

Container size - 1 x 2 L

Pyroxasulfone 85 WG (PCP# 30572): 85% pyroxasulfone formulated as a water dispersible granule.

Container size - 2 x 2 Kg

Crops and Staging:

Corn (field) - Apply prior to seeding or up to 3 days after seeding.

Soybeans - Apply prior to seeding or up to 3 days after seeding.

WARNING – application to emerged corn or soybeans will result in severe damage to the crop. *DO NOT use on peat or muck soils and soils with 7% or more organic matter content.*

Weeds and Staging:

Control of the following weeds emerging from seed (not controlled if emerged at application):

Barnyard grass	Ryegrass (Italian)
Foxtail (green, yellow)	Waterhemp (Common)
Redroot pigweed	

Control of the following weeds up to 10 cm tall (post-emergent):

Cocklebur	Purslane
Kochia	Round-leaved mallow
Lamb's-quarters	Stinkweed
Nightshade (eastern black, hairy)	Tansy mustard
Pigweed (prostrate, red-root, smooth, tumble)	Volunteer canola (all varieties)

Rates:

Aim: 30 to 36 mL per acre.

Pyroxasulfone 85 WG: 60 to 73 g per acre.
 (66 to 55 acres per case)

Note: Focus may only be applied once per growing season.

Use Agral 90 or Ag-Surf at 0.25 L per 100 L of spray solution or Merge at 1 L per 100 L of spray solution.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 14.

Application Information:

Water Volume: Minimum of 40 L per acre. Higher spray volume is required for dense weed stands. Weed control improves with the amount of coverage.

Nozzles & Pressure: Maximum 35 psi (210 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Moisture is necessary to activate the *Pyroxasulfone* component in soil for effective weed control. Dry weather following applications may reduce effectiveness.

Extremes in environmental conditions such as temperature, moisture, soil conditions, and cultural practices may affect activity.

Tank Mixes:

Herbicides:

Prior to crop emergence:

AAtrex (0.85 to 1.25 L per acre) (soil activity).

*Glyphosate** (0.5 to 1L per acre of 360 g/L formulation)

*IPA or K salt only.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall 6 to 8 hours after application may reduce activity of the *Aim* component. Heavy rainfall shortly after application may reduce weed control. Moderate rainfall beyond the above limitations will improve the activity of the *Pyrooxasulfone* component.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Grazing: DO NOT graze the treated crop or cut for feed.

Preharvest Interval: Not applicable.

Re-cropping: Conduct a field bioassay to confirm crop safety prior to seeding any rotational crops other than field corn or soybeans.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place in original container.

Buffer Zones:

Crops	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	5	3	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Warning – Contains the allergen, sulfites.

Potential skin sensitizer

For an explanation of the symbols used here see page 11.

Fortress

Herbicide Group

3 - trifluralin

8 - triallate

(Refer to page 38)

Company:

Gowan Canada (PCP# 19521)

Formulation:

10% triallate and 4% trifluralin formulated as a granular.
Container size - 22.7 kg bag, 454 kg container.

Crops and Staging:

Prior to planting wheat (spring and durum), barley, canola, flax (not including Solin), mustard.

Preplant incorporated: In fall after September 15 until soil freeze-up or in the spring prior to seeding crop.

Surface application: Apply in the fall after October 1 and when soil temperature is less than 4°C at a depth of 2 inches (5 cm) and delay incorporation until the following spring.

DO NOT apply to fields with heavy trash cover or after snow has fallen. Some wheat or barley injury may be noted on eroded knolls.

Weeds and Staging:

Pre-emergent control of wild oats, green foxtail, yellow foxtail.

Suppression of lamb's-quarters, kochia, redroot pigweed, Russian thistle, wild buckwheat.

Rates:

Apply *Fortress* according to soil organic matter content.

ORGANIC MATTER	RATE (KG/ACRE)					
	RAPESEED (CANOLA), FLAX***, MUSTARD		BARLEY		SPRING WHEAT AND DURUM WHEAT	
	Spring	Fall	Spring	Fall	Spring	Fall
< 2%	5.67	5.67	N.R.*	4.45	N.R.*	N.R.*
2 -4%	5.67	5.67	4.45	5.67	N.R.*	4.45
4 -6%	6.88	5.67	5.67	5.67	4.45	5.67
> 6%	6.88	6.88	6.88	6.88	5.67	5.67**

* N.R. -Not Recommended.

** For fall incorporated applications (not surface) apply 6.88 kg/acre when organic matter exceeds 8 percent.

*** Excluding Solin (low linolenic acid flax).

Application Information:

Fortress may be applied in the fall with or without a fall tillage operation, or in the spring as a preplant incorporated treatment. Before application of this product, the soil must be in good working condition. Application to a field that is wet, lumpy, rough or ridged will result in reduced weed control and promote crop thinning.

Fall Surface Application: Where fields are prone to water and/or wind erosion, and tillage is therefore undesirable, fall surface application should be made within 3 weeks of soil freeze-up, when the soil begins to cool (less than 4°C), which typically begins on or around October 1. Application can be made to standing stubble or to previously worked fields with incorporation delayed until spring. For best results on heavy wild oat infestations, use the incorporated treatment.

Fall Incorporated Application: *Fortress* must be applied after September 15 and before soil freeze-up. Application prior to September 15 may result in reduced weed control. Initial incorporation may be completed within 24 hours of application. The second incorporation may be done in the fall (prior to soil freeze-up) or in the spring prior to, or after, seeding. If performed after seeding, it must be completed with harrows prior to emergence of the crop. Fall incorporation is not recommended on soils where a lack of trash cover combined with the required incorporation operation could result in soil erosion.

Spring Application: *Fortress* can be applied before seeding but must be incorporated within 24 hours of application. The second incorporation must be delayed at least 48 hours after the first and may be performed at any time prior to crop emergence.

Incorporation:

Fortress applications require two incorporations, with the second incorporation at right angles to the first. Seeding with a seeder that provides soil disturbance equivalent to a cultivator may replace one incorporation. Incorporate to a maximum depth of 2 inches (5 cm) by setting disk or cultivator implements to cut a maximum of 3 inches (7.5 cm) into the soil. Mixing the product to greater depths will dilute the herbicide, decrease wild oat control, and may cause injury to cereals. If the second incorporation is conducted after seeding, it should be done with harrows or other suitable tillage equipment adjusted so as not to disturb the seed. Harrowing does not provide effective incorporation if compact soil prevents penetration of harrow teeth, if trash accumulates in the harrow sections, or if the harrows bounce.

Seeding Requirements: Accurate seeding depth control is critical. Thinning of wheat and barley has been known to occur when seeding depth has been inadequate. Ensure that cereals are seeded below the treated layer (2 to 3 inches or 5 to 7.5 cm). Do not seed deeper than 3 inches (7.5 cm). To ensure an even crop stand, increase the usual seeding rate of wheat or barley by 10 percent, especially if soil conditions are cold or dry. See product label for more information.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop injury can occur on fields where *Fortress* has been applied and heavy rainfall or cold weather occur after seeding but prior to crop emergence. Seeding under warm soil conditions (greater than 10°C and generally after May 15) will ensure optimum crop germination and emergence and will reduce the risk of crop injury. Very dry conditions in spring or prolonged cool soil temperatures at time of wild oat germination will result in reduced control. Poor results may be expected from incomplete incorporation due to wet, cloddy soil or heavy trash. Ridges left at seeding may disrupt the treated layer and allow weed escapes.

Restrictions:

Rainfall: Moisture is required for activation. Rainfall of at least 0.6 inches (1.5 cm) within 2 weeks of seeding is required to ensure optimum results.

Re-entry: DO NOT enter treated fields for at least 12 hour

Grazing: DO NOT graze or cut treated crops for livestock feed prior to crop maturity.

Re-cropping: *Fortress* will leave a residue in the soil. Oats, canaryseed, and small seeded forage grasses may be injured if planted within 24 months of application. DO NOT apply *Fortress* on land to be sown to wheat if the land has been treated with trifluralin since June 1 of the previous year.

Aerial Application: May be applied by airplane with attachments designed for applying low volumes of granules.

Storage: Store in a cool, dry place.

Hazard Rating:



Warning – contains the allergen soy.

May cause Skin and Eye Irritation

For an explanation of the symbols used here see page 11.

Frontier Max

Herbicide Group
15 - dimethanamid
(Refer to page 38)

Company:

BASF Canada (PCP# 29194)

Formulation:

720 g/L dimethanamid-P formulated as an emulsifiable concentrate.

Container size - 3 L to 1000 L.

Crops and Staging:

Pre-plant incorporated:

Corn (NOT sweet corn, popcorn, or corn grown for seed).
Dry beans (white and kidney beans only).

Pre-emergence surface:

Dry beans (white and kidney beans only).

Weeds and Staging:

Pre-emergent control of green foxtail.

Rates:

Pre-plant incorporated treatments:

Apply at 0.35 to 0.39 L per acre. Apply at the higher rate on fine-textured or high organic soils and for heavier anticipated weed problems.

Pre-emergence surface treatments:

SOIL TYPE (Texture)	RATE (L PER ACRE)		
	Less than 3% Organic Matter	3 to 6% Organic Matter	7 to 10% Organic Matter
Coarse	0.31	0.31	0.35
Medium and Fine	0.31	0.35	0.39

Application Information:

Water Volume: A minimum of 40 L per acre.

Pressure: 30 to 43 psi (200 to 300 kPa).

Nozzles: Flat fan or flood-jet. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets. Use 16 mesh suction screen, 50 mesh elsewhere on sprayer.

Incorporation: For pre-plant incorporated treatments, apply *Frontier Max* as a broadcast treatment and incorporate using a harrow, rolling cultivator or other implement capable of giving uniform, shallow incorporation into the top 5 cm (2 inches) of soil within 7 days of planting. Avoid deeper incorporation or reduced weed control and/or crop injury may result. Immediate incorporation after application is not necessary.

Beans must be planted at least 4 cm (1.5 inches) deep or crop injury may occur.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Rainfall is required within 7 to 10 days of application to activate and move *Frontier Max* into the soil zone. If dry conditions persist, a shallow cultivation or the use of a rotary hoe is necessary to move the herbicide into moist soil and control weed escapes. Shallow tillage is important to minimize dilution of the herbicide. If drought conditions persist after pre-plant incorporated or pre-emergence applications, weed control may not be adequate.

Tank Mixes:

Herbicides: None registered.

Fertilizers: May be applied with a liquid fertilizer carrier. Test compatibility with liquid fertilizer by mixing a small amount of herbicide with a proportional quantity of liquid fertilizer in a jar. May also be impregnated on dry bulk fertilizers for pre-plant incorporated treatments. A minimum of 90 kg/acre of dry bulk fertilizer should be applied. DO NOT impregnate *Frontier Max* on nitrate fertilizers, superphosphates or limestone.

Insecticides: None registered.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall after application is important for good weed control.

Re-entry: DO NOT enter treated fields for 24 hours.

Grazing: DO NOT graze or feed the treated corn crop within 40 days of application. DO NOT graze the treated bean crop or feed bean forage, hay or straw to livestock.

Re-cropping: DO NOT plant winter wheat within 120 days of application.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Must be stored under heated warehouse conditions.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	3

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on pages 15 to 16. When mixing with other pesticides, combine the method above with the method recommended for the tank mix partner if different from above for thorough cleaning.

Hazard Rating:



Caution - Poison.



Warning - Eye Irritant and Potential Skin Sensitizer.

For an explanation of the symbols used here see page 11.

Frontline 2,4-D XC

Herbicide Group
2 - florasulam
4 - 2,4-D
(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

The Frontline 2,4-D XC package has 2 components:

Frontline 2,4-D XC A (PCP#30060): 50 g/L florasulam formulated as a suspension concentrate

Frontline 2,4-D XC B (PCP#30061): 660 g/L 2,4-D LV ester formulated as an emulsifiable concentrate.

Container sizes:

Frontline 2,4-D XC A: 1.6 L.

Frontline 2,4-D XC B: 2 x 6.8 L

Crops and Staging:

Spring wheat (including durum) - 3rd leaf fully expanded to 6 leaf stage.

When mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Bluebur	Ragweed (common)
Burdock	Redroot pigweed
Chickweed	Russian thistle
Cleavers	Shepherd's-purse
Cocklebur	Smartweed
Dandelion*	Sow-thistle (annual)
Flixweed	Stinkweed
Lady's-thumb	Sunflower (annual)
Lamb's-quarters	Tartary buckwheat
Mustard (ball, wild)	Vetch
Narrow-leaved hawk's-beard***	Volunteer canola†
Plantain	Wild buckwheat
Prickly lettuce	Wild radish

Broadleaf weeds suppressed:

Canada thistle (top growth control only)	Perennial sow-thistle (top growth control only)
Hemp-nettle	

* Seedlings and overwintered rosettes

*** Up to 2 leaf stage.

† Including all herbicide-tolerant canola varieties

Rates:

Frontline 2,4-D XC A: 40 mL per acre

Frontline 2,4-D XC B: 0.34 L (340 mL) per acre.

One package treats 40 acres.

Note: Maximum one application of this product or other products containing florasulam in back to back years.

Refer to the product label for complete mixing instructions for this product and its mixes.

A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) with conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity of Frontline 2,4-D. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and re-growth may occur. Under conditions of low crop and high weed density, control may be reduced. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application may reduce weed control and increase the risk of crop injury at all stages of growth.

Tank Mixes:

Herbicides:

In spring wheat (including durum)

Assert 300 SC (0.65 L/acre) plus Acidulate

Insecticides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the Frontline 2,4-D XC label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information. DO NOT apply excessive irrigation following application as product may leach.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT allow lactating dairy animals to graze treated areas within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter. DO NOT harvest forage or cut hay within 30 days of application.

Preharvest Interval: Leave 60 days between application and harvesting mature crop.

Re-cropping: Wheat, barley, canola, oats and peas may be grown the year following an application.

Aerial Application: DO NOT apply by air.

Storage: Store in dry, heated area. If frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	5	5	30

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Warning – Poison



May cause skin and eye irritation.

For an explanation of the symbols used here see page 11.

Frontline XL (this referring text to be removed in the 2017 edition)

See florasulam + MCPA Ester on page 162.

Glyphosate

Herbicide Group
9 - glyphosate
(Refer to page 38)

Product names, Company, Formulation and Package sizes:

All products are formulated as solutions.

Product Name	Company	Salt type*	Active** content (g a.e./L)	Package sizes***
Cheminova Glyphosate (PCP# 26828)	Cheminova	IPA	356	2, 3, 7
ClearOut 41 Plus (PCP# 28322)	Farmers of North America	IPA	360	2, 7
Credit 45 (PCP# 29124)	Nufarm Agriculture	IPA	450	1, 2, 7
Crush'R Plus (PCP# 29995)	AgWest	IPA	360	1, 2, 3
Glyfos (PCP# 24359)	Cheminova	IPA	360	1, 2, 3, 7
Knockout Extra (PCP# 29266)	Great Northern Growers	IPA	360	1, 2, 5
Laji Plus (PCP# 29677)	Ray Glenn Commodities	IPA	360	1, 2
Matrix (PCP# 29775)	IPCO	DMA	480	1, 2, 3, 7
Maverick III (PCP# 28977)	Dow AgroSciences	DMA	480	1, 2, 3, 6
MPower Glyphosate (PCP# 29290)	Farmers of North America	IPA	360	2, 4, 7
NuGlo (PCP# 29470)	Nufarm Agriculture	IPA	450	3, 7
Roundup Transorb HC (PCP# 28198)	Monsanto	K+	540	1, 2, 3, 7, 10
Roundup Ultra 2 (PCP# 28486)	Monsanto	K+	540	1, 2, 3, 8
Roundup WeatherMax (PCP# 27487)	Monsanto	K+	540	1, 2, 3, 7, 10
R/T 540 (PCP# 28487)	Monsanto	K+	540	1, 2, 5, 10
Sharpshooter (PCP# 28631)	UAP	IPA	356	2, 5, 7
Sharpshooter Plus (PCP# 28623)	UAP	IPA	360	1, 2, 3, 4
StartUp (PCP# 29498)	Loveland Products (formerly Viterro)	K+	540	1, 2, 3, 4
Touchdown Total (PCP# 28072)	Syngenta Canada	K+	500	1, 2, 3, 7
Traxion (PCP# 29201)	Syngenta Canada	K+	500	1, 2, 3
Vantage Plus Max II (PCP# 28840)	Dow AgroSciences	DMA	480	2, 6, 7
Vector (PCP# 30319)	IPCO (for Federated Cooperative)	DMA	480	1, 2, 3, 6
Wise-Up (PCP# 29126)	Adjuvants Plus	IPA	356	2, 5

* Salt type: IPA = Isopropylamine, MA = Monoammonium, DA = Diammonium, DMA = dimethylamine, K+ = Potassium

** Formulation concentration is expressed as "grams of acid equivalent per litre of product (g a.e./L).

Glyphosate acid is the herbicidally active component of the formulation and is proportional to the activity of the formulation
 Note: Some products may be more effective due to formulation differences (not related to higher glyphosate content) under adverse conditions, but that benefit is reduced when applications are made under optimal conditions for activity (i.e. rapid weed growth, clean leaf surfaces). When selecting a glyphosate product, consult the product

*** Container sizes available: 1) 2 x 10 L or 1 x 20L, 2) 115 L, 3) 450 L, 4) 667 L, 5) 750 L, 6) 960 L, 7) 1000 L, 8) 1150 L, 9) 1200 L, 10) 800 L.

Product volumes (per acre) for various formulations strengths based on grams acid equivalent (gae) rate

RATE (GAE per ACRE)	Glyphosate formulation concentration (gae/L)				
	356/360	450	480	500	540
36.5	100 mL	81 mL	76 mL	73 mL	67 mL
73	200 mL	162 mL	152 mL	146 mL	134 mL
110	0.3 L	0.24 L	0.23 L	0.2 L	0.2 L
120	0.33 L	0.27 L	0.25 L	0.24 L	0.22 L
145	0.4 L	0.32 L	0.3 L	0.28 L	0.27 L
180	0.5 L	0.4 L	0.38 L	0.36 L	0.33 L
275	0.77 L	0.61 L	0.57 L	0.54 L	0.51 L
325	0.91 L	0.73 L	0.68 L	0.65 L	0.61 L
360	1.0 L	0.81 L	0.76 L	0.73 L	0.67 L
510	1.42 L	1.13 L	1.1 L	1.0 L	0.94 L
540	1.5 L	1.21 L	1.13 L	1.09 L	1.0 L
650	-	-	-	1.30 L	-
690	1.9 L	1.54 L	1.44 L	1.38 L	1.28 L
720	2.0 L	1.62 L	1.5 L	1.46 L	1.34 L
1020	2.8 L	2.27 L	2.13 L	2.02 L	1.89 L
1750	4.9 L	3.88 L	3.6 L	3.48 L	3.24 L

Crops and Uses:

1. Annual weed control prior to crop emergence or in summerfallow.
2. Quackgrass control prior to seeding or after harvest.
3. Dandelion control (other than Preharvest).
4. Canada thistle control in summer-fallow, shelterbelts and post-harvest.
5. Alfalfa control (other than Preharvest).
6. Other perennial weeds control in summerfallow, shelterbelts and post-harvest.
7. Patch treatments of perennial weeds in cereals, corn, soybean and forages.
8. Preharvest perennial weed control.
9. For use in Glyphosate tolerant crops.
10. Tank Mixes.

1. Annual weed control prior to crop emergence or in summer-fallow:

Weeds listed may not occur on all product labels. Check individual product labels for a specific list of weeds controlled.

RATE (gae per acre)	SURFACTANT*	WEEDS CONTROLLED	WEED STAGE
110	0.14 L/acre	Grasses: Green foxtail, volunteer cereals, wild oat (light infestations) Broadleaves: lady's-thumb, stinkweed, volunteer canola (NOT including glyphosate tolerant varieties), wild mustard.	Less than 3 inches (8 cm) high. Apply at the 1 to 3 leaf stage of wild oat.
145	0.14 L/acre	<i>Above weeds plus:</i> Grasses: heavy infestations of wild oat. Broadleaves: suppression of flaxweed, kochia.	1 to 3 leaves for wild oat Weeds 3 to 6 inches (8 to 15 cm).
180 to 275	Not required	<i>Above weeds plus:</i> Grasses: downy brome, Persian dandel. Broadleaves: Canada fleabane, cleavers, common ragweed, flaxweed, hemp-nettle, lamb's quarters, narrow-leaved hawk's-beard, redroot pigweed, Russian thistle, volunteer flax, wild buckwheat.	Canada fleabane, common ragweed, less than 3 inches (8 cm) high. Other weeds less than 6 inches (15 cm). Use high rate for narrow-leaved hawk's-beard 3 to 6 inches (8-15 cm) or wild buckwheat at the 3-4 leaf stage.
325	Not required	<i>Above weeds plus:</i> Grasses: annual blue grass, crabgrass. Broadleaves: annual sow-thistle, kochia, prickly lettuce, shepherd's-purse, narrow-leaved vetch**.	Less than 6 inches (15 cm) high
510	Not required	<i>Above weeds.</i>	Greater than 6 inches (15 cm) high

* Unless otherwise specified on the product label, use one of the following surfactants: Agral 90, Agsurf II, Companion or LI700.

** Note: Narrow-leaved vetch is an annual species. Establish perennial vetches such as American vetch, may not be controlled at this rate.

2. Quackgrass control prior to seeding or after harvest:

RATE (gae per acre)	QUACK GRASS STAGE
360	Season long control of light to moderate infestations. Apply when quack grass is 8 inches (20 cm) tall and has 3 to 4 actively growing leaves. Apply spring or fall.
360 to 1020	Apply when quack grass has 3 to 4 new leaves for long term control of heavy infestations. Use high rate for sod-bound quack grass (left undisturbed for at least 2 years).

DO NOT apply fall treatments if a hard frost has occurred (-5°C) or if plants are drought stressed. Spread straw to allow regrowth and good spray coverage.

Cultivation prior to application will result in reduced control. DO NOT cultivate between harvest and treatment when using fall applications. If using spring applications on fields which have been fall-tilled, delay application until the quack grass has reached the 4 to 5 leaf stage. (This will occur 1 to 4 weeks later on fall-tilled fields than in undisturbed fields).

Cultivation after application usually will improve control of quack grass. Wait a minimum of 3 days after application before cultivating. If growing conditions are poor (cold or dry), particularly in the fall, waiting longer than 5 days may improve control.

3. Dandelion control (other than Preharvest):

Apply up to and including dandelion bloom for best results.

RATE (gae per acre)	DANDELION GROWTH STAGE
360	Less than 6 inches (15 cm) diameter. Allow 3 or more days after treatment before tillage.
540 to 720	Greater than 6 inches (15 cm) diameter. Use higher rate when infestations are heavy.

4. Canada thistle control in summer-fallow, shelterbelts and post-harvest:

RATE (gae per acre)	WEED STAGING
360	Rosettes at least 6 inches (15 cm) in diameter, treated in late summer, following tillage in spring and early summer (up to August 1). Allow thistles to regrow for 5 weeks following last tillage. Wait a minimum of 10 days after application before tillage. Treatment after a mild frost is possible if leaves are green and pliable and plants are actively growing.
690 to 1020	Bud stage or beyond. Allow at least 5 days after application before tillage. -or- Post-harvest stubble treatment. Allow 8 to 10 inches (20 to 25cm) of new growth before application. Must be sprayed at least 2 weeks prior to killing frost. Straw should be removed or evenly spread to allow for proper regrowth and spray coverage.

5. Alfalfa Control (other than Preharvest):

RATE (gae per acre)	WEED STAGING
540 to 720	Fall control of alfalfa in early bud to full bloom stage. Use high rate when alfalfa populations are high or when perennial grasses are present. Allow at least 5 days before tillage. See tank mix section for minimum tillage or spring applications. Apply with 23 to 135 L per acre water.
<i>Touchdown Total and Traxion only:</i> 325 to 650	Prior to seeding or after harvest. Use higher rate for weeds beyond 3 inches (8 cm) in height or for heavy weed infestations. Wait 7 days after application for tillage. Apply in 23 to 135 L per acre water.

6. Other perennial weed control in summerfallow, shelterbelts and post-harvest:

(Refer to individual product labels for detailed application information.)

Foxtail Barley: Suppression with *Touchdown* and *Traxion* only at 0.28 L per acre.

Control from seedling to heading (all products) at 360 to 720 gae per acre. Late fall applications may provide better control of established foxtail barley plants than spring applications.

Yellow toadflax: 360 gae per acre.

Other Perennial weeds*: 1020 to 1750 gae per acre

* Perennial weeds such as absinth, blue grass spp., smooth brome grass, cattail, curled dock, field bindweed (bloom stage or beyond), hemp dogbane, hoary cress, poison ivy, purple loosetrife, perennial sow-thistle, and yellow nut-sedge applied at the early heading to early bud stage.

7. Patch treatments of perennial weeds in wheat, oat, barley, corn, soybean, forage legumes and forage grasses:

(Refer to individual product labels for detailed application instructions)

RATE (gae per acre)	WEED
360 to 1020	Quack grass 8 in (20 cm) tall
690 to 1020	Canada thistle Bud or beyond
1750	Milkweed Bud to bloom
1020 to 1750	Other perennial weeds*
36.5 to 73	Spot treatment rates for hand held equipment (per 10 L water**)

* Perennial weeds such as absinthe, blue grass spp., smooth brome grass, cattail, curled dock, field bindweed (bloom stage or beyond), hemp dogbane, hoary cress, poison ivy, purple loosetrife, perennial sow-thistle, and yellow nutsedge applied at the early heading to early bud stage.

** Use the low rate for quack grass and the high rate for all other perennials.

8. Preharvest perennial weed control:

DO NOT apply to any crops grown for seed.

Not all glyphosate products are registered for Preharvest applications on all crop species listed below. Refer to specific glyphosate labels for a list of registered uses and crop species.

RATES:

Prior to the harvest of annual grains (see staging chart below for specific crops): 360 gae per acre.

Prior to the final cut of forages to be removed from production: 360 to 720 gae per acre.

Weeds Controlled with Preharvest applications:

Quack grass 4-5 green leaves	Canada thistle and perennial sow-thistle at bud stage or beyond	Common milkweed at bud to bloom stage	Toadflax at bud to full bloom stage	Dandelion from rosette to full bloom stage
X	X	X	X	X

Crop Staging for Preharvest applications:

Apply to crops (except forage) when grain moisture is less than 30%. The following chart lists visual symptoms that can be used as guidelines to when 30% grain moisture has been reached.

CROP*	VISUAL SYMPTOMS
Wheat, Barley*, Oat*	Hard dough stage – a thumbnail impression remains on seed.
Canola, Mustard***†	Pods are green to yellow and most seeds are yellow to brown.
Flax (including Solin)	Majority (75 to 80% of bolls) are brown.
Lentil	Lowermost pods (bottom 15%) are brown and rattle when shaken.
Pea	Majority (75 to 80%) of pods are brown.
Chickpea***†	Stems are green to brown in colour: pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves).
Lupin**	Stems are green to brown in colour: pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves).
Faba bean***†	Stems are green to brown in colour: pods are mature (yellow to brown in colour); 80%-90% leaf drop (original leaves).
Soybean	Stems are green to brown in colour and pod tissue is brown and dry in appearance (80 to 90% leaf drop).
Dry Bean	Stems are green to brown in colour and pods are mature (yellow to brown) and 80 to 90% of the original leaves have dropped.
Forage	3 to 7 days prior to the last cut before rotation or forage renovation. DO NOT apply to forage stands that are to be maintained.

* Registered for application to barley grown for malt and tame oat grown for milling. Contact malt barley or milling oat buyers prior to application to confirm acceptance of glyphosate-treated grain.

** Preharvest applications on these crops are registered with Roundup Transorb HC, Roundup WeatherMax, R/T 540, StartUp and Roundup Ultra 2 only.

*** (yellow / white, brown, oriental), RoundUp Weather Max only.

† NOTE: Since these uses are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Those who apply glyphosate to chickpea, lupin, fababeans, or mustard do so at their own risk.

9. For use in glyphosate tolerant canola:

Weeds, Staging and Rates:

All applications must be made within the cotyledon to 6 leaf stage of glyphosate tolerant canola. Temporary yellowing may occur if applied at the 4 to 6 leaf stage of the crop.

Not all glyphosate products are registered for use on glyphosate tolerant canola at all rates listed. Refer to individual product labels for specific uses and rates.

Single applications of 120 gae per acre:

Weeds controlled at all stages unless indicated otherwise:

Annual grasses: barnyard grass, green foxtail, volunteer cereals, wild oat.

Annual broadleaves: annual smartweed spp.**, chickweed, corn spurry, cow cockle*, hemp-nettle, kochia, lamb's-quarters, night-flowering catchfly*, redroot pigweed, Russian thistle, shepherd's-purse*, stinkweed, volunteer canola

(except glyphosate tolerant varieties), wild mustard, wild tomato.

Single applications of 180 gae per acre:

All stages of the weeds listed above plus:

Annual broadleaves: cleavers, flaxweed, wild buckwheat, stork's-bill, narrow-leaved hawk's-beard.

Perennial weeds suppressed: Canada thistle, dandelion, perennial sow-thistle, and season long quack grass control.

Double application of 180 gae per acre plus 180 gae per acre:

Additional flushes of the weeds Listed above plus:

Annual broadleaves: round-leaved mallow

Season long control of following perennials: Canada thistle, foxtail barley, and perennial sow-thistle.

Single application of 270 gae per acre:

All weeds in single applications above plus:

Season long control of following perennials: Canada thistle and perennial sow-thistle.

* Low rates can be used only up to the 3 leaf stage of the crop otherwise use the high rate.

** Low rates can be used only when annual smartweed is in the 4 to 6 leaf stage.

NOTE: A maximum of 360 gae per acre per season is allowed in glyphosate tolerant Canola

10. For use in glyphosate corn and soybean:

Weeds, Staging and Rates:

All applications must be made within the following crop growth stages.

Corn - up to and including 8 leaf stage

Soybean - first trifoliate leaf through flowering.

Not all glyphosate products are registered for use on glyphosate tolerant corn and soybeans at all rates listed. Refer to individual product labels for specific uses and rates.

Single application of 360 gae per acre controls the following weeds:

Grasses:

Barnyard grass	Quack grass
Crabgrass spp.	Volunteer barley and wheat
Foxtail (green, yellow, giant)	Wild oats
Proso millet	

Broadleaves:

Biennial wormwood*	Night-shade, eastern-black
Canada thistle	Perennial sow-thistle
Chickweed	Pigweed (smooth, redroot)
Cleavers	Round-leaved mallow
Corn spurry	Russian thistle
Cocklebur	Shepherd's-purse
Cow cockle	Smartweed spp.
Common milkweed (suppression only)	Stinkweed (suppression only)
Common ragweed	Stork's-bill
Flixweed	Velvetleaf
Hemp-nettle	Volunteer canola (except glyphosate tolerant varieties)
Kochia	
Lamb's-quarters	Wild mustard
Narrow-leaved hawk's-beard	Wild buckwheat
Night-flowering catchfly	Wild tomato

* Registered for control in glyphosate tolerant soybean only with Roundup products and R/T 540 only.

Second applications of 360 gae per acre controls the following weeds:

Late flushes of heavy infestations of the above weeds plus control of:

Common milkweed	Round-leaved mallow
Field bindweed	Yellow nutsedge

Single application of 720 gae per acre in glyphosate tolerant soybean from the first trifoliate to flowering stage and corn up to and including 6 leaf stage:

Heavy infestations of the annual weeds listed above plus control of:

Field bindweed	Perennial sow-thistle
Common milkweed	Yellow nutsedge
Canada thistle	

** The single application rate in glyphosate tolerant corn and soybean is not labeled for all glyphosate products. Refer to individual glyphosate labels for the registration status of this rate usage in glyphosate tolerant soybean and corn.

11. Tank Mixes:

Not all glyphosate products are registered for all tank mix options below. Refer to individual glyphosate labels for registered tank mixes, glyphosate rates and registered crop species.

	RATE PER ACRE
Preseeding canola††	<i>Bromoxynil-Pardner</i> (0.51 L), <i>Koril</i> (0.48), <i>Brotex</i> (0.6 L)
Preseeding cereals***:	2,4-D [◇] (0.23 to 0.58 L)*
	<i>Banvel II</i> (0.12 L)*
	<i>Bromoxynil - Pardner</i> (0.51 L), <i>Koril</i> (0.48), <i>Brotex</i> (0.6 L)
	MCPA [◇] (0.2 to 0.4 L)*
	<i>Bromoxynil</i> / MCPA* - <i>Buctril M</i> (0.2 to 0.4 L), <i>Logic M</i> (0.25 to 0.5 L)
Preseeding corn (field and sweet) & flax:	MCPA ^{◇◇} (0.2 to 0.4 L)*
	<i>Bromoxynil</i> / MCPA* - <i>Buctril M</i> (0.2 to 0.4 L), <i>Logic M</i> (0.25 to 0.5 L)
Preseeding corn (field only)	<i>Banvel II</i> (0.12 L)*
Preseeding field pea, lentil† & chickpea†:	MCPA Amine ^{◇◇} (0.2 to 0.28 L)* [◇]
Preseeding canaryseed & seedling forage grasses ^{◇◇◇} :	<i>Bromoxynil</i> / MCPA* - <i>Buctril M</i> (0.2 to 0.4 L), <i>Logic M</i> (0.25 to 0.5 L)
Chem fallow:	2,4-D [◇] (0.5 L)*
	<i>Dicamba</i> (0.12 L)*
	<i>Bromoxynil - Pardner</i> (0.51 L), <i>Koril</i> (0.48), <i>Brotex</i> (0.6)
Canada thistle control in stubble or fallow:	<i>Dicamba</i> (0.51 L)**
Alfalfa control in spring / fall:	2,4-D (0.50 to 1.0 L)*

* Volunteer glyphosate tolerant canola control: Tank mixes of 2,4-D at 0.23 to 0.34 L per acre, MCPA and *Bromoxynil* / MCPA will control volunteer glyphosate tolerant canola up to the 4 leaf stage and 2,4-D at 0.45 to 0.68 L per acre will give control up to the 6 leaf stage. Earlier application will result in more consistent control. *Dicamba* or *Banvel II* at 0.12 L per acre will not control glyphosate tolerant canola.

** See re-cropping restrictions for *Dicamba* with fall applications.

*** 2,4-D tank-mixes in cereals are registered for winter wheat, wheat, barley, and rye; *Bromoxynil* tank-mixes in cereals are registered on wheat, oats and barley; *bromoxynil* / MCPA and MCPA tank-mixes registered on cereals include wheat, barley oat and rye; *Banvel II* tank-mixes in wheat, barley, rye oats.

† Under drought conditions, deep seeding and/or brief rain showers after seeding may cause injury to emerging seedlings in sprayer overlaps. NOT for use with *Cheminova Glyphosate*, *Credit 45*, *Glyphos*, *Knockout Extra*, *Luffi Plus*, *MPower Glyphosate*, *NuGlo*, *SharpShooter*, *SharpShooter Plus*, or *Wise Up*.

†† *Roundup WeatherMax* only.

◇ Rates based on 500 g/L formulations. All formulation concentrations are registered unless indicated otherwise.

◇◇ Use only amine formulations of MCPA prior to corn, lentil, chickpea and field peas.

◇◇◇ Forage grasses include brome grass, crested wheatgrass, intermediate wheat grass, slender wheatgrass, tall wheatgrass, Russian wildrye, timothy, orchard grass, creeping red fescue, meadow fescue, meadow foxtail, tall fescue, meadow brome grass, streambank wheatgrass and reed canarygrass.

Tank mixes in glyphosate tolerant crops:

Tank mixes or rates listed may not occur on all product labels. Refer to individual product labels for registered tank-mixes.

Canola: Lontrel at 112 mL/acre.

Soybean: Assure II at 101 to 154 mL/acre, Pursuit at 65 to 85 mL/acre.

Corn: Axtrex at 0.63 to 0.84 L/acre, 2,4-D single application (500 g ai/L formulation) (0.22 to 0.45 L/acre)*, 2,4-D split application (500 g ai/L formulation) (0.22 L/acre followed by 0.17 to 0.22 L/acre)*, DyVel DSp (0.44 L/acre).

* 2,4-D applications to corn may result in serious injury to some corn hybrids. Consult corn seed provider for varietal tolerance to 2,4-D applications. Apply prior to 4 leaf stage of corn.

Note: The above mixes are those listed on the glyphosate labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. See the general guidelines for mixing pesticides for more information.

Application Information:

Water Volume:

Ground: Use 20 to 40 L per acre in most situations; use of the lower volume may improve control when hard water (Ca or Mg) or iron (Fe) ions are present (See Effects of Growing Conditions below). For certain crop situations, perennial weeds and tank mixes may require up to 120 L per acre of clean low ion water.

Aerial: Use 8.1 to 20 L per acre for registered preharvest uses only (see Aerial Application below). Minimum 20 L per acre for preseed, fallow, glyphosate tolerant crops and post-harvest treatments with Roundup WeatherMax only.

Refer to specific weed control situations or labels for more information on water volumes and adjuvants.

Nozzles and Pressure: Use 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets for ground applications and ASABE coarse droplets for aerial applications.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Best results are achieved when temperatures are near 20°C and when weeds are actively growing. Frost which kills more than 40% of the above ground tissue will reduce control. Control will also be reduced if foliage is heavily covered with dust. "Hard water" or water containing Calcium (Ca), magnesium (Mg) or iron (Fe) ions will reduce the activity of glyphosate products proportional to the level hardness. Reducing application water volume and /or adding ammonium sulphate at

1.2 kg/acre (99% dry) or 2.4 L/acre (49% solution) will reduce the negative effects of low levels of hard water ions. If water is extremely hard (greater than 700 ppm or 40 grains), another water source should be found. Dirty water or water with suspended soil or organic matter will reduce control.

Restrictions:

Rainfall: DO NOT apply if rainfall is forecast for the time of application, as weed control may be reduced. Consult manufacturer for more information.

Grazing Interval: All portions of forage and crops treated with glyphosate products may be fed to livestock.

Re-cropping Interval: No restrictions.

Aerial application: DO NOT apply Cheminova Glyphosate, Credit 45, Crush'R Plus, Knockout Extra, Lajj Plus, Maverick III, Matrix, NuGlo, Sharpshooter, Vector or WiseUp brands of glyphosate to cropland by air. All other glyphosate products listed in the "Product names, Company, Formulation and Packaging" chart are registered for aerial application for certain pre-harvest treatments. Not all crop species listed in the pre-harvest section are registered for aerial glyphosate application. Consult manufacturer for current aerial pre-harvest registration status.

ONLY Roundup WeatherMax may be applied by air when fields are too wet to access by ground sprayer (flooded) for preseed burndown, fallow treatment, or application to glyphosate tolerant crops (canola, corn, soybean).

Storage: May be stored below 0° C.

Equipment: DO NOT mix, store or apply this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.

Buffer Zones:

Application method	Uses	Buffer Zones (metres ^{††}) Required for the Protection of:	
		Aquatic habitats	Terrestrial habitat
Ground *	All uses	15	15
Aerial	Preharvest only**	25	55
	Preharvest only***	100	100
	Glyphosate tolerant canola only [†]	5	40
	Preseed, fallow, glyphosate tolerant crops (corn, soybeans) [†]	30	70

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

** ClearOut 41 Plus, Roundup Transorb IIC, Roundup Ultra

2, Roundup WeatherMax, R/T 540, StartUp, Touchdown Total, Traxion only.

*** Glyfos, MPower Glyphosate, SharpShooter Plus, Vantage Plus Max II only.

† Roundup WeatherMax only when conditions are too wet for access by ground sprayer.

†† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Glyphosate is very toxic to non-target plants.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

Roundup Transorb HC, Roundup Ultra 2, Roundup WeatherMax, R/T 540:



Caution - Poison

ClearOut 41 Plus, Cheminova Glyphosate, Glyphos, KnockOut Extra, Roundup TransorbHC, Roundup Ultra 2, Roundup WeatherMax, R/T 540, Sharpshooter Plus:



Warning - Eye and Skin Irritant

All other products:



Caution - Skin and Eye Irritant. Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Glyphosate/dicamba

Herbicide Group

4 - dicamba

9 - glyphosate

(Refer to page 38)

Company:

Monsanto Canada (Rustler)

Nufarm Agriculture (Glykamba)

Formulation:

Rustler (PCP# 27200; Glykamba (PCP# 30870)): 194 ae/L glyphosate and 46 g/L dicamba present as isopropylamine (IPA) salts formulated as a solution.

Container sizes - 10 L, 115 L, 450 L, 750 L.

Crops and Staging:

Summerfallow.

Pre-seeding on fields to be sown to wheat, barley, oats and rye.

May also be applied prior to sowing field corn in fields with more than 2.5% organic matter (DO NOT use on sandy or sandy loam soils).

Glyphosate/dicamba **SHOULD NOT** be applied prior to broadleaf crops such as lentils, peas, canola and flax due to the risk of injury.

Weeds, Rates and Staging:

Application should be made to emerged, actively growing weeds. Application at early growth stages generally provides the best results.

Annual grasses - Apply 1 L per acre between emergence and heading.

Downy brome

Green foxtail

Persian darnel

Volunteer cereals

Wild oats

Annual broadleaves - Apply 1 L per acre up to 6 inches (15 cm) height unless otherwise indicated.

Cow cockle

Flixweed

Kochia

Lamb's-quarters

Redroot pigweed

Russian thistle

Smartweed (including lady's-thumb)

Stinkweed

Volunteer canola*

Wild buckwheat (1 to 4 leaf)

Wild mustard

* NOT including glyphosate tolerant varieties.

Foxtail barley suppression - Apply 1.26 L per acre before initiation of the seedhead or bottom leaves beginning to brown off.

Application Information:

Water Volume: 20 to 40 L per acre water. Avoid the use of extremely hard water (greater than 700 ppm calcium and/or magnesium or high levels of iron). Use of the lower water volume may improve control in situations where hard water is the only source available.

Nozzles and Pressure: Use 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE medium to coarse* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Reduced effectiveness may result if application is made to weeds that are drought-stressed, damaged by disease or insects. Poor control under cool, cloudy weather can occur. Dust on foliage can also cause reduction in control.

Tank Mixes:

Herbicides:

In wheat, winter wheat, barley and rye only:

2,4-D Ester or Amine (0.2 to 0.3 L/acre)^{†*} or (0.4 to 0.5 L/acre)^{†**}

* to control volunteer glyphosate tolerant canola up to 4 leaf stage

** to control volunteer glyphosate tolerant canola up to 6 leaf stage.

† Rates listed based on 2,4-D 600 formulations. Multiply rates by 1.2 for 500 formulations and 0.85 for 700 formulations.

Restrictions:

Rainfall: Within 6 hours may reduce weed control. Heavy rainfall within 2 hours of application may require a repeat treatment.

Re-entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT allow lactating dairy animals to graze within 7 days of treatment or cut for feed or hay within 30 days. Remove meat animals from treated areas at least 3 days prior to slaughter.

Re-cropping: No restrictions in the season following treatment. DO NOT apply in fall or spring prior to broadleaf crops such as lentils, peas, canola and flax due to the risk of injury.

Aerial Application: DO NOT apply by air.

Storage: Store above 5°C.

Equipment: DO NOT mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks.

Buffer Zones:

Product	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
<i>Rustler, Glykamba</i> (medium droplets)	1	1	15
<i>Rustler, Glykamba</i> (coarse droplets)	1	1	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

Rustler, Glykamba:



Caution – Poison



Danger – Corrosive to eyes.



Warning – Skin Irritant
Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Good Harvest

Herbicide Group
10 - glufosinate
 (Refer to page 38)

Company:

Farmers of North America (PCP# 30761)

Formulation:

150 g/L glufosinate ammonium formulated as a solution.
 Container sizes: 13.5 L, 108 L, 432 L.

Crops, Rates and Staging:

Harvest Aid Treatment:

CROP	RATE (L/acre)	STAGE
Alfalfa (seed production only)	0.81 to 1.09**	50 to 75% pod turn (brown)
Lentil*	1.09	40 to 60% pod turn (yellow to brown)
Potato*	1.21	14 to 21 days prior to harvest

* Not for crops grown for seed.

** use the higher rate when crop canopies or weed densities are heavy.

Application Information:

Water Volume:

Ground applications: Minimum 45 L per acre. When crop canopy and weed densities are heavy, apply in 69 to 89 L per acre of water.

Aerial applications: 13 to 22 L per acre.

Nozzles and Pressure:

Ground Application: Use 40 psi (275 kPa) when using conventional 80° or 110° flat fan nozzles; 45 psi (310 kPa) when using check valves. Angle nozzles forward at 45° to improve coverage of vertical leaf surfaces. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium or larger droplets.

Aerial applications: DO NOT use raindrop nozzles. Use a combination of nozzles and pressure to provide ASABE coarse or larger droplet size distribution.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Good Harvest activity is influenced by environmental conditions. Cool temperatures (less than 10°C), drought, and low humidity conditions slow weed growth. Applications made under these stressed conditions may result in reduced weed effectiveness.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Within 4 hours may reduce activity.

Re-Entry: DO NOT re-enter treated areas for 24 hours after application, without protective clothing as for spraying.

Grazing: DO NOT graze the treated crop or cut for feed.

Preharvest Interval: Leave 9 days between application and harvest of lentil and potato.

Re-cropping: No restrictions.

Aerial Application: May be applied by air.

Storage: DO NOT freeze.

Buffer Zones:

Ground: DO NOT apply within 15 metres of sensitive plants or water or wetland areas.

Aerial: DO NOT apply within 30 metres of sensitive plants, or water or wetland areas.


DO NOT apply when dead calm or when winds exceed 16 km/hr when using unprotected booms or applying by air, or exceeding 25 km/hr when using shrouded booms.


Tank Cleaning:

Refer to 'Method B' in the general tank cleaning section on page 15 to 16.

Hazard Rating:

 Warning - Poison

 Caution - Skin Irritant

 Warning - Eye Irritant

For an explanation of the symbols used here see page 11.

Gramoxone

Herbicide Group
22 - paraquat
 (Refer to page 38)

Company:

Syngenta Canada (PCP#8661)

Formulation:

200 g/L paraquat formulated as a solution.
 Container size - 4 x 5 L

Crops and Staging:

Stale seedbed: Non-selective weed control applied 3 days prior to crop emergence in beans, corn, potatoes, peas, soybeans.

Non-selective inter-row weed control: Apply as a directed spray between rows in row crops. Avoid contact with crop foliage.

Control of weed seedlings in established alfalfa and bird's-foot trefoil for hay: Apply 5 days after first cut.

Control of weed seedlings in bird's-foot trefoil for seed: Apply in spring when bird's-foot trefoil shoots are 3 to 6 inches (7.5 to 15 cm) long.

Non-selective weed control - shelterbelts: Apply as a directed spray in and around shelterbelt trees or woodlot plantings. Avoid contact with foliage.

Pre or Post seeding burndown: Apply prior to crop emergence in barley, canary seed, canola, corn (field, sweet and pop), dry beans, field peas, flax (including low linolenic types), lentils, mustard, oats, potato, rye, soybean, sunflower, triticale, wheat.

Weeds and Staging:

Annual weed burn-off. Best control when weeds are less than 2 inches (5 cm) in height or diameter.

Rates:

Pre or post seeding burndown: 0.8 to 1.6 L per acre

All other applications: If weeds are less than 2 inches (5 cm) in height, apply 1.1 L per acre (5 L treats 4.5 acres). If weeds are taller than 2 inches (5 cm), increase the rate of Gramoxone to 2.2 L per acre (5 L treats 2.2 acres).

Application Information:

Water Volumes:

Pre or post seeding burndown: Minimum 40 liters per acre.
All other applications: 135 to 500 L per acre. Use the higher water volumes within the range if weed growth is dense. Good coverage is critical for good control.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASA/E medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Best results on cloudy days or just prior to darkness.

Tank Mixes:

Corn: AAtrex, Dual II Magnum, Frontier Max and Primextra II Magnum*

Soybean: Dual II Magnum, Frontier Max, Linuron, Sencor*

* Refer to product labels for time of application and restrictions.

Restrictions:

Rainfall: Within 1 hour will reduce weed control.

Re-entry Period: DO NOT re-enter treated fields for 24 hours following application. If necessary, workers may re-enter field after 4 hours if wearing protective clothing. See label for details.

Grazing: DO NOT graze or harvest treated foliage. Regrowth from treated alfalfa or bird's-foot trefoil may be fed to livestock.

Re-cropping Interval: No restrictions

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze.

Buffer Zones:

Crop	Buffer Zones (metres†) Required for the Protection of:			
	Aquatic Habitats of Depths			Terrestrial habitat
	Less than 1 m	1 m to 3 m	Greater than 1 m	
Conservation tillage soybean	40	30	20	15
All other crops	50	40	30	20

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

DO NOT spray when conditions are dead clam or when wind is gusty or blowing faster than 16 km/hr.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Danger - Poison (may be fatal if swallowed)



Danger - Corrosive to eyes

For an explanation of the symbols used here see page 11.

Grazon

Herbicide Group
4 - picloram & 2,4-D
(Refer to page 38)

Weed Control

Company:

Dow AgroSciences (PCP#27634)

Formulation:

65 g/L picloram and 240 g/L 2,4-D formulated as a solution.
 Container size - 2 x 10 L and 110 L (through select outlets).

Note: Available only through selected retail outlets.

Crops and Staging:

Permanent grass pasture and rangeland. Apply in spring or early summer.

Weeds, Rates and Staging:

Broadleaf weeds:

Apply at 1.5 L per acre: for season long control ONLY

Canada thistle	Dandelion
Common yarrow	

Apply at 2.8 L per acre: for control of the above weeds and the following weeds

Burdock	Goldenrod
Clovers (red, sweet)	Plantain
Common ragweed	Prickly lettuce
Dock	Vetch
Fleabane	Wild carrot

Application Information:

Water volume:

Ground application: 40 to 80 L per acre.

Aerial application: Minimum 8.1 L per acre.

Nozzles and Pressure: Use nozzles that will deliver coarse droplets in a uniform pattern. Maximum 30 psi (207kPa) by ground or air when using conventional flat fan nozzles.

Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets.

Drift of even small amounts of *Grazon* into sensitive plants or areas where sensitive crops may be grown can cause injury. DO NOT apply under conditions prone to drift (i.e. high winds, dead calm and temperature inversions).

How it Works:

Refer to Table 2 on page 40.

IMPORTANT: Picloram is a very persistent and water-soluble herbicide. Treated soil should not be moved from the treated area. DO NOT apply to soils that are permeable, have sinkholes, or lie over limestone bedrock. DO NOT apply to soils whose surfaces are composed of fractured rock or unconsolidated gravel. Application to these sites may allow the movement of herbicide to underlying water sources or aquifers. When applying *Grazon* over sandy soils ensure that aquifers are not within 1.8 m of the soil surface. If shallow aquifers are present, DO NOT apply *Grazon*. *Grazon* must not be applied on range and pasture acres that are irrigated. DO NOT compost or mulch clippings or manure from grass treated with *Grazon* unless being reapplied to the treated area.

Effects of Growing Conditions:

Nothing listed on the *Grazon* label. Avoid application when pasture and target weeds are under stress from drought, flooding, extreme heat or cold, as injury to grass or unacceptable control may result. Avoid application when temperatures exceed 28°C.

Tank Mixes:

None registered.

Restrictions:

Rainfall: DO NOT apply if rainfall is forecast. No specific time frame is indicated on the label. Contact manufacturer for more information.

Re-entry: DO NOT re-enter pastures within 3 days of application.

Grazing: DO NOT allow lactating dairy animals to graze treated areas within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter. DO NOT harvest forage or cut hay within 30 days of application. Feed livestock untreated forage for 7 days prior to moving onto land that produce broadleaf crops - otherwise urine or manure may contain picloram. See restrictions in "How it Works" section above.

Re-cropping: Legumes may not be established in a pasture for several years after treatment. If legumes are essential in a pasture, DO NOT use *Grazon*. DO NOT break up treated pasture and plant to sensitive broadleaf crops for at least 5 years after application of *Grazon*.

Aerial Application: May be applied by air.

Storage: Store product in original containers in a secure, dry, cool area. DO NOT freeze.

Buffer Zones:

See page 29 for an explanation of the different habitats.

Field Sprayer (Rangeland Uses)

Use Rate (L./acre)	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
1.5	1	1	30
2.8	2	1	55

* These distances can be reduced by 30% using cones on individual nozzles and by 70% using a full shield (shroud, curtain) that extends to the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Refer to the label for buffer zone requirements for aerial application.

Heavy rains can move this product from its application site down slope toward sensitive areas. DO NOT load or mix near wells, dugouts or other water bodies.

Tank Cleaning:

There are no tank cleaning recommendations on the product label. A combination of Method A and B found on the general page on tank cleaning on page 15 to 16, or the use of a commercial tank cleaner, completed immediately after application is finished may be the best cleanout option.

Hazard Rating:



Caution – Poison



May cause skin and eye irritation

For an explanation of the symbols used here see page 11.

Harmony K

Harmony K is equivalent to a tank mix of *thifensulfuron/tribenuron* (page 272), *clodinafop* (page 122) plus *dicamba* (page 126). For other detailed information on the component products see the product pages listed above.

Company:

E. I. duPont Canada

Formulation:

The *Harmony K* package contains the following components:

Harmony Broadleaf (PCP# 30027): 53.8% dicamba sodium salt, 7.7% thifensulfuron methyl, and 3.9% tribenuron methyl formulated as a water dispersible granule.

Container size - 2.104 kg

Harmony Grass (PCP# 29202): 128 g/ L clodinafop-propargyl formulated as an emulsifiable concentrate with built in adjuvant.

Container size - 1 x 7.1 L

Crops and Staging:

Spring wheat (including durum) from the 2 to 5 leaf stage.

Herbicide Group

- 1 - clodinafop
 - 2 - thifensulfuron & tribenuron
 - 4 - dicamba
- (Refer to page 38)

Weeds and Staging:

Harmony K controls the same weeds as *Harmony SG* with the addition of Group 2 resistant kochia and dandelion (spring or fall rosettes less than 15 cm in diameter).

Rates:

Harmony Grass: 177 mL per acre

-plus-

Harmony Broadleaf: 52 g per acre

One package treats 40 acres (16 ha)

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

See *thifensulfuron/tribenuron*, *clodinafop* and *dicamba*, pages on restrictions application details and handling. Use the most limiting restrictions across all components for the mix

Note: *Harmony K* may be applied by air.

Harmony Max*

Harmony Max is equivalent to *thifensulfuron/tribenuron* (page 272) tank mixed with *clodinafop* (page 122) and *Perimeter* (= Starane or 'A' component of *Attain*, *Prestige* and *Trophy*). For other detailed information on the component products see the product pages listed above.

Company:

E. I. duPont Canada

Formulation:

The *Harmony Max* package contains the following components:

Refine SG (PCP# 28285): 33.35% thifensulfuron methyl plus 16.65% tribenuron methyl formulated as a water soluble granule.

Container size - 486 g bottle.

Herbicide Group

- 1 - clodinafop
 - 2 - thifensulfuron & tribenuron
 - 4 - fluroxypyr
- (Refer to page 38)

Harmony Grass (PCP# 29202): 128 g/ L clodinafop-propargyl formulated as an emulsifiable concentrate with built in adjuvant.

Container size - 1 x 7.09 L

Perimeter (PCP# 29586): 180 g/L fluroxypyr formulated as an emulsifiable concentrate.

Container size - 4.86 L.

* NOTE: This product is no longer manufactured, but some supplies may still remain in the distribution system. This product may be removed from future editions of this publication.

Crops and Staging:

Spring wheat (including durum) from the 2 leaf stage up to the emergence of the 4th tiller.

Weeds and Staging:

Broadleaf weeds controlled by *Refine SG plus*;

Group 2 resistant kochia.

Cleavers (up to 4 whorls)

Wild oat - 1 to 6 leaves up to the emergence of the 4th tiller.

Green foxtail - 1 to 5 leaves up to the emergence of the 3rd tiller.

For optimum control apply before the annual grasses tiller.

Rates:

Refine SG: 12 g per acre

Harmony Grass: 177 mL per acre

Perimeter: 0.12 L per acre

One package treats 40 acres (16 ha) Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Re-Entry: DO NOT re-enter treated fields for 12 hours.

Preharvest Interval: Leave 60 days between application and harvest.

Grazing: MUST NOT be grazed or fed to livestock for 7 days after treatment.

Re-cropping: Barley, canola, flax, forage grasses, lentils, mustard, oats, peas, rye or wheat or fields can be summer-fallowed the year after treatment.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen. If frozen, bring to room temperature and agitate before use. This product is COMBUSTIBLE. DO NOT store near heat or open flame.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground*	15	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Danger - Poison



Warning - Eye and Skin Irritant; contains the allergen epoxidized soybean oil.

For an explanation of the symbols used here see page 11.

Harmony SG

This product is a prepackaged tank mix of *thifensulfuron/tribenuron* (page 272) and *clodinafop* (page 122). Information listed is restricted to Crop, Weeds, Rates and Tank mixes. For other detailed restrictions and other general information on the component products see the product pages listed above.

Company:

E. I. duPont Canada

Formulation:

The *Harmony SG* package contains the following components:
Refine SG (PCP#28285): 33.35% thifensulfuron methyl + 16.65% tribenuron methyl; formulated as a water soluble granule.

Container size - 486 g bottle.

Harmony Grass (PCP#29202): 128 g/ L clodinafop-propargyl formulated as an emulsifiable concentrate, with built in adjuvant.

Container size - 1 x 7.1 L

Crops and Staging:

Spring wheat (including durum) up to the emergence of the 4th tiller.

Weeds and Staging:

Broadleaf weeds controlled or suppressed by *Refine SG* plus:

Wild oat - 1 to 6 leaves up to the emergence of the 4th tiller.

Herbicide Group

1 - clodinafop

2 - thifensulfuron & tribenuron

(Refer to page 38)

Green foxtail - 1 to 5 leaves up to the emergence of the 3rd tiller.

For optimum control apply before the annual grasses tiller.

Rates:

Refine SG: 12 g per acre

Harmony Grass: 177 mL per acre

One case treats 40 acres (16 ha)

Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 14.

Tank Mixes:

Herbicides:

MCPA Ester (0.34 to 0.45 L/acre of 500 g/L formulation).

Banxel II (44.5 or 59 mL/acre).

See component products for more information on restrictions application details and handling. Use the most limiting restrictions across all components for the mix.

Heat

Company:

BASF Canada (PCP# 29368)

Formulation:

70 % saflufenacil formulated as a water soluble granule.

Container size - 8 x 844 g containers per case.

Herbicide Group

14 - saflufenacil

(Refer to page 38)

Crops, Rates and Staging:

Prior to the seeding of; or following seeding and prior to the emergence of the following crops; fallow or after harvest:

Must be applied as part of a tank mix with glyphosate from 180 to 360 g ae per acre (0.5 to 1 L per acre of 360 g per L forms):

CROP	RATE (g of product per acre)
Barley, canaryseed, chickpea, corn (field and sweet*), field pea, oat, wheat (including spring, winter and durum)	10.4 to 28.4
Lentil† soybean**	10.4
Summer-fallow and post-harvest	10.4 to 28.4

* Some varieties may be more sensitive to *Heat* and injury may occur

† DO NOT use rates higher than 10.4 g per acre or injury could result.

Add either *Merge* or *Amigo* adjuvant at 0.2 L per acre (One 844 g container treats 80 to 30 acres)

Harvest Aid/ Desiccation:

Apply 14.4 to 28.4 grams per acre plus *Merge* at 0.2 L per acre to speed the rate of dry-down of the following crops and green weedy material. The required delay before harvest of each crop is indicated below.

CROP	PRE-HARVEST INTERVAL	APPLICATION STAGE
Field Pea	3	A majority of the pods are brown (70 – 80%)
Red lentil varieties only	3	Lower most pods (15%) are brown and rattle when shaken
Dry bean	2	Stems are green to brown, pods are mature (yellow to brown), and 80 – 90% of leaves have dropped
Soybean	3	
Sunflower	7	The backs of flower heads and bracts are turning yellow, and seed moisture is 20 – 30%.

Heat may be tank mixed with glyphosate on field pea, lentil, dry beans and soybeans for additional pre-harvest weed control. DO NOT tank mix with glyphosate when the harvested grain is to be used for seed.

Note: Tolerances for Heat residue may not be established in all countries importing the above crops. Check with your grain buyer prior to application to determine if they will accept these crops treated with Heat.

Weeds, Rates and Staging:

Apply up to the 8 leaf stage unless otherwise indicated to control the weeds controlled by glyphosate plus rapid burndown of:

Canada fleabane	Pigweed (redroot)**
Cleavers (4 whorl-stage)	Round-leaved mallow
Dandelion	Stinkweed**
Kochia (up to 15 cm)	Volunteer canola**
Lamb's-quarters	Wild buckwheat **
Flixweed	Wild mustard**
Narrow-leaved hawk's-beard	

* All varieties

** Applications at the 28.4 g per acre rate will also provide suppression of the emergence of these weeds following application.

Application Information:

Water volume:

Preseed, summerfallow or post harvest by ground only: 20 to 40 L per acre.

Desiccation:

Ground: 81 L per acre

Aerial: 20 L per acre.

Higher volumes are required for dense weed stands. Weed control improves with the amount of coverage.

Nozzles and Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium* classification droplets. Low drift nozzles may require higher pressures for proper performance. Higher pressures may be required to penetrate dense weed stands.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Rainfall shortly after application can result in slight injury to the crop. See the 'Restrictions' section below for more details. Warm, moist growing conditions promote active weed growth. Weeds hardened off by environmental stress such as cold weather, drought or excessive heat may not be adequately controlled.

Tank Mixes:

Herbicides:

Preseed, summerfallow or post harvest:

Glyphosate (180 g to 360 g ae per acre)*

* must be mixed with glyphosate.

Desiccation/Harvest Aid:

Glyphosate (360 g ae per acre)†

† NOT for use on crops to be used for seed.

(see glyphosate page for product concentrations and equivalent application rates)

Fungicides: None registered

Insecticides: None registered

Fertilizers: None registered

Note: The above mixes are those listed on the *Heat* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall shortly after application can result in slight injury to the crop. Lentils are more sensitive to injury on coarse textured (sandy or gravelly) and low organic matter soils. Injury will appear as slight leaf edge burning, which will be grown out of and yield will not be affected.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Preharvest: Leave 60 days between treatment and harvest.

Grazing: DO NOT graze fields or feed straw from fields where *Heat* is applied as a harvest aid treatment.

Re-cropping:

CROP	APPLICATION RATE (per acre) AND TIMING		
	Spring Application		Pre-harvest Application
	10.4 g	up to 28.4g	up to 28.4g
Barley	PB	PB	1
Canary seed	PB	PB	1
Canola	1	1	1
Chickpea	PB	PB	1
Corn	PB	PB	1
Dry Bean	1	1	2
Flax	1	1	1
Lentil	PB	1	1
Mustard	1	1	1
Oat	1	1	1
Field Pea	PB	PB	1
Soybean	PB	1	1
Spring Wheat (including durum)	PB	PB	1
Winter wheat	PB	PB	1

PB May be planted back in the same season

1 May only be planted the season following application

2 May only be planted the second season following application

Aerial Application: May be applied by aircraft for desiccation use only. DO NOT apply by air for any other use.

Storage: Store in dry, cool storage. May be frozen.

Buffer Zones:

Application method	Crop	Buffer Zones (metres†) Required for the Protection of Terrestrial Habitat
Ground only*	Lentil, Soybean	3
	All other crops	10
Fixed wing airplane	All desiccation uses	175
Helicopter	All desiccation uses	150

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as meters from the downwind edge of the spray boom to sensitive habitat.

DO NOT apply in areas where surface water from the treated area can run off to adjacent cropland, streams

Tank Cleaning:

HEAT can cause injury to sensitive crops at very low concentrations. Sprayers used to spray this product should be flushed out immediately after use.

Refer to 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

 Caution – Possible Skin Irritant

For an explanation of the symbols used here see page 11.

Imazamethabenz

Herbicide Group
2 - imazamethabenz
 (Refer to page 38)

Company:

Nufarm Agriculture (Assert 300SC - PCP# 21032)
 Loveland Products (formerly Viterra) (Avert - PCP#29618)

Formulation:

300 g/L imazamethabenz formulated as a suspension concentrate.

Container size - 2 x 10.8 L jugs per case.

pH adjuster: 94.5% sodium bisulfate formulated as a soluble granule.

Container size - 2 x 2.5 kg bags (1 bag per 10.8 L jug of *imazamethabenz*).

Crops, Rates and Staging:

pH adjuster: 1 packet per jug of Assert to be used.

Imazamethabenz up to 0.67 L per acre (16.1 acres per jug):
 Barley, spring wheat (including durum) - 1 to 6 leaf stage.
 Annual ryegrass (seed production only) - 4 to 6 leaf stage.

Imazamethabenz at 0.34 L per acre (32 acres per jug):
 Sunflower - plants (not under drought stress) that are in the 2 to 8 leaf stage. Crop must be less than 15 inches (38 cm) tall except for semi-dwarf varieties, which must be less than 12 inches (30 cm), and dwarf varieties, which must be less than 4 inches (10 cm). Stunting and head deformation can occur from applications made beyond recommended stages.

DO NOT apply *imazamethabenz* to the same field more than once in two years.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions. Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

Weeds, Rates and Staging:

WEEDS	STAGE	RATE	
		(L per acre)	Acres per 10.8 L jug
Stinkweed Wild mustard	Up to 6 leaves	0.34	32
<i>Weeds above plus:</i> Buckwheat (wild and tartary) (suppression) Volunteer canola (except Clearfield varieties)	Up to 4 leaves	0.54	20
Wild oats	1 to 3 leaves		
Wild oats	1 to 4 leaves	0.67	16

* Main stem leaves.

Application Information:

Good coverage of foliage is important to maintain good control

Water volume: In cereals only, *imazamethabenz* may be applied in 20 to 40 L of water per acre when applied alone or when tank mixed with *dichloroprop/2,4-D*, *2,4-D* ester, or MCPA ester. For all other applications, apply in 40 L per acre.

Nozzles and Pressure: 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets or larger.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply *imazamethabenz* 24 hours before or after a frost. It works best at warm temperatures. Performs relatively consistently under dry conditions. If cold, wet soil conditions persist in the days after application, retillering of wild oats may occur. DO NOT apply to drought stressed sunflowers.

Tank Mixes:

Herbicides:

Imazamethabenz may be applied at either 0.53 L or 0.67 L per acre in tank mixes in the brown and dark brown soils, but must be applied at 0.67 L per acre when tank mixing in the black and grey wooded soils for adequate wild oat control.

In spring wheat (including durum) and barley:

2,4-D Ester* (up to 0.32 L/acre) (700 g/L formulations)

Curtail M (0.80 L/acre)

Dichlorprop/2,4-D ester (0.7 L/acre)*†

Fenoxaprop** (0.118 L/acre)

Fenoxaprop** (0.118 L/acre) + MCPA Ester, (0.28 L/acre)

Fenoxaprop** (0.118 L/acre) + Refine SG (12 g/acre)

Frontline XL (0.65 L/acre)

Infinity (0.33 L/acre)

MCPA Ester* (up to 0.38 L/acre) (600 g/L formulations)

Refine SG (12 g/acre)

Refine SG (12 g/acre) + MCPA Ester, (0.28 L/acre)

Spectrum (20 acres per case)

Trophy (20 acres per case)

† Mix with Turboprop is NOT for use in barley. Dichlorprop-D registered with Assent only.

* Apply in 20 to 40 L of water per acre. For all other tank mixes use 40 L/acre.

** Use the 0.54 L/acre rate of *imazamethabenz* when tank mixing with *fenoxaprop* (see product labels for specific products). *Fenoxaprop* rate provides green foxtail control only.

*** When tank mixing dry broadleaf products, add products to the tank in the following order: dry broadleaf products, acidifier, *imazamethabenz*, and other liquid herbicides if required. For repeat tanks, dry broadleaf products need to be mixed with water to form a slurry prior to adding to the remaining spray solution in the tank.

Refer to *imazamethabenz* labels for specific mixing order and application details when tank-mixing.

Refer to tank mix partner for additional crop staging restrictions.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *imazamethabenz* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours will reduce control.

Re-entry: Wait at least 12 hours before entering treated fields.

Grazing: DO NOT graze treated fields or cut treated forage for silage or hay. Mature barley and wheat grain or straw from fields treated with *imazamethabenz* can be fed to livestock. DO NOT feed or graze treated annual ryegrass.

Preharvest Interval: DO NOT apply beyond the recommended crop stage.

Re-cropping:

DO NOT apply *imazamethabenz* to the same field more than once in two years.

Year After Application	Black and Grey Wooded Soils	Brown and Dark Brown Soils
Year 1	Spring wheat (including durum), barley, canola, field peas, flax, sunflowers	Spring wheat (including durum), CLEARFIELD canola, barley, sunflowers
Year 2	Spring wheat (including durum), barley, canaryseed, canola, field peas, flax, oats, sunflowers	

Conduct a field bioassay (a test strip grown to maturity) the year before growing any crop not listed in the table. Lentils are known to be particularly sensitive to *imazamethabenz* residues in the soil. The additive effect of soil residues from the use of *imazamethabenz* and sequential applications of *imazethapyr*, *metsulfuron*, or *Odyssey* herbicides on the same land area has not been determined. Crop rotation guidelines are not known and injury to rotational crops other than wheat (excluding durum) may occur. Plant only wheat (excluding durum) on fields where these herbicides have been used until a field bioassay demonstrates other crops can be grown successfully.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Shake well before using.

Buffer Zones:

CROP	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Sunflower	0	0	1
Annual ryegrass, Cereals	1	0	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

Refer to page 15.

Hazard Rating:



Caution – Poison



Warning – Eye Irritant

For an explanation of the symbols used here see page 11.

Imazethapyr

Herbicide Group
2 - imazethapyr
 (Refer to page 38)

Company:

BASF Canada (*Pursuit* - PCP# 23844)

Farmers of North America (*MPower Kamikaze* - PCP# 30127)

MANA Canada (*Phantom* - PCP# 30017)

Univar (*Gladiator* - PCP# 28923)

Loveland Products (formerly Viterra) (*MultiStar* - PCP# 29259)

Formulation:

240 g/L imazethapyr formulated as a liquid.

Container size - 2 x 3.3 L jugs per case.

Crops and Staging:

All products: DO NOT use in the brown or dark brown soil zones (except for use in dry bean and alfalfa under irrigated brown soils); rotational crops may be severely injured due to carry over in these soils.

CROP	STAGE
Field pea	May be applied up to the sixth above-ground node stage (6 true leaves).

Pursuit, Gladiator, MultiStar and Phantom only:

CROP	STAGE
Dry bean (pinto, pink and red varieties only)	Up to and including the second trifoliate leaf stage
Soybean (Manitoba only)	Up to and including the third trifoliate leaf stage
Seedling alfalfa (forage or seed production)*	Apply after the first trifoliate leaf stage.

CROP	STAGE
Established alfalfa (seed production only)**	Apply before alfalfa reaches 12 inches (30 cm) in height.
Chickling vetch grown for seed	Apply at the 5 to 7 leaf stage.

* Apply only to seedling alfalfa that will remain in production for at least 3 years following application.

Apply only once during the life of the alfalfa stand.

** DO NOT apply in the last year of established alfalfa stands.

Weeds and Staging:

In field peas. Apply up to the 4 leaf stage, unless otherwise indicated:

Chickweed	Stinkweed
Cleavers	Volunteer canola (not CLEARFIELD varieties)
Green foxtail	Wild buckwheat†
Hemp-nettle	Wild mustard
Redroot pigweed	Wild oats † (2 to 4 leaf stage)
Shepherd's-purse	
Smartweed	

In seedling and established alfalfa:

Common groundsel**	Stinkweed
Green foxtail†	Volunteer canola (not CLEARFIELD varieties)
Green smartweed *	Wild mustard
Redroot pigweed	
Shepherd's-purse**	

In dry bean:

Hairy nightshade (up to 6 leaf stage)

* Seedling alfalfa only.

† Suppression

Rates:

85 mL per acre (40 acres per jug).

A non-ionic surfactant with at least 80% active ingredient (Agral 90, Aqsurf II, Surf 92) should be added at a rate of 0.25 L per 100 L of spray solution. DO NOT over apply imazethapyr, as crop injury may result.

DO NOT apply imazethapyr more than once per season or follow imazethapyr with other products containing imazethapyr in the same year.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 40 to 160 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver proper coverage with ASABE medium droplets or larger.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT spray if temperatures of +5°C are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

None registered.

Note: There are no tank mixes listed on the imazethapyr labels. To check for other possible registered tank mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT graze or harvest seedling alfalfa within 14 days of treatment. DO NOT graze or harvest field peas for feed within 30 days. DO NOT graze other treated crops or cut for feed prior to crop maturity.

Preharvest Interval: DO NOT apply within 60 days of harvesting field peas or chickling vetch, within 75 days of harvesting dry beans, or within 85 days of harvesting soybeans.

Re-cropping: Rotate to barley, spring wheat (not durum), lentils, alfalfa, field pea or CLEARFIELD canola the year following application. The manufacturer recommends that a field bioassay (a test strip grown to maturity) be conducted the year before growing any crop other than those listed above. However, yield losses within the test strips may not be noticed unless the yield can be compared to an untreated area seeded adjacent to the imazethapyr-treated strip. In case of crop failure, only field peas or CLEARFIELD canola may be replanted in the year of application.

NOTE: Breakdown of imazethapyr may be slowed or delayed by environmental conditions such as drought, excessive cold and/or acid soils (pH less than 6.5) resulting in an increased risk of injury to rotational crops. The most tolerant crops are CLEARFIELD canola and legume crops, then cereals. Contact manufacturer for additional information on re-cropping intervals (1-877-3712273).

Aerial application: DO NOT apply by air.

Storage: DO NOT freeze. If the product is exposed to temperatures below 0°C, thaw the product completely and shake the container vigorously prior to use.

Buffer Zones:

Pursuit, MultiStar, Gladiator:

Application method	Buffer Zones (metres†) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground only*	1	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured is metres from the downwind edge of the spray boom to sensitive habitat.

Other Products: DO NOT apply within 15 m of shelter-belts, water bodies, wetlands, and woodlots.

Tank Cleaning:

There are no specific tank cleaning directions on the product label. The use of 'Method C' in the general section on tank cleaning on pages 15 to 16 is recommended for other products with similar chemistry. Contact the manufacturer for more information.

Hazard Rating:

▽ Caution – May cause skin irritation

▽ Caution – May cause eye damage

For an explanation of the symbols used here see page 11.

Impact

Herbicide Group

27 – topramezone

(Refer to page 38)

Company:

AmVac Corporation, distributed in Canada by UAP
(PCP# 28141)

Formulation:

336 g/L topramezone formulated as a suspension.
Container size - 8 L

Crops and Staging:

Corn (field, sweet[†]): From the 1 to 7 leaf stage

[†] NOTE: Tolerance of sweet corn varieties to Impact and its mix partners may be variable. When tolerance is unknown, check with the supplier of seed and/or apply to a small area first to assess tolerance.

Weeds and Staging:

The following weeds are controlled with Impact unless otherwise indicated:

Impact MUST BE applied in tank mix with one of the herbicide options indicated in 'Tank Mixes':.

Grass weeds below from the 1 to 4 leaf stage:

Barnyard grass*	Foxtail (green and yellow)*
-----------------	-----------------------------

Broadleaf weeds below from the 1 to 8 leaf stage:

Chickweed (common)*	Pigweed (redroot, green)*
Lamb's-quarters*	Ragweed (common)
Lady's-thumb*	Velvetleaf*
Nightshade (eastern black)	Wild mustard

* Suppression only

Rates:

15 mL per acre

Must be applied with either:

Merge adjuvant at 0.5 L per 100 L of spray solution

-or-

Assist at 1.25 L per 100 L plus UAN (liquid 28-0-0) at 1.25 L per 100 L of spray solution.

Maximum one application of Impact per season.

Application Information:

Water Volume: 81 L per acre.

Nozzles and Pressure: Use 20 to 40 psi (140-276 kPa) if applying without drift reduction nozzles. Drift reduction nozzles may require higher pressures for proper performance. Select the nozzle and pressure combination that produces of ASABE medium droplets while maintaining good coverage of foliage.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

When weeds are stressed because of drought, flooding, hot or cool temperatures, weeds are not actively growing, control may be reduced.

Tank Mixes:

Herbicides:

Impact must be mixed with one of the following:

Field and Sweet Corn:

AAtrex (0.42 L/acre)

Field corn only:

Frontier Max (0.3 L/acre) + AAtrex (rates above) (DO NOT use Merge adjuvant with this mix).

Glyphosate tolerant corn only:

Glyphosate – 360 gai per acre, no adjuvant required (see glyphosate page for details)

Glyphosate + AAtrex (rates above)

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the Impact label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: DO NOT apply if heavy rain is forecast. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated fields or cut for feed within 45 days of application.

Pre-harvest Interval: Leave 45 days between application and harvest.

Re-cropping: Winter wheat may be seeded a minimum four months after application. Spring wheat, field corn, navy (white) bean, soybean, pea and alfalfa may be seeded the following crop year. Check tank mix options for additional reseeding restrictions. Conduct a field bioassay (a test strip grown to maturity) the year before growing any other crop.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool (above 5°C), dry area. If product is frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	1	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds

and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Spray when winds are under 16 km/hr, but not dead calm.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on pages 15 to 16.

Hazard Rating:



Warning – Contains the allergen soy.

For an explanation of the symbols used here see page 11.

Inferno Duo

Herbicide Group

2 - flucarbazone, tribenuron
(Refer to page 38)

Company:

Arysta LifeScience Canada (PCP#30663)

Formulation:

45% flucarbazone and 25% tribenuron formulated as a water dispersible granules.

Container size - 4 x 254.5 gram pouches.

Crops and Staging:

Spring wheat (NOT including durum):

Apply to the soil surface from one week before seeding until crop emergence.

Weeds, Rates and Staging:

Apply 12.75 ml per acre of *Inferno Duo* (one 254.5 g pouch treats 20 acres) plus 180 gae per acre of IPA or K+ salts (see glyphosate page) to control:

DO NOT apply more than 6.1 g active flucarbazone per acre (equivalent of 13.5 g/acre of *Inferno Duo* or 15.4 mL/acre of flucarbazone 2.0 formulations) per growing season.

Weeds controlled by glyphosate at 180 gae/acre (see glyphosate page) plus:

Cow cockle

Dandelion†

Narrow-leaved hawk's-beard

Shepherd's-purse

† Suppression only.

Volunteer canola***

Wild oats†

Plus weeds controlled by glyphosate at 180 gae/acre (see glyphosate page)

NOTE: The entire 254.5 g pouch must be added to the spray tank. DO NOT use part pouches.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Use 30 to 50 psi (200 to 345 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop tolerance and weed control may be reduced if applications are made to plants growing under stress. Stress includes saturated or water-logged soil, drought, extreme temperatures, low fertility or visible disease symptoms at application. Adopting practices to increase crop vigor will improve crop tolerance.

Tank Mixes:

Herbicides:

Glyphosate IPA or K+ salts only.

Fertilizers: None.

Insecticides: None

Note: The above mixes are those listed on the *Inferno Duo* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: DO NOT apply if rainfall is expected within 1 hour of application.

Grazing: DO NOT graze treated fields. Mature grain or straw may be fed to livestock.

Preharvest: Leave at least 80 days from application to harvest.

Re-cropping Interval: The following crops may be planted 11 months after application.

Soil Zones and Rotational Crops			
Grey-Wooded	Black	Dark Brown	Brown
Spring Wheat Barley Canola (all varieties) Field Pea*	Wheat (Spring & durum) Barley Canola (all varieties) Field Pea* Flax Field Bean	Wheat (Spring & durum) Barley Canola (all varieties) Field Pea* Flax	Spring Wheat

* Field peas may be grown the year following application in fields where precipitation has been equal to or above the 10 year average during the growing season, and where organic matter content is above 4%, and pH is below 7.5. The company suggests a minimum of 100 mm (4 inches) of rain is needed in the 60 days following application for adequate breakdown to take place.

NOTE: Other rotational crops may also be affected if rainfall is less than the 10 year average for the area. Soils in the grey wooded, black and dark brown soil zones with a combination of low organic matter (less than 2%), light textured soils or high pH (greater than 7.5) (i.e. eroded knolls, sandy soils) may result in delayed growth and development in rotational crops. Do not plant crops other than those listed above in the year following application.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones: Leave at least 20 m from the downwind edge of the spray swath to sensitive upland plants like shelterbelts and woodlots and at least 35 m to water sources or wetland habitats. Avoid drift onto sensitive crops like canola and tame oat. DO NOT mix or load within 10 m of water sources or wetland habitats.

Tank Cleaning:

Inferno Duo residues in the spray tank can cause severe injury to sensitive crops at very low concentrations. Sprayers should be cleaned out immediately before using another product.

Refer to 'Method A' in the general section on tank cleaning on pages 15 to 16.

When mixing with other pesticides, combine the method above with the method required for the tank mix partner if it is different from above.

Hazard Rating:

⚠ Warning: Contains the allergen milk and sulphites.

⚠ Caution – Skin irritant.

For an explanation of the symbols used here see page 11.

Infinity

Herbicide Group
6 - bromoxynil
27 - pyrasulfotole
(Refer to page 38)

Company:

Bayer CropScience (PCP# 28738)

Formulation:

37.5 g/L pyrasulfotole and 210 g/L bromoxynil formulated as an emulsifiable concentrate.

Container size - 2 x 6.7 L jugs per case.

Crops and Staging:

The following crops may be treated when at the 1 leaf stage of growth until the flag leaf is just visible but still rolled:

Barley	Timothy (seed production only)
Perennial ryegrass * (seedling & established, grown for seed or forage)	Triticale
Red fescue and brome grass * (established, grown for seed or forage)	Wheat (spring, durum, winter)

* NOTE: Since the uses on forage grasses were registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Those who apply this use do so at their own risk.

Weeds, Rates and Staging:

At 0.33 L per acre (one case treats 40 acres) the following weeds are controlled at the 1 to 6 leaf stage unless otherwise noted:

Annual sow-thistle	Pale smartweed
Chickweed	Perennial sow-thistle*
Canada Thistle* (up to 30 cm)	Redroot pigweed
Cleavers (1 to 3 whorls)	Round-leaved mallow*
Cleavers (4 to 6 whorls)**	Russian thistle (up to 10 cm)
Common Ragweed	Shepherd's-purse
Dandelion* (up to 25 cm across†)	Stinkweed
Flixweed (up to 10 cm)	Volunteer canola***
Hemp-nettle	Wild buckwheat
Kochia (up to 10 cm)	Wild mustard
Lamb's-quarters	

* Suppression only

** To control cleavers from 4 to 6 whorls add ammonium

sulphate at 202 g/acre (99% dry)
 or 0.4 L/acre (49% solution).

If using an ammonium sulphate product with a different concentration, adjust the rate accordingly.

*** All herbicide tolerant varieties.

† Spring seedlings and overwintered rosettes.

DO NOT apply *Infinity* or other products containing pyrasulfotole or bromoxynil more than once in the same year.

Application Information:

Water Volume:

Ground: Minimum 19 L per acre

Aerial: Minimum 11.4 L per acre

Higher water volumes should be used under dense crop and weed canopies to ensure thorough coverage of the target weeds.

Nozzles and Pressure: Maximum 40 to 45 psi (275 to 310 kPa) with conventional flat fan nozzles. Use nozzles and pressure designed to deliver proper coverage with *ASABE* medium droplets. Angle ground sprayer nozzles forward at a 45° angle to improve coverage of vertical leaf targets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

For best results, apply to emerged, young, actively growing weeds according to the weed stages listed. Under stressed conditions and/or heavy crop canopy, early application will result in improved weed control. Weeds growing under adverse environmental conditions such as drought will be less susceptible herbicide effects

Tank Mixes:

Herbicides:

Wheat (spring, winter, and durum), barley and triticale:
Liquid Achieve (0.2 L/acre) plus *Turbocharge* adjuvant

Wheat (spring and durum) and barley only:
Puma Advance (206 to 412 mL per acre)

Wheat (spring, NOT durum) and barley only:
Axial 100EC (0.28 L/acre) plus *Adigor* adjuvant

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: DO NOT mix with fertilizers other than those indicated above.

Note: The above mixes are those listed on the *Infinity* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Bayer also supports the following mixes that are not on the *Infinity* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: 2,4-D Ester (56-112 g ai/ac), Axial+Tilt, Horizon NG, Lontrel, MCPA Ester (56-112 g ae/acre), Puma Advance + Tilt, Traxos, Varro, Traxos+Tilt.

Fungicides: Tilt.

Insecticides: Decis, Sevin XLR.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Re-entry: DO NOT re-enter treated area within 12 hours.

Grazing: DO NOT graze treated crops or cut for hay within 25 days of application.

Preharvest Interval: Leave at least 50 days for wheat and triticale and 45 days for barley from application to harvest of grain or straw.

Re-cropping: Alfalfa, barley, canaryseed, canola, field corn (Manitoba only), flax, potatoes, soybeans (Manitoba only), sunflowers, tame oat, and wheat (durum, spring) may be seeded the year following application. Field peas may be grown the season following application in black, grey-wooded and dark brown soil zones. DO NOT plant field peas the season following *Infinity* use in the brown soil zone where organic matter content is below 2.5 % and where soil pH is above 7.5. Lentils may be seeded the second season after application.

Aerial Application: May be applied by air.

Storage: Store product in original containers in a secure, dry area, away from other pesticides, food or feed above -20°C. If stored over winter, shake or mix well before using.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	1	5
Fixed wing airplane	10	1	375
Helicopter	10	1	225

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

The manufacturer recommends a cleanout process similar to "Method A" on page 15 using a combination of water and ammonia solution rinses.

For additional information, Refer to page 15.

Hazard Rating:



Warning – Warning Poison



Warning – Eye and Skin Irritant.



Warning – Contains the allergen soy.

For an explanation of the symbols used here see page 11.

Kerb

**Herbicide Group
15 - propyzamide**
(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

Kerb 50WP (PCP#25595): 50% propyzamide formulated as a wettable powder. Container size - 1.36 kg (3 x 454 g water soluble pouches).

Kerb SC (PCP#30264): 400 g/L propyzamide formulated as a suspension concentrate.

Container size - 2 x 10 L.

Crops and Staging:

Apply to the following established crops between October 1 and freeze-up or very early spring*. Temperatures should be above freezing at time of application but should not exceed 12°C after application or a reduction in control may be observed. Applications are more effective if followed by a rain. Contact manufacturer for specific staging and application guidelines prior to application.

Established alfalfa, bird's-foot trefoil, and established pastures**.

* Early spring application for seed alfalfa only.

** Severe stand thinning may occur to pastures consisting primarily of crested wheatgrass, meadow fescue and timothy. Some thinning (10 to 15%) may occur with tall fescue and creeping red fescue.

Weeds, Rates and Staging:

Apply in late fall or very early spring (seed alfalfa only) prior to the emergence of weeds.

Established grass or grass/legume pastures for control of foxtail barley:

Brown, dark brown or gray wooded soils:

Kerb 50WP: 0.36 kg per acre.

Kerb SC: 0.45 L per acre.

Black soils:

Kerb 50WP: 0.45 kg per acre.

Kerb SC: 0.56 L per acre.

Established Alfalfa† and bird's foot trefoil†:

WEED	RATE (kg/acre)	
	Kerb WP	Kerb SC
Annual grasses, volunteer cereals, wild oat	0.71 kg†	0.89 L†
Quackgrass, orchardgrass, timothy, chickweed	0.91 to 1.32* kg	1.13 to 1.62* L
Dodder (fall application only)	1.3 kg	1.62 L

Note that complete control may not be achieved.

* Maximum 0.91 kg per acre† Including fall application on spring seeded crops. with spring application. Low temperatures and adequate moisture following application are needed for efficacy.

† Including fall application on spring seeded crops.

Caution: DO NOT use on soils with more than 6% organic matter. DO NOT apply to soils prone to flooding. DO NOT apply to pastures that contain high proportions of timothy, crested wheat grass or meadow fescue. Consult the manufacturer for other forage grass species sensitivities to Kerb.

Application Information:

Water Volume: 120 to 200 L per acre.

Nozzles and Pressures: Maximum 30 to 40 psi (200 to 275 kPa) with conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Dry soil conditions at time of weed emergence may result in reduced control. Approximately 3 inches of total precipitation is required for adequate activation. Best results when soil temperatures are low but above freezing.

Tank Mixes:

None Registered

Restrictions:

Rainfall: Surface applications are most effective if followed by 0.5 to 1 inch (1.25 to 2.5 cm) of rain within 2 days of application. Avoid application when heavy rain is forecast.

Re-entry: DO NOT re-enter treated areas for 24 hours.

Grazing: DO NOT graze or harvest for livestock feed within 90 days of the 1.32 kg/acre rate of *Kerb 50WSP* or the 1.62 L/acre rate of *Kerb SC*, and 60 days of application for lower rates.

Re-cropping: May be replanted to leafy vegetable crops after 30 days of treatment and root or tuber vegetables within 90 days of treatment. DO NOT re-plant to any other crops within 1 year of treatment.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. Do not freeze.

Buffer Zones: DO NOT contaminate domestic or natural water sources or wetlands.

Crop	Buffer zone* (meters†) for terrestrial habitat
Established grass pastures, established grass /legume pas- tures, alfalfa or trefoil grown for seed	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

Kerb WSP:



Caution – Poison

For an explanation of the symbols used here see page 11.

Ko-Act

Herbicide Group
2- tribenuron
4- 2,4-D
(Refer to page 38)

This product is a prepackaged tank mix of *Spike* (Tribenuron, page 283) and 2,4-D (page 66). Information listed is restricted to Crop, Weeds, Rates and Tank mixes. For other detailed restrictions and other general information on the component products see the product pages listed above.

Company:

Nufarm Agriculture

Formulation:

The *Ko-Act* package contains the following components:

Spike (PCP# 29653): 75% Tribenuron methyl, formulated as water soluble granule.

Container size - 2x160 g

2,4-D 700 ester (PCP# 27820): 2,4-D 660g/L as emulsifiable concentrate.

Container size - 2 x 8.69 L.

Crops and Staging:

Pre-seed burn-off prior to seeding the following crops:

Barley

Wheat

Rye

May also be applied to chem-fallow.

Weeds, Rates and Staging:

Spike at 4 g per acre plus 2,4-D 700 ester at 212 mL per acre.

Weeds controlled up to 10 cm or 3 leaf rosette or less, unless specified, include:

Chickweed

Dandelion

Flixweed

Hemp-nettle

Kochia*

* All biotypes

One case treats 80 acres.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

Mustard

Narrow-leaved hawk's-beard

Shepherd's-purse

Volunteer canola*

Tank Mixes:

Herbicides:

Glyphosate (360 g ae per acre*)

* see glyphosate page for rate conversion

See component products for more information on restrictions application details and handling. Use the most limiting restrictions across all components for the mix.

Liberty 150SN

Herbicide Group
10 - glufosinate
(Refer to page 38)

Company:

Bayer CropScience (PCP#28837)

Formulation:

150 g/L glufosinate ammonium formulated as a solution.

Container sizes - 13.5 L, 108 L, 432 L.

Crops and Staging:

Liberty Link Canola - cotyledon to early bolting stage. Temporary crop discolouration (bronzing) may be observed after application.

Note: A valid Liberty and Trait Agreement is required to purchase Liberty 150SN.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

WEED	WEED STAGE (from emergence to stage)	RATE (L/ACRE)	ACRES PER 13.5 L
Cow cockle	4 leaf	0.54	25
Green foxtail	6 leaf (max. 3 tillers)		
Barnyard grass	4 leaf	0.81	16.6
Wild mustard	5 leaf		
Lamb's-quarters, smartweed (lady's-thumb)	6 leaf		
Stinkweed	8 leaf		
Volunteer flax	2.5 inches (6 cm)		
Russian thistle	3 inches (8 cm)		
Wild buckwheat	3 leaf	1.08	12.4
Redroot pigweed, round-leaved mallow, quackgrass*	4 leaf		
Light to moderate infestations† of volunteer wheat, volunteer barley*	4 leaf (max. 2 tillers)		
Hemp-nettle (1 to 3 leaf pairs), shepherd's-purse	6 leaf		
Common chickweed (max. 4 leaf pairs), sow-thistle	8 leaf		
Kochia	3 inches (8 cm)		
Canada thistle*, scentless chamomile	4 inches (10 cm)		
Cleavers	2 whorls (nodes)	1.35	10
Stork's-bill and heavy populations of wild buckwheat	3 leaf		
Quackgrass (light to moderate** or heavy infestations*)†, volunteer wheat, volunteer barley*, wild oat	4 leaf (max. 2 tillers except quackgrass)		
Hemp-nettle	8 leaf (1 to 4 leaf pairs)		
Dandelion rosettes	6 in. (15 cm) across		
Flixweed, Canada thistle*	4 inches (10 cm)		
Quackgrass***	4 leaf	1.6	8.3
Canada thistle**	4 inches (10 cm)		

* Temporary top growth control. Plants may return from surviving growing points.

** Extended top growth control.

*** Season long control.

† The company does not provide guidelines for weed densities. When in doubt as to the infestation level, use the high rate or contact the manufacturer.

Second Application:

A second application of up to 1.35 L per acre may be made to fields that were treated initially with *Liberty* to a maximum total combined rate of 2.97 L per acre (1.62 L + 1.35 L). DO NOT apply more than 2.97 L per acre of *Liberty* in one season.

Application Information:

Water Volume:

Ground applications: 45 L per acre.

Aerial applications: 22 L per acre.

Nozzles and Pressure:

Ground Application: Use 40 psi (275 kPa) when using conventional 80° or 110° flat fan nozzles; 45 psi (310 kPa) when using check valves. Angle nozzles forward at 45° to improve coverage of vertical leaf surfaces. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium or larger droplets.

Aerial applications: DO NOT use raindrop nozzles. Use a combination of nozzles and pressure to provide ASABE coarse or larger droplet size distribution.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Liberty 150SN activity is influenced by environmental conditions. Cool temperatures (less than 10°C), drought, and low humidity conditions slow weed growth. Applications made under these stressed conditions may result in reduced weed control.

Tank Mixes:

Herbicides:

Clethodim (*Centurion* or *Select* only) (25.5 mL/acre or 120 acre/case).

Consult *Liberty 150SN* label for exact weeds controlled. For *Centurion* or *Select* tank mix add *Amigo* to the tank first at 0.5 L per 100 L spray solution followed by *Liberty* and then *Centurion* or *Select*. Consult label for specific mixing instructions.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Liberty 150SN* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Bayer CropScience also supports the following mixes that are not on the *Liberty* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Centurion* (50 mL/acre).

Insecticides: *Decis*, *Sevin XLR*.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Re-Entry: DO NOT re-enter treated areas for 24 hours after application, without protective clothing as for spraying.

Grazing: DO NOT graze the treated crop or cut for feed.

Re-cropping: No restrictions.

Aerial Application: May be applied by air.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	0	1
Fixed wing airplane or Helicopter	1	0	30

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.


DO NOT apply when dead calm or when winds exceed 16 km/hr when using unprotected booms or applying by air, or exceeding 25 km/hr when using shrouded booms.


Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

 Warning - Poison

 Caution - Skin Irritant

 Warning - Eye Irritant

For an explanation of the symbols used here see page 11.

Liberty 200 SN

Herbicide Group
10 - glufosinate
(Refer to page 38)

Company:

Bayer CropScience (PCP#25337)

Formulation:

200 g/L of glufosinate ammonium formulated as a solution.

Container sizes - 10 L

Crops and Staging:

Liberty 200 SN tolerant Corn only: 1 to 8 leaf stage. Refer to product label for appropriate method of determining crop leaf stage.

Liberty tolerant soybean varieties only: up to the start of flowering and prior to canopy closure.

Weeds Rates and Staging:

Weeds controlled with 0.61 L per acre rate

WEED	WEED STAGE (from emergence to stage)
Cocklebur	4 leaf
Green foxtail, Proso millet, Ragweed	5 leaf
Redroot pigweed, Shepherd's-purse	6 leaf
Chickweed	8 leaf

Weeds controlled with 0.81 L per acre rate

WEED	WEED STAGE (from emergence to stage)
Perennial sow-thistle, Wild buckwheat, Wild mustard, Wild oat, Yellow foxtail	4 leaf
Barnyard grass, Eastern black nightshade	5 leaf
Canada thistle*, Field bindweed*, Lady's-thumb, Lamb's-quarters, Wormseed mustard	6 leaf

WEED	WEED STAGE (from emergence to stage)
Ragweed	7 leaf
Stinkweed	8 leaf

* season long suppression.

Weeds controlled with 1.0 L per acre rate

WEED	WEED STAGE (from emergence to stage)
Quackgrass**	4 leaf

** season long suppression, apply with ammonium sulfate, 2.4 L per acre (49% solution) or 1.2 kg per acre (99%).

Second Application:

A second application may be made to fields treated initially with up to 1 L per acre, if weeds and crop are at the correct leaf staging. DO NOT apply more than 2 L per acre *Liberty 200SN* to a crop in a single season.

Split Application Program:

For season long control of the weeds above a split application of *Liberty 200SN* may be employed. The first application must be a minimum of 0.81 L per acre made at the correct weed staging. For the second application of a 0.51 L per acre rate may be used. The second application timing must be made as soon as the second flush of weeds occurs and before the maximum leaf stage for the crop.

Application Information:

Water Volume: A minimum of 45 L per acre.

Nozzles and Pressure: Use 25 to 40 psi (175 to 275 kPa) when using conventional 80° or 110° flat fan nozzles. Angle nozzles forward at 45° to improve coverage of vertical leaf surfaces. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Liberty 200SN activity is influenced by environmental conditions. Cool temperatures (less than 10°C), drought and low humidity conditions slow weed growth. Applications made under these stress conditions may result in reduced weed control.

Weed control may also be reduced when heavy dew, fog, or mist are present at the time of application.

Tank Mixes:

None registered.

Restrictions:

For use on glufosinate-ammonium tolerant field corn only.

Rainfall: Within 4 hours of application may reduce control.

Re-Entry: DO NOT re-enter treated areas for 24 hours after application, without protective clothing as for spraying.

Grazing: DO NOT graze treated fields within 20 days of application.

Preharvest Interval: Leave 86 days between application and grain harvest.

Re-cropping: No re-cropping restrictions the year after treatment.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	0	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack applications do not require a buffer.

Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Caution - Poison



Caution - Skin Irritant



Warning - Eye Irritant

For an explanation of the symbols used here see page 11.

Linuron*

**Herbicide Group
7 - linuron**
(Refer to page 38)

Company:

Tessenderlo Kerley Inc. (*Lorox L*)

United Agri Products (*Linuron 400*)

The following recommendations are a blend of recommendations of all linuron products. Consult the individual product labels for specific recommendations.

* NOTE: The Pest Management Regulatory Agency has re-evaluated linuron products and have proposed to phase out the use of all linuron products. As a result of this phase out, this page may not appear in future editions of this publication.

Formulations:

Linuron 400 (PCP#15544): 400 g/L linuron formulated as a suspension concentrate.

Container size - 10 L.

Lorox L (PCP#16279): 480 g/L linuron formulated as a suspension concentrate.

Container size - 10 L.

Crops, Rates and Stages:

Post-emergent applications only:

CROP	STAGE	LINURON 400 (L/acre)	LOROX L (L/acre)
Spring wheat (including durum), oats and barley*	2 to 4 leaf stage	0.20 to 0.26	0.17 to 0.22
Field corn (post-emergent** directed spray, do not spray over top of corn)	Apply when corn is at least 15 inches (38 cm) high (highest leaf on free standing plant)	1.16 to 2.18	0.97 to 1.82
Caraway, coriander	Apply when in the 2 to 4 leaf stage	—	0.50 to 0.67
Dill†	Apply when dill has at least 2 full leaves developed	—	0.77 to 1.9
Shelterbelts (caragana, green ash, Siberian and American elm, Manitoba maple, poplar, willow, white spruce, Colorado spruce, Scots pine)	Apply as an overall spray to dormant stock or as a directed spray if buds have broken.	2.18	1.82
Short Rotation Intensive Poplar	Apply as a directed spray under plants that have been established for 1 year or more	—	1.82

* Only when tank mixed with MCPA amine at 0.34 to 0.45 L per acre (500 formulation) or 0.4 to 0.57 L per acre MCPA K (400 formulation).

** Use lower rate when weeds do not exceed 2 inches (5 cm) and higher rate for weeds up to 8 inches (20 cm) in height, preferably before they are 5 inches (13 cm) high. Requires the addition of a mineral oil surfactant blend at 1 to 2 L per 100 L or spray solution or spray oil at 1 to 2 L per 10 L of spray solution. See oil labels for directions. DO NOT apply if *linuron* has been applied pre-emergent.

† A split pre-emergent/post emergent application of *linuron* may be made in dill. See below for more information.

Pre-emergent surface (not incorporated) applications for use on loam to clay soils only:

	LINURON 400 (L/acre)		LOROX L (L/acre)	
	<i>less than 2%</i>	<i>from 2 to 5%</i>	<i>less than 2%</i>	<i>from 2 to 5%</i>
Soil organic matter				
Field corn	1.09*	1.58	0.91*	1.31*
Soybeans	1.09 to 1.58	1.58 to 2.18	0.91 to 1.31	1.31 to 1.82
Sweet white lupins	1.01	1.50	0.85	1.25
Potatoes	1.11 to 1.72	1.72 to 2.22	0.91	1.82
Dill†	—	—	0.53 to 0.77	0.77 to 1.0

* Must be tank mixed. Refer to specific labels for registered tankmix partners.

† A split pre-emergent/post emergent application of *linuron* may be made in dill. See below for more information.

If used on sandy soils, severe crop injury may result.

Seed the crop at least 2 inches (5 cm) deep. Make only one application per year to field crops.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Split applications:

This product may also be applied to dill as a split pre/post-emergent application. A pre-emergent surface application of up to 0.77 L per acre, followed by a second post-emergent application, no sooner than two weeks after the first, of up to 1.0 L per acre. Minimum staging for post-emergent applications given above still applies.

Banded Applications:

This product may also be applied in a narrow band directly over the row in wide rowed crops if another method is to be used for weed control in between the rows. For band treatment, use proportionately less; for example, for 10 inch band on 30 inch row, use 1/3 of the broadcast rate.

Weeds and Staging:

Post-Emergence

When tank mixed with MCPA amine in cereals, the following weeds are controlled:

Chickweed	Lamb's-quarters
Corn spurry	Ragweed (common, giant)
Cow cockle	Redroot pigweed
Flixweed	Shepherd's-purse
Green foxtail (suppression possible)	Stinkweed
Green smartweed	Stork's-bill
Hemp-nettle	Tartary buckwheat
Lady's-thumb	Wild buckwheat

Apply when annual broadleaf weeds are in the 2 to 4 leaf stage and when green foxtail is in the 1 to 3 leaf stage.

In shelterbelts, apply when weeds are less than 4 inches (10 cm) tall.

Pre-emergent surface treatments and Post-emergent applications in corn and shelterbelts:

Sufficient moisture (1 to 2 inches or 3 to 5 cm) in the form of rainfall or irrigation is necessary within 7 to 10 days of a pre-emergence application or poor weed control will result.

Barnyard grass*	Purslane
Common chickweed	Ragweed (common)
Common groundsel [■]	Shepherd's-purse
Corn spurry [■]	Smartweed (annual)
Crabgrass*	Sow-thistle (annual, perennial [■] seedlings only)
Dandelion (seedlings only) [■]	Stinkweed [○]
Foxtail (green and yellow)*	Wild buckwheat
Goosefoot	Wild radish [■]
Knotweed	Witchgrass
Kochia [■]	Wormseed mustard
Lamb's-quarters	
Pigweed (prostrate [■] , redroot)	
Plantain (seedlings only) [■]	

*partial control

[■] Not registered with Loro L.

[○] Not registered with Linuron 400.

Application Information:

Water Volume:

Post-emergent:

Cereals, coriander & caraway: 40 L per acre.

Field Corn directed spray: 69 to 138 L per acre.

Dill: 89 to 178 L per acre

Pre-emergent surface: 81 to 162 L per acre.

Shelterbelts: 90 to 180 L per acre.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional stainless steel flat fan nozzles. Low drift nozzles may require higher pressures for proper

performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift. Use a 50 mesh or coarser line strainers and screens.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

In post-emergent applications the best weed control occurs when temperatures are moderate, when relative humidity is high and when soil moisture is adequate. Injury to cereals (crop lightening) will occur when the crop is under stress because of drought or disease. This injury is worse when the product is applied at advanced leaf stages. In pre-emergent surface treatments, rainfall or irrigation (1 to 2 inches or 3 to 5 cm) is required to move linuron into the root zone of germinating seeds. Insufficient moisture will result in poor weed control. Drought conditions after application will result in little to no weed control. If rainfall does not occur within 7 to 10 days of application and prior to crop emergence, a shallow rotary hoeing (0.75 to 1.5 inches/ 2 to 4 cm) should be made to mix the top layer of soil to help activation. Avoid covering treated ground with un-treated soil. If unusually heavy rain follows application, severe crop injury may result from herbicide in the root zone of the crop. DO NOT use on sandy soils or severe crop injury will result.

Tank Mixes:

Herbicides:

In Cereals: For post-emergent applications in cereals, linuron must be tank mixed with MCPA amine or MCPA K. DO NOT tank mix with other herbicides.

In Corn: Atrazine, Dual II Magnum and Primextra II Magnum. Not all linuron products have the same tank mix options, refer to specific labels.

Fertilizers: None registered.

Insecticides: None registered.

Note: The above mixes are those listed on the linuron labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label for post-emergent applications; required interval may be up to 8 hours. Pre-emergent applications require rainfall for activation. Contact manufacturer for more information.

Grazing: DO NOT graze treated crops or cut for feed prior to crop maturity.

Preharvest Interval:

Sweet Corn: DO NOT harvest within 50 days of treatment.

Field Corn: DO NOT harvest within 60 days of treatment or until after tassel emergence.

Caraway, Coriander & Dill: DO NOT harvest within 60 days of treatment.

Re-cropping: If the intended crop fails, fields treated with pre-emergent surface applications of linuron, may be seeded back to corn, soybeans, sweet white lupins, or potatoes. Till the soil thoroughly before reseeding. No restrictions 1 year after treatment.

Aerial Application: DO NOT apply by air.

Storage: DO NOT store liquid *Linuron* formulations at temperatures below 5°C. *Lorox L* may be frozen.

Tank Cleaning:

Refer to page 15.

Lontrel 360**Herbicide Group**

4 - clopyralid

(Refer to page 38)

Company:

Dow AgroSciences (PCP# 23545)

Formulation:

360 g/L clopyralid formulated as a solution.

Container size - 3.4 L, 4.45 L, 110 L, 115 L, 208 L, 454 L, Bulk

Crops Rates and Staging:

Applications of 0.17 to 0.23 L per acre:

Barley, spring wheat (NOT including durum), oat - 3 to flag leaf emergence stage.

Applications of 0.23 to 0.34 L per acre:

Flax, solin (low linolenic acid flax) - 2 to 4 inches (5 to 10 cm) in height.

Applications of 0.17 to 0.34 L per acre:

Canola - 2 to 6 leaf stage. Apply *Lontrel* to canola (Argentine - *B. napus* and Polish - *B. rapa*) varieties only; application to any other canola type oilseeds may result in injury to the crop.

Seedling forage grasses* - 2 to 4 leaf stage.

Established grasses* - at the shot blade stage, or in the fall after harvest or in early spring.

Seedling and established grasses* for forage and seed production include:

Bromegrass (smooth)
Fescue (creeping red, meadow, tall)
Kentucky bluegrass
Meadow foxtail
Orchardgrass

Reed canarygrass
Timothy
Wheatgrass (crested, intermediate, slender, streambank, tall**)
Wildrye (Altai, Russian)

Applications of 0.34 L per acre:

Summerfallow - Stage according to weeds.

Shelterbelts* containing villosa lilac, acute willow, Colorado spruce, white spruce, buffaloberry and chokecherry.*

Plantation poplar (including hybrid poplar)*

* **NOTE:** Since these uses are registered under the User Requested Minor Use Label Expansion program, the manufacturer assumes no responsibility for herbicide performance. **Users of this product for these uses do so at their own risk.**

** for forage use only

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

The following weeds will be controlled at 0.17 L per acre when young and actively growing:

Alsike clover Vetch (*Vicia* sp.)

The following weeds will be controlled from 0.23 to 0.34 L per acre when young and actively growing:

Common groundsel	Scentless chamomile
Common ragweed	Wild buckwheat
Perennial sow-thistle (top growth only)	Volunteer alfalfa -2 to 20 inches (5 to 50 cm) tall

Canada thistle - after all thistles have emerged and when the majority are in the rosette to pre-bud stage;

0.17 L per acre Provides top growth control of Canada thistle for 6 to 8 weeks.

0.23 L per acre Provides season long control of Canada thistle. Not all root stalks will be killed and some regrowth may occur by the end of the growing season.

0.34 L per acre Provides season long control of Canada thistle with suppression into the following year.

Spotted and diffuse knapweed is controlled up to flower emergence at 0.28 L per acre.

Sheep soren and oxe-eye daisy is controlled at 0.34 L per acre.

Application Information:

Water Volume: 40 to 89 L per acre.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) with conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressures designed to deliver thorough, even coverage with ASABE coarse droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Poor control may occur under dry conditions. Injury to flax may occur when tank mixing with MCPA. To reduce the risk of crop injury, DO NOT apply tank mixes if temperature exceeds 27°C.

Tank Mixes:

Herbicides:

Lontrel applications following applications of products containing bromoxynil (*Approve*, *Badge*, *Bromotril*, *Buctril M*, *Koril*, *Logic M*, *Mextrol*, *Pardner*, *Thumper*) should be delayed by 14 days to allow the Canada thistle to recover from leaf burn.

Recommended rates of *Lontrel* may be used for each crop unless otherwise indicated.

In Canola:

Poast Ultra plus Merge adjuvant.

Select plus Amigo Adjuvant (*Lontrel* at 0.17 to 0.34 L/acre).

In Canola (ROUNDUP READY varieties only):

*Roundup Transorb** (*Lontrel* at 0.112 L/acre).

In Canola (CLEARFIELD varieties only):

*Odyssey*** (*Lontrel* at 0.17 to 0.23 L/acre).

* Roundup Ready varieties only

** CLEARFIELD varieties only

In Flax:

Poast Ultra plus Merge adjuvant.

Poast Ultra + MCPA Ester (rates as above)

Select plus Amigo adjuvant.

Lontrel at 0.17 L/acre for the following mix:

MCPA amine or ester (0.28 to 0.38 L/acre - 600 g ae/L formulation)

In Spring wheat (NOT including durum) and barley:
Lontrel at 0.11 to 0.17 L/acre for the following mix:

2,4-D or MCPA (amine or ester) at label rates.

In Oat:

MCPA amine or ester (*Lontrel* at 0.11 to 0.17 L/acre).

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *Lontrel* label only. To check for other possible mixes see the blue fold out chart inside the back cover.

Dow AgroSciences also supports the following mixes that are not on the *Lontrel* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Assure II*, *Muster*, *Muster+any of Assure II*, *Clethodim*, *Poast Ultra*, *Pursuit*, *Solo*.

Adding ingredients in the correct order is critical for optimum performance.

Check labels of products to be mixed for directions.

General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfall period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT enter treated fields for 12 hours.

Grazing: Crops or areas treated with this product may be grazed immediately following treatment.

Re-cropping: *Lontrel* residues in the soil may affect succeeding crops. The year after application, replant to wheat, barley, oats, rye, flax, forage grasses, mustard or canola.

DO NOT use manure from animals fed or bedded with *Lontrel*-treated straw, except on fields that are to be sown to *Lontrel*-tolerant crops.

Aerial Application: DO NOT apply by air.

Storage: Store in heated storage. If product is frozen, bring to room temperature and agitate before use.

Buffer Zones: Avoid contamination of or drift toward non-target land, water or irrigation ditches.

Tank Cleaning:

No cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Caution - Poison



Danger - Eye Irritant

For an explanation of the symbols used here see page 11.

MCPA

Herbicide Group

4 - MCPA

(Refer to page 38)

Company and Formulation

IPCO (600 Ester** - PCP#27802; 500 Amine* - PCP#20308; 400 (K) Potassium salt* - PCP#20305; 300 (Na) Sodium salt* - PCP#20306)

Nufarm Agriculture (600 Ester** - PCP#27803; 600 Amine* - PCP#28384; 300 (Na) Sodium salt* - PCP#14718)

UAP (600 Ester** - PCP#27804; 500 Amine* - PCP#9516; 300 (Na) Sodium salt* - PCP#9858)

Loveland Products (formerly Viterro) (600 Ester** - PCP#29001; 500 Amine* - PCP#29244)

Farmers of North America (MPower MCPA Ester 500 ** - PCP#30462; MPower MCPA 500 Amine* - PCP#30461)

* formulated as a solution.

** formulated as an emulsifiable concentrate.

Crops, weeds and rates and other application details may differ with different product labels. Consult specific product labels for more information.

Crops, Rates and Staging:

The maximum safe rates for various crops are given below. Higher rates used for harder to control weeds (see "Weeds, Rates and Staging") may cause crop injury. Application rates for individual products may vary from those listed. Refer to the label for product specific use rates. Rates greater than these for harder to control weeds may cause crop injury. When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

CROP	STAGE	RATE (L/ACRE)			
		AMINE 500	AMINE or ESTER 600	K SALT	Na SALT
Wheat (spring and durum), barley	3 leaf to early flag leaf.	0.45	0.42	0.53	0.81
Oats	3 leaf to early flag	0.45	0.36	0.53	0.81
Spring rye	2 leaf to early flag leaf.	0.45	0.42	0.53	0.81
Flax (not Solin varieties)	2 inches (5 cm) in height to prebud stage. Apply at 2 to 4 inches (5 to 10 cm) in height for maximum crop tolerance.	0.4	0.28 (E) or 0.34 (A)	0.65	0.71
Winter wheat (WW), fall rye (FR)	In spring, apply from the time growth commences until the early flag leaf stage.	0.45	0.42	0.61 (WW) 0.40 (FR)	0.81
Corn	As a broadcast spray up to 6 to 7 in. (15 to 18 cm) tall or 6 leaf stage. Up to 3 weeks before tassling as a directed spray using drop nozzles.	0.45	0.37 (Amine only)	0.51	0.61
Peas	Vines 4 to 7 inches (10 to 18 cm) long. For short-statured, determinate flowering peas, apply at the early stages within this range.	0.22*	0.17 (Amine only)	NR	0.36*
Cereals underseeded to alfalfa (not Flemish varieties)	Apply when the majority of seedling legumes are in the 1 to 3 trifoliate leaf stage.	0.22	0.19 (Amine only)	NR	0.4
Underseeded alsike, ladino and red clover	Apply when the majority of seedling legumes are in the 1 to 3 trifoliate leaf stage.	0.28	NR	NR	0.4

(E) or (A) indicates Ester or Amine formulations.

Crops, Rates and Staging continued on next page.

Crops, Rates and Staging *continued*

CROP	STAGE	RATE (L/ACRE)			
		AMINE 500	AMINE or ESTER 600	K SALT	Na SALT
Red clover [◇] Seedling (seed and forage) Established [◇] (seed only)	Seedlings: 1 to 3 trifoliate stage. Do not feed to livestock in the first year. Established: Apply at the breaking of dormancy in the spring up to 7.5 cm.	0.23	0.19 (Amine only)	NR	NR
Grass pastures	Spring or fall.	1.42	1.13 (E) or 1.42 (A)	NR	0.71
Seedling forage** grasses (not for seed)	Apply from the 3 leaf stage to the shot blade stage.	0.45**	NR	NR	NR
Established forage** grasses (not for seed)	Apply in the spring up to the shot blade stage or in the fall after harvest.	0.45**	NR	NR	NR

(E) or (A) indicates *Ester* or *Amine* formulations.

NR = Not Registered

* The rates given are lower than the registered rates for peas. Less than the maximum label rates are recommended because of crop injury concerns.

** MCPA is NOT registered for use on forage grasses, any such use is entirely at the risk of the user.

◇ Nufarm Agriculture MCPA *Amine* only.

Formulation Characteristics:

Formulation	Risk of Vapour Drift	Activity on Weeds	Risk of Crop Injury
LV Ester	Medium	Fast	Medium
Amine	Very Low	Medium	Low
Salts	Very Low	Slow	Very Low

Weeds, Rates and Staging:

Apply at lower rates when weeds are small (2 to 4 leaf stage) and actively growing. Higher rates are needed when weeds are larger, in heavy populations, or growing under stressful conditions (excessively cold, hot, dry or wet).

NOTE: The following rates are a general range for all products. Rate ranges for individual products may differ slightly. Consult the product label for specific rates for each application.

+ Not controlled by MCPA K salt formulation

++ Not controlled by MCPA K or Na salt formulations

Susceptible weeds:

*Amine 500 formulations – 0.28 to 0.45 L per acre**Amine and Ester 600 formulations – 0.24 to 0.36 L per acre**K formulations – 0.61 to 0.71 L per acre.**Na formulations – 0.5 to 0.81 L per acre.*

Burdock

Cocklebur

Flixweed (late fall applications
or small seedlings)*

Kochia

Lamb's-quarters

Mustards (except dog and tansy)

* Winter annual weeds

Prickly lettuce+

Ragweed

Russian pigweed

Shepherd's-purse*

Stinkweed*

Wild Radish

Wild Sunflower+

Harder to control weeds:

Amine 500 formulations – 0.45 to 0.71 L per acre.

Ester 600 formulations – 0.42 to 0.61 L per acre.

K formulations – 0.71 to 0.81 L per acre.

Na formulations – 0.81 to 1.1 L per acre.

Annual sow-thistle+	Hemp-nettle
Biennial wormwood+	(suppression only)
Bluebur+	Mustard (dog and tansy)+
Common peppergrass+	Plantain+
Curled dock+	Purslane+
Flixweed (overwintered	Redroot pigweed+
rosettes prior to bolting)+	Smartweed (annual)+
Goat's-beard+	Tumble pigweed+
Goosefoot+	

Top growth control only (rates as for harder to control weeds):

Blue lettuce+	Field horsetail++
Canada thistle	Hedge bindweed+
Corn spurry++	Hoary cress+
Dandelion++	Leafy spurge+
Gumweed+	Perennial sow-thistle+
Field bindweed+	Russian thistle++

Application Information:**Water Volumes:**

Cereals, flax, pastures, forage grasses: 40 to 81 L per acre.

Peas: Minimum 61 L per acre.

Cereals Underseeded to Forage Legumes: 61 to 81 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets.

Tank Mixes:**Herbicides:****In Wheat and barley:**

Avenge (500 ester only).

Banvel II (amine and K salt only).

Pardner (K salt only).

Linuron and *Sencor* (500 amine only).

In Oats:

Pardner (K salt only).

Linuron (500 amine only).

Not all brands are labelled for tank mixing. Check the product label prior to use for registered mixes and rates. Follow all precautions and restrictions on both labels.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: Liquid nitrogen (28-0-0) may be used in place of water as a carrier with certain amine formulations for application in spring to winter wheat or fall rye.

Note: The above mixes are those listed on the *MCPA*

labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours of *MCPA Na salt* or *MCPA K salt*, 4 hours of *MCPA amine*, or 2 hours of *MCPA ester* application will reduce control.

Re-Entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze within 7 days of application.

Re-cropping: No restrictions the year after application.

Aerial Application: Some products may be applied by air to specific crops. Check the label for detailed instructions.

Storage: *MCPA ester* may be frozen. DO NOT freeze *MCPA amine*, *MCPA sodium salt* and *MCPA K*.

Buffer Zones:

Crop	Application method	Buffer Zones (metres†)		
		Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Cereals, Flax	Ground*	1	1	4
	Fixed wing aircraft	1	0	60
	Helicopter	1	0	50
Legume forages	Ground*	1	1	4
	Fixed wing aircraft	1	1	25
	Helicopter	1	1	25
Stubble, pastures	Ground*	1	1	4
	Fixed wing aircraft	15	0	60
	Helicopter	15	0	50

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack applications do not require a buffer.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Best weed control occurs when temperatures are above 21°C (daytime) or 10°C (night time) and humidity is above 70 percent. DO NOT apply if temperature exceeds 27°C. If applying to flax, injury and a delay in maturity may result from application under hot or humid conditions. Extremely hard water may reduce performance or cause problems in spraying the product.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B'

found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

MCPB / MCPA

Herbicide Group
4 - MCPB & MCPA
(Refer to page 38)

Company:

IPCO (*Clovitox Plus* – PCP# 24336)
Nufarm Agriculture (*Tropotox Plus* – PCP#8211)
UAP (*Topside* – PCP#22003)

Formulation:

375 g/L MCPB, 25 g/L MCPA present as sodium salts and formulated as a solution. Container size - 10 L.

Crops, Rates and Staging:

Registered for all products:

Apply 1.11 to 1.72 L per acre. Apply only that needed to control the target weeds.

DO NOT make more than one application of this or other product containing the same ingredients per year.

CROP	STAGE
Pea	3 to 6 expanded leaves.
Clover (alsike, ladino, red, white Dutch, wild white)	Monofoliolate to 3 trifoliolate leaf stage (with or without a cover crop).
Oats, wheat, rye or barley (alone or as a companion crop)	2 leaf to flag leaf stage.
Field corn	45 cm high to the start of tassling – use drop nozzles.
Established pasture	After grazing or cutting when weeds have regrown to a susceptible stage.

Seedling Forage Grasses:

Apply at 1.11 to 1.42 L per acre from the 2 to 4 leaf stage:

Bromegrass (smooth, meadow)	Timothy
Fescue (altai, red, meadow, tall)	Wheatgrass (crested, creeping intermediate, northern, pubescent, slender, stream-bank, tall, western)
Green needlegrass	Wild rye (altai, Russian)
Reed canarygrass	

Registered for *Tropotox Plus*, and *Clovitox Plus* only:

Seedling alfalfa for seed production* at the 3 to 6 trifoliolate stage.

NOTE: Seedling alfalfa vigour may be reduced in the year of treatment, however, the crop recovers and yield will not normally be affected.

* Since this use is registered under the User Requested Minor Use Label Expansion program, the manufacturers assume no responsibility for herbicide performance. Users of this treatment on seedling alfalfa do so at their own risk.

Weeds, Rates and Staging:

WEEDS	STAGE	RATE (L/ACRE)
Ball mustard, Lamb's-quarters, Stinkweed, Wild mustard, Wormseed mustard	Seedlings	1.11
Annual sow-thistle*, Hemp-nettle*, Redroot pigweed, Ragweed, Shepherd's-purse, Volunteer rapeseed (including canola), Wild radish*	Seedlings	1.72

WEEDS	STAGE	RATE (L/ACRE)
Curled dock, Perennial sow-thistle**, Plantain	Rosette	1.72
Bull thistle	Rosette to early bud	1.72
Buttercup (Creeping, Tall), Field bindweed	In spring during rapid growth	1.72
Canada thistle	6 inches (15 cm) to early bud	1.72
Horsetail*	6 inches (15 cm)	1.72

* Suppression only

** Top growth control only

Application Information:

Water Volume:

Clovitox Plus: 71 to 91 L per acre.

Tropotox Plus, Topside: 61 to 81 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Damage to peas or seedling forage legumes may occur if the crop is sprayed when under drought or disease stress. Under extremely hot or humid conditions, crop injury may be severe. DO NOT apply when temperatures are over 27°C. Best activity on weeds occurs in warm weather.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze crop treated with *Topside* or cut for hay. DO NOT graze or cut seedling forage grasses in the

year of treatment. Cereals treated with *Tropotox* or *Clovitox* may be used for grazing or cut for greenfeed or hay 30 days after application. Forage legumes and peas treated with *Clovitox* may be used for animal feed 30 days after application. Withdraw meat animals from fields treated with *Tropotox* or *Clovitox* at least 3 days before slaughter.

Re-cropping: No restrictions listed. Phenoxy herbicides can persist in soils for weeks, particularly if dry or cool weather persists. DO NOT seed sensitive crops immediately after spraying.

Aerial Application: *Clovitox* may be applied by air to established pasture and cereal crops (not underseeded to clover).

Storage: DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	4

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:

Clovitox Plus:



Danger – Poison



Danger - Corrosive to eyes

Tropotox Plus & Topside:



Caution – Poison

Tropotox Plus:



Warning – Contains the allergen caseinate (milk)

For an explanation of the symbols used here see page 11.

Mecoprop-p

Herbicide Group
4 - mecoprop-p
 (Refer to page 38)

Company:

United Agri-Products (*Mecoprop-P* - PCP# 27891)
 Nufarm Agriculture (*Comptox* - PCP# 27824)

Formulation:

150 g/L mecoprop-p present as potassium salt.
 Container size - 10 L.

Crops and Staging:

Spring wheat (including durum), barley and oats - 3 leaf to flag leaf stage.

Weeds and Staging:

Apply to weeds from the 2 to 4 leaf stage.

Black medic*	Clover (volunteer)
Canada thistle (top growth control only)	Corn spurry
Chickweed	Lamb's-quarters*
Cleavers	Plantain
	Wild mustard*

* not *Comptox*

Rates:

2.2 to 2.8 L per acre.

Use the high rate for weeds in an advanced stage of growth.

Application Information:

Water Volume: 81 to 121 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Apply in warm weather under good growing conditions. Avoid spraying in very hot weather or in drought conditions.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Grazing: DO NOT graze or feed treated crop to livestock prior to crop maturity.

Re-cropping: No restrictions the year after application.

Aerial application: DO NOT apply by air.

Storage: DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	0	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Caution - Poison

For an explanation of the symbols used here see page 11.

Metribuzin

Herbicide Group

5 - metribuzin

(Refer to page 38)

Company:

Bayer CropScience (*Sencor Solupak 75 DF, Sencor 75 DF*)

United Phosphorus Inc. (*TriCor 75 DF*)

Formulations:

Sencor Solupak 75 DF (PCP# 20968): 75% metribuzin formulated as a dispersible granule. Container size - 2.5 kg (5 water soluble bags, 500 g each).

Sencor 75 DF (PCP# 17242): 75% metribuzin formulated as a dispersible granule. Container size - 2.5 and 5 kg.

TriCor 75 DF (PCP# 30661): 75% metribuzin formulated as a dispersible granule. Container size 2.5 kg.

Crops, Rates and Staging:

CROP	APPLICATION TIMING	RATES (g/acre)
Barley	2 to 5 leaf stage.	80 to 152
Spring wheat (including durum)	2 to 5 leaf stage.	80 to 111
Pea (field and processing)*	Preplant incorporated (when tank mixed with <i>Rival</i> or <i>Treflan</i> E.C.).	Spring: 152 to 192 Fall: 190 to 223
Pea (field only)*	Post-emergence - up to 6 inches (15 cm) of vine length. For short-statured, determinate flowering peas, apply at the early stages within this range.	111 to 152
	Split post-emergent applications**.	First split application: 60 to 80 Second split application: 7 to 10 days later with rates within these ranges.
Lentil*	Single or split applications**: Plants up to 6 inches (15 cm) of vine length. For maximum crop tolerance, apply at the 1 to 4 above ground node stage.	111
		First split application: 60 to 80 Second split application: 7 to 10 days later with rates within these ranges.
Chickpea*	Up to 2.5 inches (6 cm) in height, when vines have 1 to 3 above ground nodes. <i>Note: application past recommended growth stage may result in severe crop injury.</i>	111
Potato (except Belleisle or Tobique)***	Preplant incorporated (with <i>Eptam</i>).	152 to 223
	Pre-emergence in sprinkler irrigation systems (apply only in a tank mix with <i>Eptam</i> 8-L).	152 to 223

Crops, Rates and Staging continued on next page.

Crops, Rates and Staging *continued*

CROP	APPLICATION TIMING	RATES (g/acre)
Potato (except Atlantic, Belleisle, Eramosa, Tobique and red-skinned or early maturing varieties)***	Early post-emergence (up to 4 inches or 10 cm in height).	151
Soybean***	Preplant incorporated (tank mixed with <i>Treflan</i> EC).	111 to 223
Fababean	Preplant incorporated (tank mixed with <i>Treflan</i> EC).	Spring: 111 to 223 Fall: 190 to 223

* DO NOT use on lentils, peas or chickpeas seeded less than 2 inches (5 cm) deep or in soils with less than 4 percent organic matter.

** Under certain field or weather conditions a split application may provide better weed control and crop tolerance than single applications. The first application should be made at the cotyledon to 2 leaf stage of the weeds. The second application should be made when a second flush of weeds have emerged or if weeds which were more advanced at the time of the first application have started to show regrowth. The split applications are normally 7 to 10 days apart.

*** Consult manufacturer or seed supplier for varietal tolerances to *metribuzin* applications in soybean and potato.

Note: When *metribuzin* is tank mixed with *Trifluralin* in peas, fababeans, and soybeans, refer to product label for maximum rates that can be applied on light textured soils.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Post-emergence applications should be made when weeds are small – 2 inches (5 cm) in height or diameter.

Split applications (postemergence on lentils and peas) – 1st application at cotyledon to 2 leaf stage of weeds. 2nd application (if necessary) 7 to 10 days after the first.

Post-emergence at 81 g per acre:

Weeds controlled in spring wheat, barley, field pea and suppressed in lentil and chickpea:

Chickweed	Stinkweed
Green smartweed	Volunteer canola
Hemp-nettle*	Wild mustard
Lamb's-quarters	

Additional weeds controlled in spring wheat and barley only:

Lady's-thumb	Redroot pigweed
--------------	-----------------

Post-emergence at 111 g per acre:

Weeds controlled in spring wheat, barley, potato, field pea, and suppressed in lentil and chickpea:

Weeds above plus:

Ball mustard	Hemp-nettle
Corn spurry	Tartary buckwheat

Additional weeds controlled in spring wheat and barley only:

Common groundsel	Wormseed mustard
Night-flowering catchfly	

Post-emergence at 152 g per acre in spring wheat and barley only:

Weeds above plus:

Henbit	Russian thistle
--------	-----------------

Post-emergence at 152 g per acre in potatoes only:

Weeds listed for peas above plus:

Lady's-thumb	Shepherd's-purse
Redroot pigweed	

Preplant Incorporated in fababeans, lentils, field pea and soybean:

Must be applied in tank mix with *Treflan* EC or *Rival*:

Chickweed	Lamb's-quarters
Corn spurry	Stinkweed
Green smartweed	Volunteer canola
Hemp-nettle	Wild mustard

Plus weeds controlled by either *Rival* or *Treflan* EC.

* Use the high rate for best control.

** Suppressed only in lentils and chickpeas.

Application Information:

Water Volume:

Preplant incorporated: 40 L per acre.

Post-emergence applications:

Cereals - 40 L per acre.

Lentils, peas, chickpeas - 70 L per acre.

Nozzles and Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets. If using conventional flat fan nozzles use a maximum of 30 to 40 psi (200 to 275 kPa) with opening no smaller than 8002 or TK2 with 50 mesh screens. For lentils, peas and chickpeas use nozzles no smaller than 8003 or TK3. Angle nozzles 45° forward to achieve better coverage of vertical weed targets.

Incorporation: All plant growth and stubble should be thoroughly worked into the soil before treatment. Apply directly to the soil surface. Two incorporations are required at right angles for thorough mixing. The first incorporation must be made within 24 hours of spraying. For fall applications, it is preferred that both incorporations be done in the fall. The second incorporation may be delayed until spring to conserve trash; however, both incorporations must be done the recommended depth.

Incorporate with a tandem disc, discer or field cultivator (Vibrashank type). Set equipment to work at a depth of 3 to 4 inches (8 to 10 cm). Operate disc implements at 4 to 6 mph (7 to 10 km/hr), cultivators at 6 to 8 mph (10 to 13 km/hr).

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions

Crop height reductions or yellowing may occur if high temperatures occur within 48 hours of application. Cold, cloudy weather or frost within 3 days of application will also aggravate injury. If frost occurs, allow 4 to 5 days for crop to recover prior to applying *metribuzin*. Heavy rainfall soon after application to peas, lentils and chickpeas can result in stand reduction on soils with less than 4 percent organic matter.

Tank Mixes:

Herbicides:

In spring wheat or barley:

Dicamba, *Target*, MCPA amine or 2,4-D amine.

In potatoes (post emergent) Sencor 75 DF and TriCor only:
*Prism**

In potatoes (preplant incorporated):

Eptam 8-E (Required).

In fababeans (preplant incorporated):

Treflan I/C (Required).

In soybeans (preplant incorporated):

Treflan I/C (Required).

In peas:

Treflan (PPI)

Rival (PPI).

Sencor 75 DF or *TriCor* at 77 g/acre plus 0.19 L/acre MCPA sodium salt (300 g/L).

* Consult manufacturer or seed supplier for varietal tolerances to *metribuzin* and *Prism* tank-mix applications in potato.

Fertilizers: None registered.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *Sencor* or *TriCor* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Allow 5 days between application of *metribuzin* and application of other pesticides.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours may reduce control.

Re-Entry: DO NOT re-enter treated areas for 12 hours after application.

Grazing: DO NOT graze treated cereal crops within 30 days of application, or peas, chickpeas or lentils within 70 days of application.

Preharvest Interval: DO NOT harvest barley, wheat or potatoes within 60 days of application. DO NOT harvest lentils, chickpeas, or field peas within 70 days of application. DO NOT harvest processing peas or chickpeas within 40 days of application.

Re-cropping: Preplant incorporated treatments may leave a residue in the soil that will affect succeeding crops when using higher rates of product. DO NOT seed canola, sunflowers, onions, celery, peppers, cole crops, lettuce, spinach, red beets, turnips, pumpkin, squash, cucumbers or melons the year after treatment. Fall seeded crops may be injured when seeded the same year as preplant or post-emergence applications of these products.

Aerial Application: No restrictions on label. While aerial application is not specifically prohibited, it is not recommended by the manufacturer.

Storage: May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:			
	Aquatic Habitats of Depths			Terrestrial habitat
	Less than 1 m	1 to 3m	Greater than 3 m	
Ground only*	5	2	1	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured is metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Use 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

Keep out of reach of children.

Metsulfuron

Herbicide Group**2 - metsulfuron**

(Refer to page 38)

Company:

E. I. duPont Canada (*Ally* - PCP#24388)

Cheminova Canada (*Accurate* - PCP#29242)

Formulation:

60% metsulfuron methyl formulated as a water dispersible granule.

Container size -

Ally: 122 g package (4 x 30.5 g water soluble bags).

Accurate: 120 g container.

Crops, Rates and Staging:

The following are maximum rates by crop. See the Weeds, Rates and Staging section for weeds controlled by rates less than the maximum.

Cereals - up to 3 g per acre, plus surfactant:

Wheat (spring and durum), barley: 2 leaf up to emergence of the flag leaf.

Established forage grasses for forage or seed production* - up to 3 g per acre, plus surfactant:

Apply from the 2 leaf to flag leaf stage and before canopy is dense enough to prevent thorough leaf coverage.

Crested wheatgrass*

Intermediate wheatgrass*

Creeping red fescue*

Orchardgrass*

Timothy*†

† Fall application with *Ally* only.

Pasture and Rangeland - up to 12 g per acre

* NOTE: - Since applications to forage grasses have been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. **Application to forage grasses is at the risk of the user.**

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:**Cereals and forage grasses:**

Unless otherwise indicated, apply to weeds at the 2 to 4 leaf stage.

Alone or in tank mix with a grass control herbicide use 3 g per acre (One 122 g package treats 40 acres). A rate of 2 to 3 g per acre may be used when mixing with certain other herbicide (See **Tank Mixes**). Add a non-ionic surfactant such as Agral 90, Ag-Surf II, Companion, Super Spreader or Citowett Plus at 0.2 L per 100 L spray volume.

Weeds Controlled:

Ball mustard	Lady's-thumb
Bluebur	Prostrate pigweed
Chickweed	Redroot pigweed
Common groundsel	Scentless chamomile
Corn spurry	Shepherd's-purse
Cow cockle	Stinkweed
Flixweed	Stork's-bill
Green smartweed	Tartary buckwheat
Hemp-nettle	Volunteer canola*
	Wild mustard

* CLEARFIELD varieties will be controlled only with the addition of 2,4-D or MCPA.

Weeds Suppressed:

Canada thistle**	Sow-thistle (annual and perennial)**
Lamb's-quarters♦	Toadflax
Russian thistle♦	Wild buckwheat***

** Apply when thistles are less than 6 inches (15 cm) tall.

*** Apply to wild buckwheat up to the 3 leaf stage.

♦ Apply when weeds are less than 3 inches (8 cm) tall.

Pastures and Rangelands:

Western snowberry - 10 g per acre

Wild rose - 12 g per acre

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:**Water Volume:**

Cereals and forage grasses: Minimum 40 L per acre.

Pastures and Rangelands: 40 to 91 L per acre.

Nozzles and Pressure: No application pressures are recommended by the manufacturer. Typical application pressures for standard flat fan nozzles are from 35 to 40 psi (240 to 275 kPa). Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressures designed to deliver proper coverage with *ASABE medium* when applying to annual crops and *ASABE coarse* droplets when applying to range and pasture. Use a 50 mesh and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Metsulfuron may injure crops stressed by heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations

in day/night temperatures, drought, or water-saturated soils, either before or after application. Weed control will be reduced under dry, cold conditions.

Tank Mixes:

DO NOT mix the soluble bags with any substance containing boron or which releases chlorine.

Herbicides:***In wheat and barley:***

2,4-D Amine or Ester (up to maximum rate for susceptible weeds on 2,4-D page), plus surfactant*.

Avenge (1.72 L/acre), no surfactant required*.

Avenge + MCPA Ester, no surfactant required*.

MCPA Amine or Ester (0.28 to 0.45 L/acre - 500 g/L formulation), plus surfactant.

In creeping red fescue:

Assure II (0.2 to 0.3 L/acre) plus *Sure-Mix* adjuvant*.

Consult tank mix partner labels for additional crop staging and variety restrictions.

* Use with the 3 g per acre rate of metsulfuron only.

Insecticides: None registered.

Fungicides: None registered.

Fertilizer: None registered. DO NOT mix the soluble bags with fertilizers.

Note: The above mixes are those listed on the *Metsulfuron* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rain within 4 hours of application of tank mixes with 2,4-D amine, 2 hours of application of tank mixes with 2,4-D ester, will reduce weed control.

Re-entry: DO NOT enter treated fields for 12 hours.

Grazing: No restrictions.

Re-cropping: Refer to table on the opposite page. The following re-cropping intervals, based on soil pH, should be considered as guidelines only. *Metsulfuron* residues may affect crops for a longer period of time than outlined in the following table. Add 12 months to recommendations if less than 5 inches (130 mm) of rainfall in brown and dark brown soils or less than 10 inches (250 mm) rainfall in black or grey wooded soils in any year following application.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Use	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat**
	Less than 1 m	Greater than 1 m	
Cropland	1	1	15
Pasture and Range use	1	1	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

** Terrestrial buffers are not required for transport and utility rights of way

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack applications do not require a buffer.

Caution:

Metsulfuron residues can persist for long periods, potentially limiting re-cropping options. Degradation of *metsulfuron* is dependent on the pH, moisture, and temperature of the soil. Refer to the label for details on rotation and minimum re-cropping intervals.

MINIMUM RE-CROPPING INTERVAL (MONTHS)

SOIL PH	BARLEY, WHEAT	OAT*	CANOLA*	FLAX*	LENTILS	CANARY-SEED	YELLOW MUSTARD
less than 7.0	10	10	10 (22)	10 (22)	34	48	48
7.0-7.9	10	10 (22)	22 (34)	34	48	48	48

* Figures in brackets refer to re-cropping intervals in brown and dark brown soil zones.

ALL OTHER DATA refer to re-cropping intervals in all soil zones. On black and grey wooded soils with pH of 7.5 or less, fescue may be planted 10 months after application and alfalfa, red clover, peas and flax may be planted 22 months after application. DO NOT use on soils with pH greater than 7.9.

Effects of *metsulfuron* residues on crops other than those listed in the table have not been fully evaluated. Because of the length of re-cropping restrictions and the lack of information on many rotational crops, *metsulfuron* is not recommended for use on farms where special crops are grown (such as fababeans, beans, sunflowers, buckwheat, corn, potatoes, sugar beets, etc.).

Tank Cleaning:

Metsulfuron can cause severe injury to sensitive crops at very low concentrations. Sprayers used to apply *metsulfuron* should be flushed out immediately after *metsulfuron* is used. Refer to 'Method B' found in the general tank cleaning section on page 15 to 16. All nozzles, screens and filters should be removed and cleaned after applying this product.

Hazard Rating:

Caution - Poison



Caution - Eye Irritant

For an explanation of the symbols used here see page 11.

Momentum

Herbicide Group
4 - clopyralid & fluroxypyr
 (Refer to page 38)

Company:

Loveland Products (formerly Viterra) (PCP# 30456)

Formulation:

90 g/L clopyralid and 90 g/L fluroxypyr formulated as an emulsifiable concentrate.

Container size: 8.99 L.

Crops and Staging:

Apply at the 3 leaf to just before the flag leaf stage of barley, wheat (spring, durum).

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

The following weeds are controlled at the 1 to 4 leaf/whorl stage unless specified:

Canada thistle**	Stork's-bill (1-8 Leaf)*
Cleavers	Volunteer flax (1 to 12 cm)
Kochia (2 to 8 leaf)	Wild buckwheat

* Suppression only.

** Season long control, some regrowth may occur in the fall. Apply from the 4 inch (10 cm) to pre-bud stage.

Rates:

0.45 L per acre

Apply a maximum of one application of this product or other products containing either clopyralid or fluroxypyr.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Use 30 to 40 psi (200 to 275 kPa) if applying without drift reduction nozzles. Drift reduction nozzles may require higher pressures for proper performance. Select the nozzle and pressure combination that produces of ASABE coarse droplets while maintaining good coverage of foliage.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

When weeds are stressed because of drought, flooding, hot or cool (less than 8°C) temperatures, weeds are not actively growing and control may be reduced. DO NOT apply to weeds stressed longer than 20 days from lack of moisture as poor control can result.

Tank Mixes:

Herbicides:

MCPA Ester 500 (0.34 to 0.45 L/acre)

MCPA Ester 600 (0.28 to 0.38 L/acre)

Momentum alone or tank mixed with MCPA ester rates above may be mixed with the following:

In spring wheat (including durum) and barley:

Tralkoxydim (0.20 L/acre) plus registered adjuvant

Imazamethabenz (0.52 to 0.64 L/acre) plus water pH adjuster

Fenoxaprop 120 EC (0.16 to 0.31 L/acre).

In spring wheat (not including durum) and barley:

Axial 100EC (0.24 L/acre) plus Adigor adjuvant.

In spring wheat (including durum):

clodinafop (label rates)

Simplicity (0.2 L/acre)

Triaxos (label rate)

† Note: The manufacturer may not support all brand of these products. See the label or contact Loveland Products (formerly Viterra) for more information.

Check the labels of mix partners for additional crop staging restrictions.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the Momentum label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Loveland Products (formerly Viterra) also supports the following mixes that are not on the Momentum label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Everest 2.0, Puma Advance, Varro, Avert, Refine SG and 2,4-D Ester

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfall period is specified on the label; required interval between application and rain without loss of control may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated fields or cut for hay within 3 days of application.

Preharvest Interval: Leave 60 days between application and harvest.

Re-cropping: Wheat, barley, oats, rye, flax, canola, mustard and peas may be planted the year after application or the field may be fallowed. DO NOT under-seed crops to forage legumes the year after treatment. DO NOT sow *any other* crops until the second year after application. Apply manure bedded with straw from treated crops only to the crops listed above.

DO NOT seed to field peas for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field peas grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field peas an additional 12 months (22 months following application). Contact your local Loveland Products (formerly Viterro) representative or retailer for more information before seeding field peas following drought conditions in the previous year.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool (above 5°C), dry area. If product is frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	15	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Spray when winds are under 16 km/hr, but not dead calm.

Tank Cleaning:

No specific cleaning recommendations are provided on the Momentum label. As a petroleum based emulsifiable concentrate, 'Method B' in the general section on tank cleaning on page 15 to 16 may be the most effective. Check with the manufacturer for more information.

Hazard Rating:



Warning – Eye Irritant

For an explanation of the symbols used here see page 11.

Muster Toss-N-Go

Herbicide Group
2 - ethametsulfuron
 (Refer to page 38)

Company:

E. I. duPont Canada (PCP#23569)

Formulation:

75% ethametsulfuron-methyl formulated as a water dispersible granule.

Container size - 320 g package (containing 4 x 80 g water soluble bags).

Crops, Rates and Staging:

NOTE: NOT for use on Yellow mustard (*Brassica alba*).

CROP	RATE (g/acre)	STAGE
Canola	8 to 12	Minimum 2 leaf stage (main stem) to the start of bolting.*
Mustards: Brown & Oriental condiment as well as oilseed quality (<i>Brassica juncea</i>)	8	4 leaf stage but prior to bolting.
Ethiopian Mustard (<i>Brassica carinata</i>)		
Sunflower	8 to 12	2 to 8 leaf stage (15 to 45 cm)

* NOTE: DO NOT apply prior to this stage as severe crop injury can occur.

Muster applied alone requires the addition of Agral 90, Agsurf II, or Citovett at 0.2 L per 100 L of spray solution.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Weeds, Rates and Staging:

Apply from the 2 to 6 leaf stage. Stinkweed must be sprayed in the 1 to 4 leaf stage

At the 8 g per acre rate (one 320 g package treats 40 acres):

Flixweed *	Stinkweed **
Green smartweed	Wild mustard
Hemp-nettle	

The 12 g per acre rate (one 320 g package treats 26.7 acres) controls above weeds plus:

Redroot pigweed **	Stinkweed
--------------------	-----------

* Spring seedlings only.

** Suppression with Muster alone but control with Assure II plus Sure-Mix or a Poast Ultra plus Merge tank mix where permitted.

Muster applied alone requires the addition of Agral 90, Agsurf II, or Citovett at 0.2 L per 100 L of spray solution. Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Equipment, Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets. Use a 50 mesh or coarser screen and filter system. Sprayer must be equipped with continuous agitation. Maintain the spray boom at 24 inches or less above the crop canopy.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT use on crops that are stressed because of drought or flooding. Less than acceptable control will occur in fields where high weed populations exist and where stressful environmental conditions prevail (drought, cold weather). Heavy rainfall soon after application may result in visual crop injury or possible yield reduction. Thin crop stands or application prior to the 2 leaf stage, sandy soils or soils with low organic matter may increase the severity of the injury.

Tank Mixes:

DO NOT mix with substances that contain boron or that release chlorine.

Herbicides:

Canola, Brown and Oriental Mustards (*Brassica juncea* only): *Assure II*. The adjuvant used with *Assure II* is all that is required for this tank mix.

Canola only:

Poast Ultra. The adjuvant used with this product is all that is required for this tank mix.

NOTE: No tank mixes are registered for *Brassica carinata*.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered. DO NOT mix soluble bags with liquid fertilizers.

Note: The above mixes are those listed on the *Muster* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

E.I. duPont also supports the following mixes that are not on the *Muster* label. Mixes must be applied according to the most restrictive use limitations for either product:

Herbicides: *Assure II* plus *Lontrel*, *Lontrel*, *Lontrel* plus *Poast*.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 4 to 6 hours may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze or feed crop to livestock within 60 days of application. DO NOT graze treated sunflowers.

Preharvest: Leave 60 days from application to harvest.

Re-cropping: DO NOT sow wheat, barley, oats or flax within 10 months of application. DO NOT seed canola,

lentils, peas, fababeans, tame mustard, alfalfa, canaryseed, dry beans, fescues or red clover within 22 months of application. All other crops must not be sown until a "field bioassay" is performed at 22 months (or more) after application. Growers may experience reduced yields if other crops (such as corn) are grown without following these guidelines.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones:

CROP (By ground only*)	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Canola, Sunflower, <i>Brassica carinata</i>	4	2	55
Mustard (Condiment and Oilseed types)	3	2	40

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Sprayers used to apply *Muster* should be flushed out immediately after *Muster* is used. Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

None indicated.

Odyssey

Company:

BASF Canada (PCP# 25111)

Formulation:

35% imazamox and 35% imazethapyr formulated as a dispersible granule.

Container size - 8 x 86.5 g water soluble packs per 40 acre case.

Crops and Staging:

Field pea - 1 to 6 above-ground nodes (1 to 6 true leaves).
 CLEARFIELD canola, CLEARFIELD oilseed mustard (*Brassica juncea*), and Fababean - 2 to 6 leaf stage.
 CLEARFIELD lentil - 1 to 9 above ground nodes.
 Soybean - 1 to 3 true leaves.
 Fenugreek (seed and forage production) - 1 to 4 leaf stage.
 Alfalfa and bird's-foot trefoil (seed production only, seedling or established).

Temporary crop yellowing may be observed shortly after application in field pea, and CLEARFIELD canola.

Weeds, Rates and Staging:

At 17.3 g per acre (40 acres per case), Odyssey will control:
 Grasses - 1 to 4 main stem leaves, until tillers are visible:

Barnyard grass	Volunteer cereals (wheat
Green foxtail	excluding CLEARFIELD
Persian darnel	varieties, barley, oats)
	Wild oat

Broadleaf Weeds - cotyledon to 4 leaf stage unless otherwise indicated:

Chickweed	Stinkweed
Cleavers (up to 4 whorls)	Stork's-bill
Flixweed	Volunteer canola (not
Green smartweed	CLEARFIELD varieties)
Hemp-nettle*	Volunteer tame mustard
Lamb's-quarters**	(not CLEARFIELD oil-
Redroot pigweed	seed (<i>B. juncea</i>) varieties)
Russian thistle†	Wild buckwheat*
Shepherd's-purse	Wild mustard

* Suppression only in field peas and CLEARFIELD lentils.

** Suppression only in field peas and CLEARFIELD canola, not controlled in CLEARFIELD lentils.

*** Suppression only.

† Suppression only in CLEARFIELD lentils.

Merge adjuvant (sold separately) must be used with Odyssey.

Herbicide Group

2 - imazamox & imazethapyr

(Refer to page 38)

CLEARFIELD Canola, CLEARFIELD Lentil, Soybean, Alfalfa (seed production only), and Pea: Use only Merge adjuvant at a rate of 0.5 L of Merge in 100 L of spray solution.

DO NOT apply Odyssey more than once or follow Odyssey with any related product in the same year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift. Use 50 mesh or coarser filter screens.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT spray if temperatures of +5°C are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

Herbicides:

In CLEARFIELD canola only:

Lontrel (0.17 to 0.23 L/acre).

In field peas, CLEARFIELD canola, CLEARFIELD oilseed mustard and CLEARFIELD lentils only:

Equinox (67 to 101 mL/acre).

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the Odyssey label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall within 3 hours of application may reduce control.

Re-Entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT graze treated canola or soybean or cut for hay. Field pea may be fed to livestock 30 days after application. DO NOT harvest forage or cut for hay.

Preharvest Interval: DO NOT apply within 60 days of harvesting canola, fababean, oilseed *Brassica juncea*, field pea, and lentil. DO NOT apply within 85 days of harvesting soybean.

Re-cropping: Field pea, lentil, CLEARFIELD canola, canaryseed, oat, barley, field corn, chickpea and spring wheat (including durum) may be seeded the first full season after application. Flax, canola and sunflower may be seeded the second full season after application. The company recommends that a field bio-assay (a test strip grown to maturity) be conducted the year before growing any crops other than those listed above.

NOTE: Breakdown of Odyssey may be slowed or delayed by environmental conditions such as drought, excessive cold and/or acid soils (pH less than 6.5) resulting in an increased risk of injury to rotational crops. The most tolerant crops are CLEARFIELD

canola and legume crops, then cereals. Contact manufacturer for additional information on re-cropping interval (1-877-371-2273).

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Store in a cool, dry place above 5°C.

Buffer Zones: Avoid spraying in situations where drift may occur. Leave a buffer zone of at least 14 m between the outside boundary of the sprayed area and sensitive areas such as shelterbelts, hedgerows, wetlands, woodlots, vegetated ditch banks, ponds, streams, and sloughs.

Tank Cleaning:

Refer to 'Method C' in the general tank cleaning section on page 15.

Hazard Rating:



Warning – Eye and Skin Irritant



Warning – Contains allergen "sulfites"

For an explanation of the symbols used here see page 11.

Odyssey DLX

This product is a prepackaged tank mix of *Equinox* (page 150) and *Odyssey* (page 234). Information listed is restricted to Crop, Weeds, Rates and other important details. Refer to the component products listed above for detailed information on restrictions and the effects of growing conditions.

Herbicide Group

1 - tepraloxymid

2 - imazamox & imazethapyr
(Refer to page 38)

Company:

BASF Canada

Formulation:

The *Odyssey DLX* package contains three components:

Odyssey (PCP# 25111): 35% imazamox and 35% imazethapyr formulated as a dispersible granule.

Container size - 8 x 86.5 g water soluble bags

Equinox (PCP# 27603): 200 g/L tepraloxymid as an emulsifiable concentrate.

Container size - 2.67 L jug.

Merge adjuvant (PCP# 24860):

Container size - 8.1 L jug.

Crops and Staging:

Field peas: 1 to 6 above ground nodes (1 to 6 true leaves).

CLEARFIELD lentil: 1 to 9 above ground nodes.

Temporary crop yellowing may be observed shortly after application in field peas and CLEARFIELD lentils.

CLEARFIELD canola: 2 to 6 leaf stage.

CLEARFIELD oilseed mustard (*Brassica juncea*): 2 to 6 leaf stage.

Weeds and Staging:

Broadleaf and grass weeds controlled at the stages found on the *Odyssey* page plus grasses controlled by *Equinox* at the annual grass rate up to the six leaf stage.

Plus:

Japanese brome (spring seedlings*)

* NOTE: While the label does not indicate the upper limit on what constitutes a seedling, BASF recommends a maximum of 4 leaves at application.

Rate:

Odyssey: 17.4 g per acre.

Equinox: 67 mL per acre.

One case of *Odyssey DLX* will treat 40 acres.

Merge adjuvant: Merge must be used with *Odyssey DLX* at a rate of 0.5 L of Merge in 100 L of spray solution.

Add components in the order listed above.

Odyssey DLX with any related product (*Odyssey*, *Equinox*, *Solo*, *Viper* or imazethapyr) in the same year.

Refer to the product label for complete mixing instructions for this product. A general guide to mixing can be found on page 14.

Restrictions and Application Information:

See the *Equinox* and *Odyssey* components on restrictions application details and handling. Use the most limiting restrictions across all components for the mix.

NOTE: Pay particular attention to the recropping statements on the *Odyssey* page.

Optica Trio

Company:

United Agri Products (PCP#29662)

Formulation:

160 g/L MCPA + 130 g/L mecoprop-p + 310 g/L dichlorprop-p formulated as a solution

Container size - 10 L.

Crops and Staging:

CROP	STAGE
Barley, Oat, Spring wheat (including durum)	2 to 5 leaf
Winter wheat	Spring application only; up to 12 inches (30 cm) high (top leaf extended)

Weeds, Rates and Staging:

Weeds controlled at the 2 to 3 leaf stage unless otherwise indicated.

Apply at 0.61 L per acre to control:

Lamb's-quarters
Stinkweed

Volunteer canola
Wild mustard

Herbicide Group

4 - MCPA, mecoprop & dichlorprop
(Refer to page 38)

Apply at 1.0 L per acre to control the weeds listed above plus:

Canada thistle*
Chickweed (Common)
Cleavers (1 to 2 whorls)
Kochia

Lady's-thumb (suppression)
Ragweed (Common)
Redroot pigweed
Wild buckwheat

* Top Growth Control only.

DO NOT apply Optica Trio more than once or follow application with any related product in the same year.

Application Information:

Water Volume: Minimum 20 L per acre.

Nozzles and Pressure: 30 to 43 psi (200 to 300 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Less than satisfactory control may result if weeds are not actively growing such as under conditions that are extremes of hot or cold, dry or wet weather prior to spraying.

Tank Mixes:

Herbicides:

Spring wheat (including durum)

Signal (93 mL/acre) plus supplied adjuvant.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Optica Trio* label only. To check for other possible registered mixes see the blue chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT feed treated crops to milking animals or harvest for forage within 7 days of application. Meat animals grazing treated crops must be removed 3 day prior to slaughter.

Preharvest: No pre-harvest interval indicated on label when *Optica Trio* is used alone.

Re-cropping: No information provided on label. Contact manufacturer for information.

Aerial Application: DO NOT apply by air.

Storage: Keep from freezing.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	2

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Caution – Poison



Danger – Corrosive to eyes

For an explanation of the symbols used here see page 11.

Option 35 DF/Option 2.25 OD

For use in Manitoba only.

Herbicide Group
2 - foramsulfuron
(Refer to page 38)

Company:

Bayer CropScience

Formulations:

Option 35 DF (PCP# 27425): 35% foramsulfuron formulated as a dispersible granule. Container size - 8 x 100 g packets per case.

Option 2.25 OD (PCP# 27424): 22.5 g/L foramsulfuron formulated as an oil dispersion. Container size - 6.3 L jug.

Crops and Staging:

Field corn at the 1 to 8 leaf stage or 5 to 6 visible collars

Weeds and Staging:

Annual Grasses:

WEED	LEAF STAGE
Barnyard grass	1 to 6 (to early tillering)
Foxtail (green and yellow)	2 to 5 (to early tillering)
Proso millet	2 to 5 (to early tillering)
Witchgrass	2 to 4

Broadleaf Weeds:

WEED	LEAF STAGE
Chickweed, common	4 to 6
Lamb's-quarters	4 to 8
Mustard, wild	5 to 7
Mustard, wormseed	5 to 9
Nightshade, eastern black	1 to 5
Pigweed, redroot	1 to 7
Ragweed, common*	2 to 4

* Suppression only.

Rates:

Option 2.25 OD: 0.63 L per acre (10 acres per jug) plus 28% UAN (liquid 28-0-0) at 1.0 L per acre.

Option 35 DF: 40.5 g per acre (20 acres per case) plus 28% UAN (liquid 28-0-0) at 1.0 L per acre plus *Hasten* Adjuvant at 0.71 L per acre.

NOTE: *Option 35 DF* and *Option 2.25 OD* must be tank mixed with *Banvel II* at 121 mL per acre.

Add *Option 35 DF* or *Option 2.25 OD* to a half full tank, followed by *Banvel II*, then 28% UAN then *Hasten* adjuvant (for use with *Option 35 DF* only).

Application Information:

Water Volume:

Option 35 DF: 89 L per acre.

Option 2.25 OD: 60 L per acre

Nozzles and Pressure: Use 25 to 40 psi (175 to 275 kPa) when using conventional 80° or 110° flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets. Use with 50 mesh or larger screens.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Under optimum conditions weed growth ceases within 1 to 3 days and yellowing of the growing point occurs in 5 to 10 days. Warm moist conditions provide for the best activity. Activity may be reduced or delayed if applied under cool and/or dry conditions or in the presence of heavy dew, fog, mist or rain or if weeds are dust covered. If the crop or weeds are under stress due to environmental conditions, delay application until the both crop and weeds have resumed active growth.

Tank Mixes:

Herbicides:

Barvel II (121 mL/acre)*.

Insecticides: Avoid application to corn that has been treated with organophosphorous insecticides.

Fungicides: None registered.

Fertilizers: DO NOT use any fertilizers or additives other than 28% UAN (1 L/acre), recommended*.

* *Option 35 DF* must be applied to corn in Manitoba as a tank-mixture with *Barvel II*, 28% UAN and *Hasten* adjuvant. *Option 2.25 OD* must be applied to corn in Manitoba as a tank-mixture with *Barvel II* and 28% UAN. See 'Rates' section above.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours may reduce control.

Re-entry: DO NOT enter treated fields until residues have dried.

Grazing: DO NOT graze treated corn crops or cut for forage within 45 days of application.

Preharvest Interval: Leave 70 days between application and harvest of grain.

Re-cropping: The following crops may be grown the season following application: alfalfa, barley, bean (dry common), canola, clover (red), corn (field and sweet), oat, pea, potato, soybean, timothy, spring wheat. Winter wheat may be seeded 4 months after application.

Aerial Application: DO NOT apply by air.

Storage: Keep dry.

Buffer Zones:

Application method (ground only*)	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Option 35 DF	1	1	3
Option 2.25 OD	1	1	3

See page 29 for an explanation of the different habitats.

* Buffer zones for ground applications can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Option residues in the spray tank can cause severe injury to sensitive crops at very low concentrations. Sprayers should be cleaned out immediately before using another product. Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

Option 35 DF:

⚠ Warning – Skin and Eye Irritant.
Potential Skin Sensitizer

Option 2.25 OD:

⚠ Caution – Eye Irritant

⚠ Warning – Skin Irritant
Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Overdrive

Herbicide Group

4 - dicamba

19 - diflufenzopyr

(Refer to page 38)

Company:

Engage Agro on behalf of BASF Canada (PCP# 30065).

Formulation:

20% diflufenzopyr and 50% dicamba sodium salts formulated as water dispersible granules.

Container size - 4 x 3.4 kg.

Crops and Staging:

Established permanent grass pasture, non-cropland sites and rangeland. DO NOT apply *Overdrive* on annual crops or newly seeded grasses.

Weeds and Staging:

Biennial wormwood	Redroot pigweed
Canada thistle*	Sweet clover*
Dandelion**	Tall water-hemp
Lady's-thumb	Velvetleaf
Lamb's-quarters	Vetch*
Leafy spurge**	Wild buckwheat
Ragweed (common)	

* Top growth control

**Top growth suppression

Rates:

115 g per acre. (One package treats 118 acres)

MERGE Adjuvant at the rate of 0.25 L per 100 L of spray solution or a nonionic surfactant at 0.25 L per 100L of spray solution plus ammonium nitrate (UAN 28%) at 1.25L per 100L of spray solution must also be added. Use of an anti-foam agent is suggested.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum 89 L per acre. Use higher water volumes when treating dense or tall vegetation.

Nozzles and Pressure: Maximum 20 psi (150 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of application equipment and pressure that is designed to deliver an even coverage of ASABE coarse droplets that are less prone to drift. Non-target broadleaf plants are very sensitive to *Overdrive* drift. Avoid conditions that are conducive to drift. (See page 12 for drift control suggestions).

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT spray if temperatures are expected to exceed 27°C. DO NOT spray in high humidity or fog. DO NOT spray if wind velocity exceeds 8 km/h. Established grasses growing under stress conditions can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Heavy rain within 4 hours of application may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT permit lactating dairy animals to graze fields within 7 days after application. DO NOT harvest forage or cut hay within 30 days after application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place.

Buffer Zones:

Hand-held or backpack sprayer and spot treatment DO NOT require a buffer zone from sensitive habitat, but efforts should be made to minimize exposure to sensitive plants and open water or wetlands.

Application method	Buffer Zones (metres†) Required for the Protection of:	
	Freshwater habitat	Terrestrial habitat
Field sprayer*	15	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Caution - Poison



Caution - Eye Irritant
Potential Skin Sensitizer



Warning - Contains the allergen sulfites

For an explanation of the symbols used here see page 11.

Pinnacle SG

Herbicide Group
2 - thifensulfuron
(Refer to page 38)

Company:

E. I. duPont Canada (PCP# 22002)

Formulation:

Pinnacle SG (PCP# 29349): 50% thifensulfuron methyl as a water soluble granule.
Container size - 8 x 12 g water soluble pouches.

Crops and Staging:

Soybean - First fully expanded trifoliate leaf to flower initiation.

Weeds, Rates and Staging:

Apply up to 4 inches (10 cm) tall or wide:

Pinnacle SG at 3.3 g per acre will control:

Lady's-thumb	Wild Mustard
Redroot pigweed	

Pinnacle SG at 4.8 g per acre will control the weeds above plus:

Lamb's-quarters	Velvetleaf *
-----------------	--------------

(one container treats 28.5 to 20 acres).

Requires the addition of a non-ionic surfactant such as *Agral 90*, *Ag surf II*, or *Citowett* at 1 L per 1000 L of spray solution. Oil surfactant blends such as *Assist*, or *Sure-Mix* may be used as adjuvants (check label for use rates).

* The addition of 28-0-0 liquid fertilizer at 4 L per 100 L of spray solution or 2.4 kg of 46-0-0 dry urea fertilizer may improve control of velvetleaf. Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 14.

Application Information:

Water Volume: Minimum of 45 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE medium* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Pinnacle applied to crops that have been under stress before application may result in crop injury. Stress conditions within 3 days after application may also result in crop injury.

Weeds under stress conditions at the time of application may not be adequately controlled.

Stress conditions are severe weather conditions, frost, low fertility, drought, water-saturated soils, and disease or insect damage.

Injury symptoms can be crop discoloration (yellowing, purpling or reddening), or stunting.

Tank Mixes:

Herbicides:

Assure II (0.2 L/acre) plus *Sure-Mix**.

Basagran (0.71 or 0.91 L/acre) plus *Assist* adjuvant*.

Basagran Forté (0.71 or 0.91 L/acre)*.

Assure II (0.25 L/acre) plus *Basagran Forté* (0.71 or 0.91 L/acre) plus *Sure-Mix* adjuvant*.

* Refer to appropriate labels for *Pinnacle* and adjuvant rates of application.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Pinnacle* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: If rainfall occurs soon after application control may be reduced. Several hours of dry weather are needed after application to allow uptake by the plants.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze or cut for feed.

Preharvest: Leave 60 days from application to harvest.

Re-cropping Interval: DO NOT plant any crop other than wheat or barley for 30 days after application.

Aerial Application: DO NOT apply by air.

Storage: Store in closed original container in a dry area away from food or feed.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	0	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Pinnacle can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray *Pinnacle* should be flushed out immediately after use. Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Warning: Contains the allergen milk.

For an explanation of the symbols used here see page 11.

Poast Ultra

Herbicide Group
1 - sethoxydim
(Refer to page 38)

Company:

BASF Canada (PCP#24835)

Formulation:

450 g/L sethoxydim formulated as an emulsifiable concentrate.
Container size - 2 x 7.7 L jug of *Poast Ultra*.

Crops, Rates and Staging:

Crops are tolerant at all growth stages. However, the Preharvest interval outlined in the "Restrictions:" sec-

tion must be followed to avoid unacceptable residues of sethoxydim in harvested crops.

To a maximum rate of 0.19 L per acre:
Chickpea

To a maximum rate of 0.23 L per acre:
Tame buckwheat

To a maximum rate of 0.26 L per acre:

Alsike clover**

Caraway

Cicer milkvetch**

Coriander

Dill

Safflower

Sainfoin**

Solin (low linolenic flax)

Sweet clover**

To a maximum rate of 0.45 L per acre:

Alfalfa
Alsike clover*
Canola
Chickling vetch
Cicer milkvetch*

Creeping red fescue (for seed only)
Dry beans (adzuki, kidney, lima, mung, pinto, white)
Dry field peas
Fababeans

Fenugreek
Lentil
Lupin
Mustard
Potatoes
Sainfoin*

* Established stands

** Seedling stands

Flax (not including Solin)
Shelterbelts
Soybeans
Sunflower
Sweet clover*

Weeds, Rates and Staging:

Optimum yield response occurs when weeds are controlled early.

WEEDS AND STAGES	STAGING	RATE (L/ACRE)	ACRES TREATED PER 7.7 L CONTAINER
Green or yellow foxtail, barnyard grass, volunteer corn, Persian dandel, proso millet, witchgrass, large crabgrass	1 to 6 leaf	0.13	60
Wild oats, volunteer wheat, oats and barley	1 to 6 leaf stage except for low rate (See footnote*)	0.13* or 0.19	60 or 40
Quackgrass suppression	1 to 3 leaf stage	0.19	40
Quackgrass (season long control)	1 to 3 leaf stage	0.45	17
Foxtail barley suppression	prior to tillering	0.45	17

* Use the low rate in canola, flax and peas only when

- wild oat, volunteer wheat and volunteer barley are from 1 to 4 leaves (best results prior to tillering)
- under ideal growing conditions (adequate moisture, good fertility and moderate temperatures (15 to 28°C). DO NOT apply under stress conditions.
- with water volumes between 20 to 40 L per acre.

Merge Adjuvant (sold separately): Must always be used with *Poast Ultra*. When *Poast Ultra* is applied alone use *Merge* at 0.5 L to 1.0 L per 100 L of total spray solution. When applying to quackgrass and/or foxtail barley use *Merge* at 1.0 L per 100 L of spray solution. See the tank mix section for *Merge* rates for tank mixing. *Merge* should be added at rates of 0.10 to 0.20 L per acre when applied by air.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre

40 L to 81 L per acre if crop or weed growth is dense, and when spraying quackgrass.

Aerial: 10 to 20 L per acre

Nozzles and Pressure: Use 40 to 45 psi (275 to 300 kPa) with conventional 80° or 110° flat fan nozzles tilted forward at an angle of 45°. Low drift nozzles may require higher pressures for proper performance. Contact the herbicide manufacturer regarding the suitability of low drift nozzles for use with this product. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Most effective control is achieved when grasses are actively growing. Weeds stressed by drought, flooding, hot or prolonged cool temperatures (<15°C) and poor fertility are more difficult to control. Use the higher of the recommended rates for grasses stressed for less than 20 days. DO NOT apply to grasses stressed more than 20 days because of lack of moisture. Control may be reduced if temperatures are below 15°C. Subsequent tillering may occur under stress conditions or if fertility is low.

Tank Mixes:

Herbicides: The following tank mixes can be applied with 0.13 to 0.19 L/acre of *Poast Ultra*.

Merge Adjuvant (sold separately): Use at 0.75 to 1.0 L of *Merge* per 100 L of mixed spray solution for most mixes except when mixing with *Pursuit* use 1.0 L per 100 L of solution.

In Flax:

Buctril M (0.4 L/acre)(including Solin).

Logic M (0.5 L/acre)(including Solin).

Lontrel 360 (0.23 to 0.34 L/acre).

Lontrel 360 (0.23 to 0.34 L/acre) + *MCPA Ester* (0.34 to 0.45 L/acre - 500 g/L formulations).

MCPA Ester (up to 0.45 L/acre - 500 g/L formulations)

The above tank mixes may reduce grass control, especially under adverse weather conditions.

In Canola:

Lontrel (0.17 to 0.34 L/acre).

Muster (8 to 12 g/acre).

Lontrel (0.17 L/acre) + *Muster* (8 g/acre) + *Merge* (0.4 L/acre).

In Liberty Link Canola only:

Poast Ultra (0.90 L/acre) + *Liberty* (1.08 L/acre)

In Field Pea:

Poast Ultra (0.19 L/acre) plus *Merge* (0.4 L/acre) may be tank mixed with:

Pursuit (40 mL/acre) to control:

Chickweed

Cleavers

Hemp-nettle (peas only)

Redroot pigweed (light infestations only)

Smartweed

Stinkweed

Volunteer canola (non-

CLEARFIELD varieties)

Wild buckwheat (light infestations only)

Wild mustard

The company does not provide guidelines for weed densities under light infestations. When in doubt, use the higher rate below or contact the manufacturer.

Pursuit (85 mL/acre) for all weeds on the *Pursuit* label.

Check label directions for mixing order and additional timing restrictions for broadleaf partners.

Allow 4 days between application of *Poast Ultra* and application of herbicides other than those registered for tank mixing. Allow 5 days between application of *Sencor* and *Poast Ultra*. Allow 14 days for regrowth when applied in sequence with a grass control herbicide.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Poast Ultra* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Re-Entry: DO NOT enter treated field for 12 hours.

Grazing: DO NOT graze the treated crop or cut for feed prior to crop maturity. Forage legumes may be cut after the specified Preharvest interval.

PREHARVEST INTERVAL (DAYS)	CROPS
30	Forage legumes (excluding alfalfa)
60	Dry peas, fenugreek, flax
65	Lentil, chickpea
70	Canola, chickling vetch, alfalfa
76	Mustard
80	Potato, dry bean, soybean, fababean, lupin
85	Buckwheat
86	Solin
90	Safflower
105	Sunflower

Re-cropping: DO NOT plant cereals or grass within 14 days of application.

Aerial Application: May be applied by air.

Storage: May be frozen.

Buffer Zones:

Application method	Crops	Buffer Zones (metres [†]) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
Ground*	All	1	0	2
Fixed wing airplane	Food or feed crops	1	0	7
	Shelter- belts	5	0	150
Helicopter	Food or feed crops	1	0	60
	Shelter- belts	1	0	85

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance from downwind edge of spray boom and non-target area.

Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15 to 16. Empty and clean spray tank using this method if an oil film accumulates.

Hazard Rating:



Caution - Poison



Caution - Eye and Skin Irritant

For an explanation of the symbols used here see page 11.

PrePass XC (this referring entry will remain until the 2016 edition)

See florasulam + glyphosate on page 160.

Prestige XC

Herbicide Group
4 - fluroxypyr, clopyralid & MCPA
(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

The *Prestige XC* package has 2 components:

Prestige XC A (PCP# 29462): 333 g a.e./L fluroxypyr
Container size - 3.3 L or in bulk package 4 x (2 x 9.9 L)

Prestige XC B (PCP# 29465): 50 g/L clopyralid and 280 g/L MCPA ester.

Container size - 2 x 8.0 L or bulk package 4 x 96 liter.

All of the above components are formulated as an emulsifiable concentrates.

Crops and Staging:

Cereals:

Spring wheat (including durum), winter wheat (apply in the spring), barley, oat and canaryseed*

Forage Grasses* grown for seed production:

Seedling and established stands - 4 leaf until the emergence of the flag leaf.

Bromegrass (meadow, smooth)	Wheatgrass (crested, intermediate)
Fescue (creeping red, tall)	
Timothy	

* NOTE: Since these uses are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Users of this product on forage grasses and canary seed do so at their own risk.

Weeds, Rates and Staging:

Unless otherwise stated, the following weeds will be controlled if sprayed in the 2 to 4 leaf stage.

Prestige XC A at 0.13 L per acre; *Prestige XC B* at 0.6 L per acre (one case treats 27 acres or bulk treat 640 acres) controls:

Burdock	Ragweeds
Canada thistle (light infestations)	Shepherd's-purse
Cleavers (1-8 whorls)	Stinkweed
Field horsetail†	Stork's-bill (1-8 leaf)
Flixweed (spring seedlings only)	Vetch
Kochia	Volunteer flax (1-12 cm)
Lamb's-quarters	Volunteer sunflower
Plantain†	Wild annual sunflower
Prickly lettuce	Wild buckwheat (1-8 leaf)
	Wild mustard
	Wild radish

The 480 acre per bulk container rate of *Prestige XC* or the 20 acre per case rate (*Prestige XC A* at 0.17 L per acre; *Prestige XC B* at 0.8 L per acre) controls the above weeds plus:

Annual sow-thistle	Redroot pigweed
Canada thistle* (moderate to heavy infestations)	Round-leaved mallow (1 to 6 leaf)
Chickweed (up to 6 cm)	Russian pigweed
Common groundsel	Scentless chamomile
Dandelion**	Smartweed
Hemp-nettle (2-6 leaf stage)	Tartary buckwheat
Perennial sow-thistle*	Volunteer canola

* Spray when 4 to 6 inches (10 to 15 cm) high. Season long control, with some regrowth in the fall.

** Spring rosettes only.

† Top growth control only.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre.

Aerial: 12 to 20 L per acre. Consult label for buffer zones.

Nozzles and Pressure: Maximum 30 to 40 psi (200 to 275 kPa) with conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver proper coverage with ASABE coarse droplets. Tilt nozzles forward at a 45° angle to improve coverage of vertical targets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

The activity of the *Prestige* is influenced by weather conditions. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions (drought or heat stress) or if heavy infestations exist.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Liquid Achieve (0.2 L/acre) plus *Turbocharge* adjuvant.

Assert (0.53 to 0.65 L/acre) plus acidifier.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Prestige* XC label only. For other possible registered mixes see the blue fold out chart inside back cover.

Dow AgroSciences also supports the following mixes that are not on the *Prestige* XC label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Avenge*

Adding ingredients in the correct order is critical for optimum performance.

Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours of post-emergent application may result in reduced weed control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT cut or graze treated fields of wheat, barley or canaryseed for 7 days after application. DO NOT cut treated forage grass fields for hay or forage. DO NOT graze treated forage grass fields.

Preharvest Interval: DO NOT harvest crop within 60 days of application.

Re-cropping: Wheat, oat, barley, rye (not under-seeded to forage legumes, clover or alfalfa), flax, canola, field pea* and mustard may be seeded the season following application.

*NOTE: DO NOT seed to field pea for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field pea grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field pea an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field peas following drought conditions in the previous year.

DO NOT seed legume forages or crops other than those listed above until the second season following application.

Aerial Application: May be applied by air.

Storage: Store product in original containers in a secure, dry, heated area. If the product is frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†)		
	Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	0	1
Fixed Wing aircraft	4	0	60
Helicopter	1	0	50

See page 29 for an explanation of the different habitats.

* Buffer zones for ground applications can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method C' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Danger - Poison.



Warning - Eye and Skin Irritant

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Primextra II Magnum

Herbicide Group
15 - metolachlor
5 - atrazine
(Refer to page 38)

Company:

Syngenta Canada (PCP#25730)

Formulation:

400 g/L of s-metolachlor + 320 g/L of atrazine formulated as a liquid.

Container size - 14 L.

Crops and Staging:

Corn - Preplant incorporated (PPI) or Pre-emergence if irrigated within 10 days of application.

Weeds and Staging:

Apply prior to the emergence of weeds. Weeds that have emerged prior to application will not be controlled.

American nightshade	Purslane
Barnyard grass	Ragweed
Buckwheat	Redroot pigweed
Eastern black nightshade	Smartweed (lady's-thumb)
Foxtail (green, yellow)	Wild mustard
Lamb's-quarters	Witchgrass
Prostrate pigweed	Yellow nutsedge*

* Herbicide must be incorporated for best control.

Rates:

WEED POPULATIONS	RATE (L/ACRE)	ACRES TREATED PER 14 L CONTAINER
Light infestations	1.2	11.7
Medium infestations	1.4	10
Heavy infestations	1.6	8.8

DO NOT apply to soils with less than 1% or more than 10% organic matter.

Application Information:

Water Volume: 61 L per acre

Pressure: 30 to 45 psi (200 to 300 kPa).

Nozzles: Flat fan with 50-mesh nozzle screens.

Incorporation:

Incorporate using S-tine or C-tine cultivators or tandem disk. Do not incorporate deeper than 4 inches (10 cm).

To ensure that the product remains in the top 2 inches (5 cm) of soil, apply to a firm seedbed free of large clods or lumps. If using tandem disks, set disks to work the soil at a depth of 4 inches (10 cm) and operate at a speed of 4 mph (6 km/hr). If using an S-tine cultivator, set the implement to work the soil to a depth of 4 inches (10 cm) and operate at a speed of 6 mph (10 km/hr).

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Extended periods of dry soil conditions may result in reduced weed control. Moderate rainfall after application will enhance activity. Heavy rainfall following application of *Primextra II Magnum* may dilute the metolachlor deeper than 2 inches (5 cm) and result in reduced weed control, particularly on light textured soils.

Tank Mixes:

Herbicides: None registered.

Insecticides: None registered.

Fertilizers: May be tank mixed with liquid fertilizer for preplant incorporated applications. Conduct a compatibility test by performing a jar test prior to mixing the products in the tank. May be impregnated onto dry bulk fertilizers (except nitrate or superphosphate fertilizers or limestone).

Note: The above mixes are those listed on the *Primextra* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Moderate rainfall shortly after application will enhance activity. Heavy rainfall reduces weed control by leaching the chemical out of the top few centimeters of soil. Inadequate rainfall after application (within 10 days) will cause reduced weed control.

Re-entry: DO NOT re-enter treatment area within 12 hours of application.

Grazing: DO NOT graze or cut corn for feed before ear emergence.

Re-cropping: This product contains *Atrazine*. All crops except corn and triazine-tolerant canola may be affected the year following the use of *Atrazine*. Other more sensitive crops may be affected two or more growing seasons after application.

Aerial Application: DO NOT apply by air.

Storage: Store in a dry place.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic habitat	Terrestrial habitat
Ground only*	29	10

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

DO NOT mix or load this product within 30 m of any sensitive aquatic habitats

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Ratings



Caution Poison



Caution - Eye Irritant

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Prism SG

Herbicide Group

2 - rimsulfuron

(Refer to page 38)

Company:

E. I. duPont Canada Inc. (PCP#30057)

Formulation:

25% rimsulfuron formulated as a water soluble granule.

Container size - 480 g.

Crops and Staging:

Irrigated potato* prior to flower initiation.

Potato tolerance differs by variety. Limit first use to a small area of each variety prior to widespread adoption in the field. Delay cultivation for 7 to 10 days after application.

* NOTE - Since applications to irrigated potato in western Canada has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to irrigated potato in western Canada is at the risk of the user.

Weeds and Staging:

WEED	STAGE
Barnyard grass Green foxtail Yellow foxtail Witchgrass	1 to 6 leaf stage, maximum 2 tillers
Quackgrass	3 to 6 leaf stage (less than 10 inches or 25 cm leaf extended).
Redroot pigweed Lamb's-quarters (suppression only)	4 to 6 leaf stage (less than 4 inches or 10 cm tall or across).

Rates:

24 g per acre (one package treats 20 acres).

Add a recommended non-ionic surfactant such as *Citowett Plus*, *Agsurf II*, or *Agral 90* at 0.2 L per 100 L spray solution.

Make only one application per growing season.

Refer to the product label for complete mixing instructions for this product and its mixes.

A general guide to mixing can be found on page 14.

Application Information:

Water Volume: Minimum 40 L per acre.

Nozzles and Pressure: 25 to 40 psi (175 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE medium* droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Apply when the temperature 24 hours before and after application is between 5°C and 28°C. Temperatures beyond this range increase the potential for crop injury. Rapid fluctuations in temperature will stress the crop (greater than a 20°C difference within 24 to 36 hours). Allow 48 to 72 hours for the crop to acclimatize before spraying if severe temperature fluctuations occur. Crop injury may result if applications are made when potatoes are stressed by abnormally hot, humid, or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage. If potatoes have been injured by frost, wait 48 to 72 hours after normal growing conditions have resumed before applying.

Warm, moist conditions after application promote good weed control while cool and/or dry conditions may reduce or delay activity. Weeds hardened off by cold weather or drought stress may not be controlled.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Within 2 to 4 hours may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Preharvest: Leave 30 days from application to harvest.

Grazing: DO NOT graze the treated crop or cut for hay.

Re-cropping: Spring barley, soybean, white bean, red clover, sorghum, potato and field corn may be planted the year after application. Winter wheat may be planted 4 months after application. For all other crops, a field bioassay is recommended before planting.

Aerial application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

DO NOT apply in areas where surface water from the treated area can run off to adjacent cropland, streams, irrigation water or wells.

Tank Cleaning:

Refer to 'Method A' found in the general tank cleaning section on page 15 to 16. Check the label or contact the manufacturer for more specific tank cleaning information.

Hazard Rating:



Warning – Eye Irritant

Warning – Contains the allergen sodium sulfite.

For an explanation of the symbols used here see page 11.

Pulsar

Herbicide Group
4 - dicamba & fluroxypyr
 (Refer to page 38)

Company:

Syngenta Canada

Formulation:

86.9 g/L dicamba and 113.3 g/L fluroxypyr formulated as an emulsifiable concentrate.

Container size - 2 x 9.82 L or 78.6 L

Crops and Staging:

Barley and spring wheat (including durum) from the 2 to 5 leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Unless otherwise indicated apply when weeds are at the 2 to 3 leaf stage and rosettes are less than 2 inches (5 cm) across.

At 246 mL per acre (80 acre per case) *Pulsar* controls:

Cleavers	Wild buckwheat*
Kochia (up to 9-leaf)	

At 371 mL/acre (53 acre per case) *Pulsar* controls the weeds above plus:

Lamb's-quarters*	Stork's bill*
Redroot pigweed*	Volunteer flax
Russian thistle	Wild buckwheat (up to 9-leaf)

* Suppression only

Application Information:

Water Volume: Minimum 44.5 L per acre.

Nozzles and Pressure: For conventional flat fan nozzles use a maximum pressure between 40 and 45 PSI (275 to 310 kPa) for the preformulated product. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium droplets*.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury or reduced weed control may result.

Tank Mixes:

Herbicides:

Barley, Spring wheat, and durum only:

MCPA LV600 ester (0.23 L/acre)

Spring wheat, and durum:

Horizon NG (376 mL/acre)

Horizon NG (376 mL/acre) plus MCPA LV600 ester (0.23 L/acre)

Traxos (label rate)

Traxos (label rate) + MCPA Ester (rates above)

Fertilizers: None registered

Note: The above mixes are those listed on the *Pulsar* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Syngenta also supports the following mixes that are not on the *Pulsar* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides:

2,4-D Ester, Express SG, Refine SG, Sierra 2.0 (flucarbazone).

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-Entry: DO NOT enter treated fields for at least 12 hours.

Preharvest: Leave 60 days between treatment and harvest.

Grazing: Treated crops may be grazed, or cut for hay or silage after 7 days when used alone, or a minimum of 12 days when mixed or longer if the intervals are longer for the tank mix partner.

Re-cropping: Wheat, barley, oats, rye, forage grasses, flax, canola, mustard, lentils and peas may be grown the following season. There are no recropping restrictions the second year after application.

Aerial Application: DO NOT apply by air.

Storage: May be frozen. If frozen, bring to room temperature and agitate before use. This product is combustible. DO NOT store near heat or open flame.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic habitat	Terrestrial habitat
Ground only*	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Legumes are particularly sensitive to *Pulsar*.

Handheld or backpack applications do not require a buffer.

Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Danger – Poison



Warning – Eye Irritant



Caution – Skin Irritant.

For an explanation of the symbols used here see page 11.

Quizalofop

**Herbicide Group
1 - quizalofop**
(Refer to page 38)

Company:

E. I. duPont Canada (*Assure II* - PCP# 25462)

Gowan Canada (*Yuma GL* - PCP# 30100)

Formulation:

96 g/L quizalofop-P-ethyl formulated as an emulsifiable concentrate.

Container size:

Assure II - 8 L *Assure II* + 8 L *SureMix* adjuvant, or 1000 L tote.
Yuma GL - 8 L *Yuma GL* + 8 L *XA Oil Concentrate*.

Crops and Staging:

No leaf stage restrictions, but do not apply beyond Preharvest intervals listed in the table:

CROP	PREHARVEST INTERVAL (DAYS)
Camelina*††♦♦	64
Canola	64
Chickpea	85
Dry Edible Bean*†♦♦	30
Ethiopian Mustard (<i>Brassica carinata</i>)*††	64
Faba Bean*††♦	30
Flax, or Solin (low linolenic acid flax)	82
Lentil	65
Oriental mustard (condiment types and oilseed quality types <i>Brassica juncea</i>)	64
Pea (field and processing)	65
Soybean	80
Sunflower*††	60
Yellow and Brown Mustard††	64

† NOTE: While *Quizalofop* has been registered for use on all dry field bean types (except faba bean) not all types have

been tested for tolerance. When using *Quizalofop* on a new dry bean type or variety for the first time evaluate tolerance on a small area first before applying large acreages and check with seed supplier for variety sensitivity.

Forage Crops (no Preharvest interval restrictions):

Alfalfa (seed production)

All seedling legume forage species for seed production*♦♦

Creeping red fescue for seed (seedling or established)*.
Established Alsike and Red Clover for seed production only††♦

†† *Assure II* only.

* NOTE - Since applications to these crops have been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to these crops is at the risk of the user.

♦ If using *Assure II* with *Merge* adjuvant use low rate only.

♦♦ If using *Assure II* use with *SureMix* adjuvant only.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Add one of the following registered adjuvants to the spray tank when applying *quizalofop*:

SureMix (0.5 L per 100 L of spray solution)**

Merge (0.5 to 1.0 L per 100 L of spray solution)

XA Oil Concentrate (0.5 to 1.0 L per 100 L of spray solution)***

Use the higher rate of *Merge* or *XA Oil Concentrate* when wild oats or quackgrass are present in the field or when growing conditions are poor.

** *SureMix* is packaged with small sizes of *Assure II*.

*** Use *XA Oil Concentrate* with *Yuma GL* only.

WEED	STAGE	RATE	
		L/ACRE	ACRES PER 8 L
Wild oat*	1 to 5 leaf prior to tillering	0.15	54
	up to 2 tillers	0.20	40
Green foxtail	2 leaf to early tillering	0.15	54
Volunteer wheat, barley & oat*	2 leaf to early tillering	0.15	54
Volunteer corn	2 to 6 leaf stage	0.15	54
Barnyard grass, yellow foxtail, proso millet, old witchgrass	2 leaf to early tillering	0.20	40
Quack grass suppression	2 to 6 leaf stage		
Foxtail barley†	3 to 4 leaf max 3 tillers		
Quack grass season long control	2 to 6 leaf stage	0.30	26

* Best results are likely to occur if applications are made before tillering begins. Apply at the 2 to 3 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering.

† Control with *Assure II* only and suppression with *Yuma GL*. Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

Application Information:

Water Volume:

Ground: Minimum 40 L per acre. Up to 162 L per acre of water may be used under heavy populations to improve coverage.

Aerial: Minimum 10 L per acre to a maximum of 20 L per acre.

Nozzles and Pressure: 30 to 40 psi (210 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium* droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop injury may occur if crops are stressed because of drought or flooding. Less than acceptable weed control may be expected if weeds are under stress because of drought, flooding or cool weather.

Tank Mixes:

Herbicides:

In Canola: *Muster* (8 to 12 g/acre).

In Dry Bean (Pinto, Pink, Great Northern and Small Red): *Basagran* (label rates with *Quizalofop* at 0.25 L/acre plus *SureMix* adjuvant)

In Oriental Mustard (B. juncea condiment and oilseed): *Quizalofop* (0.15 to 2.0 L/acre) plus *Muster* (8 g/acre) plus *Merge* or *SureMix* adjuvants. DO NOT use on yellow mustard as injury will result.

In Soybean:

Pinnacle (2.2 to 3.3 g/acre).

Quizalofop (0.25 L/acre) plus *Pinnacle* (2.2 to 3.3 g/acre) plus *Basagran Forté* (0.71 to 0.91 L/acre) plus *Merge* or *SureMix* to control grasses controlled by *Quizalofop* alone at 0.20 L/acre plus the broadleaf weeds controlled by *Pinnacle* and *Basagran Forté*.

In Established creeping red fescue for seed: *Quizalofop* at 0.2 to 0.3 L/acre may be tank mixed with *Ally* (3 g/acre). Allow 24 hours after application before applying a broadleaf herbicide. If the broadleaf herbicide is applied first, wait 7 days before application of *Quizalofop*.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Quizalofop* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Quizalofop manufacturers may also support mixes with pesticides that are not on the *quizalofop* labels. Check with each manufacturer for the products they support. Mixes must be applied according to the most restrictive use limitations for all products added to the tank.

Herbicides: *Ares*, *Buctril M**, *Curtail M**, *Eclipse**, *Glyphosate*, *Liberty*, *Lontrel**, *Lontrel+Muster**, *Odyssey*, *Pursuit*, *Solo**.

**Assure II* only.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour of application may reduce control.

Grazing: DO NOT graze treated crops or cut for feed in the year of treatment.

Preharvest Interval: See 'Crops and Staging' chart above.

Re-cropping: No restrictions the year after treatment.

Aerial Application: May be applied by air when used alone.

Storage: DO NOT freeze.

Buffer Zones:

Rates (L/acre)	Application method	Buffer Zones (metres†) Required for the Protection of:		
		Aquatic Habitats of Depths		Terrestrial habitat
		Less than 1 m	Greater than 1 m	
All rates	Ground *	1	0	3
Up to 0.15	Winged aircraft	0	0	70
	Helicopter	0	0	55
Up to 0.20	Winged aircraft	0	0	85
	Helicopter	0	0	70
Up to 0.30	Winged aircraft	1	0	125
	Helicopter	1	0	100

See page 29 for an explanation of the different habitats.

* Buffer zones for ground applications can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Cleanout is recommended but no specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Danger – Corrosive to eyes



Skin irritant, Potential skin sensitizer

For an explanation of the symbols used here see page 11.

Reclaim

Herbicide Group

2 - metsulfuron

4 - aminopyralid & 2,4-D

(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

Reclaim A (PCP# 29751): 52.5% aminopyralid + 9.45% metsulfuron methyl formulated as a water dispersible granule. Container size - 1.84 kg

Reclaim B (PCP# 29750): 564 g/L 2, 4-D Ester formulated as an emulsifiable concentrate.

Container size - 2 x 8 L

NOTE: Limited availability through selected retail outlets.

Crops and Staging:

Rangeland and pastures - Apply in spring or early summer.

Weeds, Rates and Staging:

Apply when weeds are young and actively growing in the vegetative stage.

Reclaim A: Apply at 92 g per acre.

Reclaim B: Apply at 0.8 L per acre.

One case treats 20 acres.

Weeds Controlled:

The following broadleaf weeds, invasive plants and shrubs will be controlled for 24 months after application.

Canada Thistle

Prairie sage

Dandelion

Wild Rose

Wolf willow (Silver-berry)

Shrubby cinquefoil

Buckbrush (western snow-berry)

Pasture sage (fringed sage)

Application Information:

Water Volume:

Ground application: 45 L per acre minimum. For better coverage apply at 80 L per acre.

Aerial application: 20 L per acre minimum

Nozzles and Pressure: Use a combination of application equipment and pressures that will apply *ASABE coarse* droplets in a uniform pattern. Drift of even small amounts of *Reclaim* into sensitive plants or areas where sensitive crops may be grown can cause injury. DO NOT apply under conditions prone to drift (i.e. high winds, dead calm, or temperature inversions).

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Application should be avoided when pasture and targeted weeds are under stress of drought, excess moisture, extreme heat or cold or other environmental stresses. Target weeds must be actively growing. Avoid applications when temperatures exceed 28°C.

Tank Mixes:

None registered

Restrictions:

Rainfall: No rainfast period is specified on the label. Contact manufacturer for more information.

Re-entry: DO NOT re-enter treated areas for 12 hours.

Grazing: DO NOT allow lactating dairy animals to graze treated areas within 7 days of application. Withdraw meat animals from treated areas and feed untreated feed for at least 3 days before slaughter. DO NOT harvest forage or cut hay within 30 days of application.

Re-cropping: DO NOT apply to pastures where legumes are an essential component. DO NOT break up treated pasture and plant to sensitive broadleaf crops for at least 3 years after application.

Aerial Application: May be applied by air.

Storage: Store product in original, labeled containers in a secure, dry, cool area. DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic habitat	Terrestrial habitat
Ground*	10	15
Fixed wing airplane	80 to 175**	250 to 750**
Helicopter	70 to 150**	175 to 650**

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

** Distance varies depending on spray droplet size. Consult the *Reclaim* label to determine buffer zone size when applying by air.

† Distance is measured from the downwind edge of the boom to sensitive areas.

DO NOT apply this product directly to any water body or mix or load near water or wells. DO NOT apply when heavy rains are forecast or on moderate to steep slopes toward sensitive areas or to light soils with shallow water table. Contact the provincial environment department for permits to apply near water.

Tank Cleaning:

Refer to 'Method A' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product such as All Clear or Clean Out spray cleaner. The inclusion of detergent in 'Method B' may provide improved cleaning. Contact the manufacturer for more information.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Reglone

(this referring entry will remain until the 2015 edition)

See diquat on page 134.

Reflex*

Herbicide Group

14 - fomesafen

(Refer to page 38)

*(For use only in the Red River Valley of Manitoba)

Company:

Syngenta Canada (PCP#24779)

Formulation:

240 g/L fomesafen formulated as a solution.

Container size - 10 L.

Crops and Staging:

Soybeans and dry beans[♦] in the 1 to 2 trifoliate leaf stage. DO NOT use before the 1st trifoliate leaf stage or increased risk of crop injury may result. For use in the Red River Valley of Manitoba only.

[♦] NOTE - Since applications to beans in the Red River Valley has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to beans is at the risk of the user.

Weeds and Staging:

Broadleaf weeds controlled by Basagran at the 0.71 L per acre rate plus improved control of the following weeds up to the 4-leaf stage:

Cocklebur	Ragweed (common)
Eastern black nightshade	Redroot pigweed*
Lady's-thumb	Volunteer canola
Lamb's-quarters	Wild mustard

* Suppression only

Rate:

Reflex is registered in the Red River Valley of Manitoba only as a tank mix at a rate of 235 mL per acre Reflex plus 0.71 L per acre Basagran plus Agral 90 at 1 L per 1000 L of spray solution.

Application Information:

Water Volume: Minimum 81 L per acre. Increase water volume to 142 L per acre for fields with heavy weed densities or with weeds at the upper limit of their recommended stage.

Pressure: 275 kPa (40 psi). Increase pressure to 420 kPa (60 psi) for fields with heavy weed densities or with weeds at the upper limit of their recommended stage.

Nozzles: Use nozzles capable of delivering appropriate pressures and volumes.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Weed control and crop tolerance may be reduced under certain stress conditions such as cold temperatures, excess moisture, drought and injury from hail or previous herbicide applications.

Tank Mixes:

Herbicides: Reflex is only registered for use in a Basagran tank mix. See Rates.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Grazing: DO NOT graze treated crop or cut for hay. There is insufficient data to support such use.

Preharvest: Leave at least 84 days from application to harvest.

Re-cropping: Winter wheat may be sown 4 months after application. Spring wheat, dry beans, soybeans and field corn may be grown the year following an application.

DO NOT apply Reflex to any field more often than once every 2 years.

These re-cropping restrictions refer only to the Red River Valley of Manitoba. Use outside this region is not registered as re-cropping options have not been determined.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool place away from food or feed.

Buffer Zones: Leave a buffer zone of at least 15 m between the last spray swath and the edge of sensitive terrestrial areas such as shelterbelts, hedgerows and shrublands as well as aquatic areas such as ponds, streams, rivers, prairie potholes and sloughs. DO NOT apply when winds are greater than 15 km/hr.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Danger - Corrosive to Eyes

For an explanation of the symbols used here see page 11.

Restore*/Restore II

Herbicide Group
4 - aminopyralid & 2,4-D
(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

The *Restore** package contains two components:

Restore A (PCP# 28551): 240 g/L aminopyralid formulated as a solution. Container size - 3 L, or 450 L bulk.

Restore B (PCP# 28552): 564 g/L 2,4-D formulated as dimethylamine solution. Container size - 2 x 7.65 L or 450 L bulk -or-

Restore II (PCP# 30632): 40 g/L aminopyralid and 400 g/L 2,4-D both present as amine salts formulated as a solution. Container size - 2 x 10 L

Note: Limited availability through selected retail outlets.

* NOTE: This product is no longer manufactured, but some supplies may still remain in the distribution system. This product may be removed from future editions of this publication.

Use a maximum of one application of *Restore* products or other products containing aminopyralid per season.

Crops and Staging:

Rangeland and pastures - Apply in spring or early summer.

Weeds, Rates and Staging:

Apply when weeds are young and actively growing in the vegetative stage for control unless indicated otherwise.

Restore A at 0.11 L/acre plus Restore B at 0.6 L/acre or Restore II at 0.57 L/acre will control:

Annual sow-thistle	Gumweed†*
Blue lettuce†*	Hoary cress†*
Bull thistle	Horse-nettle**
Burdock (<4 leaf)	Nodding thistle**
Buttercup (hairy, tall)	Ox-eye daisy**
Canada fleabane**	Perennial sow-thistle
Canada thistle††**	Prickly lettuce
Canada goldenrod††*	Scentless chamomile*
Common broomweed**	Spotted knapweed (††*)
Common plantain	Stinging nettle
Curled dock (<4 leaf)*	Sweet clover
Daisy fleabane**	Yellow star-thistle**
Goat's beard	

As well as other annual "Susceptible Weeds" controlled by 2,4-D on the 2,4-D page.

Restore A at 0.15 L plus Restore B at 0.77 L or Restore II at 0.86 L/acre will control:

Weeds listed above plus:

Absinthe††*	Hawkweed**
Canada goldenrod††	Hairy fleabane**
Cudweed**	Heal all**
Curled dock (<4 leaf)	Scentless chamomile
Dandelion††	Sulphur cinquefoil††**
Dog mustard**	Tansy ragwort**
Groundsel**	Western ragweed**

As well as other annual "Hard to Control Weeds" controlled by 2,4-D on the 2,4-D page.

Restore A at 0.2 L/acre plus Restore B at 1 L/acre or Restore II at 0.97 L/acre will control:

Weeds listed above plus:

Absinthe	Fuller's teasel**
Biennial wormwood**	Gumweed††**
Bindweed (Field and Hedge)††**	Hoary cress††**
Blue Lettuce††**	Ironweed (tall)**
Burdock**	Mouse-eared chickweed††**
Common tansy††*	Russian snowweed†††**
Common tansy †††**	Western snowberry (buck brush)††*
Dandelion††**	Yarrow (common)†††**
Diffuse knapweed†††**	Yellow rocket††**

* *Restore* only.

** *Restore II* only.

† Top growth control only.

†† Season long control.

††† Suppression only.

Application Information:

Water volume:

Ground application: 40 L per acre minimum.

Aerial application: 20 L per acre minimum

Nozzles and Pressure: Use a combination of application equipment and pressures that will apply *ASABE coarse* droplets in a uniform pattern.

Drift of even small amounts of *Restore* into sensitive plants or areas where sensitive crops may be grown can cause injury. DO NOT apply under conditions prone to drift (i.e. high winds, dead calm, or temperature inversions).

Avoid applications closer than the drip line or outer edge of the canopies of trees or injury may occur to the tree.

NOTE: Use closed handling systems when using bulk containers and/or if handling more than 663 L of product per day. Handheld

applications are limited to 20 L. of product per day. Respirators must be worn if applying more than 12.5 L. per day using hand-held equipment.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Application should be avoided under conditions of drought or other environmental stress.

Tank Mixes:

None registered.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT re-enter treated areas for 12 hours.

Grazing: DO NOT allow lactating dairy animals to graze treated areas within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter. DO NOT harvest forage or cut hay within 30 days of application. Allow 3 days of grazing on an untreated pasture (or feed untreated hay) before transferring livestock to areas where sensitive broadleaf crops may be grown.

Re-cropping: If legumes are essential in a pasture, DO NOT use *Restore*. DO NOT break up treated pasture and plant to sensitive broadleaf crops for at least 3 years after application of *Restore*.

Aerial Application: May be applied by air.

Storage: Store product in original, labelled containers in a secure, dry, cool area. DO NOT freeze.

Buffer Zones:

Handheld equipment is exempt from the buffer zones indicated below when implementing Early Detection and Rapid Response measures on isolated plants or patches. DO NOT apply to water.

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground *	10	10
Fixed wing airplane	80 to 175**	80 to 175**
Helicopter	70 to 150**	70 to 150**

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

** Distance varies depending on spray droplet size. Consult the *Restore/Restore II* labels to determine buffer zone size when applying by air.

† Distance is measured from the downwind edge of the boom to sensitive areas.

DO NOT apply this product directly to any water body or mix or load near water or wells. DO NOT apply when heavy rains are forecast or on moderate to steep slopes toward sensitive areas or to light soils with shallow water table. Contact the provincial environment department for permits to apply near water.

Tank Cleaning:

'Method A' found in the general tank cleaning section on page 15 to 16.

Hazard Rating:

Restore:



Warning – Poison

Restore II:



Danger – Eye and Skin Irritant

For an explanation of the symbols used here see page 11.

Retain SG

This product is a prepackaged tank mix of the equivalent of *thifensulfuron/tribenuron* (page 272) and *fluroxypyr + 2,4-D* (page 168). Information listed is restricted to Crop, Weeds and Rates. For other detailed information on the component products see the product pages listed above.

Herbicide Group

2 - thifensulfuron & tribenuron

4 - fluroxypyr & 2,4-D

(Refer to page 38)

Company:

Loveland Products (formerly Viterro)

Formulation

The *Retain SG* package has 3 components:

Retain A (PCP#30129): 33.35% thifensulfuron + 16.65% tribenuron formulated as a water soluble granule.

Container size - 486 g

Retain B (PCP#29557): 180 g/L fluroxypyr formulated as an emulsifiable concentrate.

Container size - 4.8 L

Loveland Products (formerly Viterro) 2,4-D ester 700 (PCP#29006): 660 g/L 2,4-D ester formulated as an emulsifiable concentrate.

Container size - 6.8 L

Crops and Staging:

Spring wheat (including durum), barley - 4 leaf to flag leaf stage.

Weeds and Staging:

Apply from the seedling to 4 leaf or whorl stage of the following weeds:

Weeds controlled by *thifensulfuron/tribenuron* plus cleavers*

* Not Group 2 resistant biotypes

Rates

Retain A: 8 g per acre.

Retain B: 0.12 L per acre

Loveland Products (formerly Viterro) 2,4-D: 0.2 L per acre

One case per 40 acres

Add *Agral 90*, *Agsurf II*, or *Citovett Plus* at 0.2 L per 100 L of spray solution. *Retain SG* may degrade if left in the sprayer for an extended period. Apply within 24 hours of mixing.

Tank Mixes:

Loveland Products (formerly Viterro) supports the following mixes that are not on the *Retain SG* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Axial*, *Everest 2.0*, *Traxos*, *Simplicity*, *Varro*, *WildCat*, *Puma Advance*.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 14.

See component products for more information on restrictions application details and handling. Use the most limiting restrictions across all components for the mix.

Reward

Company:

Syngenta Canada, distributed by True North Specialty Products (PCP#26271)

Formulation:

240 g/L diquat formulated as a solution.

Container sizes: 4 x 3.78L.

Use:

For use in farm dugouts and other clear, slow moving water bodies to control water weeds, such as:

Canada water weed	Pond weeds
Coontail	Water milfoil
Duckweed	

Offers temporary control of certain species of algae. High levels of suspended organic matter or clay particles in water will reduce control.

NOTE: A permit must be obtained from Saskatchewan Environment or Manitoba Conservation for application of pesticides directly to or within a set distance of water bodies that are not wholly contained within a private parcel of land.

Timing:

Mid-May through late June when water weeds or algae are actively growing. Apply before weeds have developed a heavy mat of growth for effective control.

Rates:

Dugouts less than 5 feet (1.5 m) deep: Apply *Reward* at 7.4 L per acre.

At this rate, 2.2 L of *Reward* will treat a dugout that is 160 feet by 80 feet (49 m x 24.4 m).

Dugouts more than 5 feet (1.5 m) deep: Apply *Reward* at 10.1 to 11.8 L per acre.

At these rates, a dugout that is 160 feet by 80 feet (49 m x 24.4 m) will require 3.0 to 3.5 L of *Reward*.

Herbicide Group

22 - diquat

(Refer to page 38)

Milfoil can be controlled in early stages by 3.7 L per acre in early stages of growth.

Application:

Dilute 1 part *Reward* with 4 parts clean water.

Spray over the water surface, inject below the water surface or pour directly onto the water surface from a moving boat or for small water bodies, apply from the banks. See label for detailed instructions. Note: *Reward* is bound rapidly to soil, so material must enter the water directly to be effective.

How it Works:

Refer to Table 2 on page 40.

Restrictions:

Grazing: DO NOT use water for animal consumption for 24 hours after application.

Irrigation: DO NOT use water for irrigation for 5 days after application.

Domestic Use: DO NOT use water for human consumption for 5 days after application. DO NOT swim in water for 24 hours after treatment.

Storage: DO NOT freeze.

Environment: If weed growth is dense, protect fish by not treating more than one-fourth of dugout at a time.

Equipment Clean Out:

Refer to page 'Method C' in the general section on tank cleaning on 15.

Hazard Rating:



Warning - Poison



Caution - Skin Irritant

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Simazine

Herbicide Group
5 - simazine
 (Refer to page 38)

Company:

Syngenta Canada distributed by True North Specialty Products (*Princep Nine-T* - PCP#16370)
 United Agri Products (*Simazine 480* - PCP#23181)

Formulations:

Princep Nine-T: 90% simazine formulated as a water dispersible granular. Container sizes: 5 kg.

Simazine 480: 480 g/L simazine formulated as a solution. Container sizes: 2 x 10L.

Crops and Staging:

Established alfalfa or bird's-foot trefoil (*Princep Nine-T* only):

DO NOT use in year of seeding. Apply after final cut in fall until freeze-up. DO NOT apply to the same field more than three consecutive years. Residues may build up with yearly applications.

Corn (Field and Sweet): Apply one week prior to seeding and incorporate to a depth of 1 inch (2.5 cm), OR apply no later than 4 days after seeding corn. Rainfall is required to activate herbicide.

Established shelterbelts (elm (American, Siberian), caragana, green ash, Manitoba (boxelder) maple): Apply in fall or early spring before weeds begin growth. Injury may occur to shelter belts growing under saline conditions.

DO NOT apply to frozen ground

Weeds and Staging:

Simazine is applied prior to the emergence of the weeds and kills them when they are exposed to the treated layer of soil.

Barnyard grass
 Lamb's-quarters
 Perennial species starting from seed
 Purslane
 Ragweed

Smartweed (including lady's-thumb)
 Volunteer clovers
 Wild buckwheat
 Wild oats
 Yellow foxtail

Rates:

Forage crops:

Princep Nine-T: 0.45 kg per acre.

Corn*:

Princep Nine-T: 0.61 to 0.81 kg per acre.

Simazine 480: 1.4 to 3.4 L per acre.

Shelterbelts:

Princep Nine-T: 1.8 kg per acre.

Simazine 480: 3.8 to 5.7 L per acre.

* Rate of application to corn is dependent on soil texture. Refer to specific labels for correct application rates on corn.

Application Information:

Water Volume: Minimum 121 L per acre. In shelterbelts, use a minimum of 202 L per acre.

Pressure and Nozzles: For conventional flat fan nozzles use a maximum pressure of 30 to 45 psi (200 to 300 kPa). Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

Use 50 mesh or coarser nozzle screens and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

When applying to forage stands, dry soil conditions at the time of weed emergence may result in reduced weed control.

Tank Mixes:

None registered.

Note: The above mixes are those listed on the simazine labels only. For additional mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Moderate rainfall after application enhances activity.

Re-Entry: DO NOT re-enter treated areas for 12 hours.

Grazing: In forage stands, allow 30 days between application and grazing, 60 days between application and cutting for feed. DO NOT graze or cut corn for feed prior to ear emergence.

Re-cropping: *Simazine* is persistent and residues may persist for several years depending on soil pH, available soil moisture, number of yearly applications, and the sensitivity of the following crop. *Simazine* will break down in soil more slowly under conditions of high pH and/or low rainfall. Corn will tolerate soil residues of *simazine* and may be planted the year of application. White beans, onions, peas may be injured 12 month after application.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze *Simazine* 480. *Princep Nine-T* may be frozen. Store in a cool, dry place.

Buffer Zones:

Crops	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Alfalfa	1	1	4
Bird's-foot trefoil, sweet corn	1	1	5
Field corn	1	1	10
Shelter-belts	2	1	20

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Caution – Poison. (*Simazine* 480)

For an explanation of the symbols used here see page 11.

Simplicity

Herbicide Group

2 - pyroxsulam

(Refer to page 38)

Company:

Dow AgroSciences (PCP#28887)

Formulation:

30 g/L pyroxsulam formulated as an oil dispersion.
Container size - 2 x 8 L jug.

Crops and Staging:

Wheat (Spring and durum): 3 leaf stage until prior to the emergence of the flag leaf (up to 6 leaf plus 2 tillers).

Winter wheat: 1 to 3 leaf stage in fall or 2 to 7 leaf plus 4 tillers in spring.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds, Rates and Staging:

Wild oats (less than 75 plants per sqm):

0.15 L per acre (one case treats 106 acres)

All weeds listed below: 0.20 L per acre (one case treats 80 acres).

Shake *Simplicity* jug well before adding to spray tank.

Grasses:

WEED	STAGE
Wild oat	up to the 4 leaf, 2 tillers
Barnyard grass, Yellow foxtail Green foxtail*	1 to 5 leaf
Japanese brome	1 to 6 leaf
Downy brome†	

Broadleaves:

Canada thistle* (up to 30 cm, before budding)	Hemp-nettle (1-8 leaf)
Cleavers (up to 6 whorl)	Redroot pigweed (1-8 leaf)
Cow cockle (up to 8 leaf)	Round-leaved mallow (up to 6 leaf or 10 cm)
Common chickweed (up to 10 cm)	Russian thistle* (up to 10 cm)
Corn spurry (up to 2 whorl or 10 cm tall)	Shepherd's-purse (up to 30 cm)
Dandelion* (spring rosettes <20 cm diameter)*	Smartweed (1-5 leaf)
Flixweed (up to 10 cm)	Stinkweed (up to 30 cm)
	Volunteer canola (1-6 leaf)**
	Wild buckwheat (1-4 leaf) *

* Suppression only.

** Not Clearfield varieties

† Control with fall application in winter wheat ; suppression only in spring applications on both winter and spring wheat.

When applied alone, *Simplicity* must be applied using a non-ionic surfactants (*Agral 90*, *AgSurf II* or *Surf 92*) at 0.25 L per 100 L.

Winter wheat (*Simplicity* alone) use one of the following adjuvants:

Assist at 0.8 L per 100 L of spray

Merge at 0.5 L per 100 L of spray (spring application only)

* Adjuvants purchased separately.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre.

Aerial: 12 L per acre

Nozzles and Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets. See the label for detailed instructions on aerial application.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury may result and/or weed control may be reduced.

Tank Mixes:

Herbicides:

The addition of an adjuvant is not required in tank mixes unless the adjuvant is required by the tank mix partner.

TANK-MIX PARTNER	PRODUCT RATES
<i>Buctril M</i>	0.4 L per acre
<i>Curtail M</i>	0.6 L per acre
<i>Frontline 2,4-D</i>	60 acres per case
<i>Frontline XL</i>	0.5 L per acre
<i>MCPA ester (500 formulation)</i>	0.28 to 0.45 L per acre
<i>Prestige XC</i>	27 acres per case
<i>Spectrum</i>	20 acre per case
<i>Thumper</i>	0.4 L per acre

Fungicides:

*Tilt** (label rates)

*Stratego** (label rates)

MCPA + *Tilt**

MCPA + *Stratego**

Fertilizers: None registered

* High rate of *Simplicity* only.

Note: The above mixes are those listed on the *Simplicity* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Dow AgroSciences also supports the following mixes that are not on the *Simplicity* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: 2,4-D ester, *Attain XC* (low use rate), *Attain XC* + either *Tilt* or *Stratego*, *Barricade*, *Bromoxynil*, *OcTtain*, *Retain*, *Stellar*, *Thifensulfuron/tribenuron* (incl. +*Perimeter MEGA*).

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Preharvest: Leave 60 days between treatment and harvest.

Grazing: Must NOT be grazed or fed to livestock for 7 days after treating crop.

Recropping: Barley, condiment and oilseed quality brown mustard (*B. juncea* types), canola, chickpea, dry bean, flax, lentil, oat, field pea, potato, spring wheat, soybean, sunflower and yellow mustard may be seeded 11 months following treatment.

Aerial Application: May be applied by air.

Storage: Avoid freezing, store above -9°C. Allow product to warm above 7°C before using and thoroughly mix the product prior to use.

Buffer Zones:

Application method	Buffer Zones (metres†)		
	Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground*	1	1	2
Helicopter	1	1	60
Fixed wing aircraft	1	1	70

See page 29 for an explanation of the different habitats.

* These distances can be reduced by 30% using cones on individual nozzles and by 70% using a full shield (shroud, curtain) that extends to the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Equipment used to apply *Simplicity* should not be used to apply other pesticides to sensitive crops without thorough cleaning. To avoid subsequent injury to crops other than cereals, all spraying equipment must be thoroughly cleaned both inside and out, as follows:

1. Immediately after spraying drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Add *All Clear* tank cleaner at 0.5 L per 100 L of water or *Clean-Out* tank cleaner at 0.25 L per 100 liters of water while filling the tank ½ full with clean water. Agitate for at least 15 minutes ensuring the cleaning solution comes in contact with interior surfaces. Flush the boom and hoses with the cleaning solution and be sure to remove caps at the end of booms to allow cleaning solution to reach all areas of the boom. Leave the spray solution in the sprayer for an extended period if possible (eg. overnight). Thoroughly drain the sprayer.
4. Remove nozzles and screens and clean separately with *All Clear* cleaning solution (50 mL in 10 L water).
5. Rinse the tank with clean water and flush through the booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.

Refer to page 15 to 16 for additional information on tank cleaning.

Hazard Rating:



Warning – Poison



Warning – Contains the allergen soy



Caution – Eye and Skin Irritant, Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Solo

Herbicide Group
2 - imazamox
 (Refer to page 38)

Company:

BASF Canada

All crops listed under 'Crops And Staging:' below
 (PCP#25496)

CLEARFIELD canola only (PCP#28741)

Formulation:

70% imazamox as a water dispersible granule.

Container size - 4 x 117 g water soluble bags.

Crops and Staging:

CLEARFIELD sunflower: 2 to 8 leaf stage.

CLEARFIELD canola: 2 to 6 leaf stage.

CLEARFIELD lentil: 2 to 6 leaf stage.

CLEARFIELD oilseed mustard (*Brassica juncea*): 2 to 6 leaf stage

Dry Beans: Up to the second trifoliate leaf stage. Must be mixed with *Basagran Forté* - See tank mix section.

Temporary crop yellowing may be observed shortly after application in CLEARFIELD canola.

Weeds and Staging:

Grasses - 1 to 4 main stem leaves, early until tillering.

Barnyard grass	Volunteer oat
Green foxtail	Volunteer wheat (not CLEARFIELD varieties)
Japanese brome*	Wild oat
Persian darnel	Yellow foxtail
Volunteer barley	
Volunteer canaryseed	

Broadleaf Weeds - cotyledon to 4 leaf stage.

Cleavers*	Shepherd's-purse
Cow cockle	Stinkweed
Green smartweed	Volunteer canola (not CLEARFIELD varieties)
Lamb's-quarters	Wild buckwheat*
Redroot pigweed	Wild mustard

* Suppression only.

Rates:

Solo: 11.7 g per acre (40 acres per case)

Merge adjuvant (sold separately): Must be used with *Solo* at a rate of 0.5 L of *Merge* per 100 L of total mixed spray solution.

DO NOT apply *Solo* more than once or follow *Solo* with any products containing imazamox in the same year.

Refer to the product label for complete mixing instructions for this product and its mixes.

A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 40 L per acre.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage and a minimum of fine droplets that are prone to drift. Use 50 mesh or coarser filter screens.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT spray if temperatures of +5°C are forecast within 3 days of application. Treat crops during warm weather when weeds are actively growing and soil moisture is adequate for rapid growth. Under cool or dry conditions, control of some weeds may be severely reduced.

Tank Mixes:

Dry Beans:

Solo must be mixed with *Basagran Forté* (0.5 L/acre) plus UAN (liquid 28-0-0) at 0.81 L/acre. The addition of 28% UAN is required for proper activity.

Restrictions:

Rainfall: DO NOT spray if there is a forecast of rain during or soon after application as it may reduce control.

Re-Entry: DO NOT enter treated fields for 12 hours.

Grazing: DO NOT graze treated canola or lentil or cut for feed within 20 days of application.

DO NOT graze treated sunflower or cut for straw.

Preharvest Interval: DO NOT apply to canola or lentil within 60 days of harvest.

DO NOT apply to sunflower within 70 days of harvest.

Re-cropping: Winter wheat may be seeded 3 months after application. Barley, canaryseed, canola, chickpea, field corn, field pea, flax, lentil, oat, sunflower, and spring wheat (including durum) may be seeded the first spring after application and tame mustard (condiment types only) the second season after application. The company recommends that a field bioassay (a test strip grown to maturity) be conducted the year before growing any crops other than those listed above.

Contact manufacturer for additional information on recroping intervals (1-877-371-2273).

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Store in a cool, dry place above 5° C.

Buffer Zones: Avoid spraying in situations where drift may occur. Leave at least 11 m between the outside edge of the sprayed area and sensitive non-target areas such as shelterbelts, hedgerows, wetlands, woodlots, vegetated ditch banks, ponds, streams, and sloughs.

Tank Cleaning:

Refer to 'Method C' in the general tank cleaning section on page 15.

Hazard Rating:



Warning - Eye and Skin Irritant.

May cause eye damage.

For an explanation of the symbols used here see page 11.

Spectrum

Company:

Dow AgroSciences

Formulation:

Each case of *Spectrum* contains 2 components:

Spectrum A (PCP# 27031): 50 g/L florasulam formulated as a suspension concentrate. Container size - 0.8 L.

Spectrum B (PCP# 27032): 50 g/L clopyralid and 280 g/L of MCPA ester formulated as an emulsifiable concentrate. Container size - 12 L.

Crops and Staging:

Spring wheat (including durum), barley and oats in the 2 to 6 leaf stage.

Forage Grasses* (seedling and established) grown for seed production: No staging indicated for forage grasses.

Bromegrass (meadow, smooth, hybrid)	Perennial Ryegrass
Fescue (chewings, creeping red, hard, tall)	Timothy
	Wheatgrass (crested, intermediate)

* NOTE: Since these uses are registered under the User Requested Minor Use Label Expansion (URMULE) program, the manufacturer assumes no responsibility for herbicide performance. Users of this product on forage grasses do so at their own risk.

Herbicide Group

2 - florasulam

4 - clopyralid & MCPA

(Refer to page 38)

Weeds and Staging:

Broadleaf weeds controlled at the 2 to 4 leaf stage:

Canada thistle	Smartweed
Chickweed (common)	Sow-thistle (annual)
Cleavers	Sow-thistle (perennial) [†]
Dandelion**	Stinkweed
Flixweed*	Stork's-bill
Hemp-nettle	Volunteer canola (all varieties)
Lamb's-quarters	Wild mustard
Redroot pigweed	Wild buckwheat
Shepherd's-purse	

Broadleaf weeds suppressed:

Dandelion***

* Spring seedlings only.

** Seedlings and overwintered rosettes < 15 cm.

*** Overwintered rosettes > 15 cm; mature plants.

[†] Top growth control only.

Rates:

Spectrum A: 40 mL per acre

Spectrum B: 600 mL per acre

One case treats 20 acres.

Note: Maximum one application of this product or other products containing florasulam in back to back years.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 14.

Application Information:

Water Volume: 40 L per acre.

Nozzles & Pressure: Maximum 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity of *Spectrum*. Weeds hardened off by cold weather or drought stress may not be adequately controlled or suppressed and regrowth may occur. Under conditions of low crop and high weed density, control may be reduced. Extreme growing conditions such as drought or near freezing temperature prior to, at, or following time of application, may increase the risk of crop injury at all stages of growth.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:
Assert (0.65 L/acre) plus *Acidulate*

In spring wheat (NOT durum) and barley:
Axial 100EC (243 mL/acre) plus *Adigor* adjuvant

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Spectrum* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Dow AgroSciences also supports the following mixes that are not on the *Spectrum* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Simplicity*, *Avenge*.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT re-enter treated fields for 12 hours.

Grazing: DO NOT graze treated crop or cut for feed within 7 days of application. Withdraw meat animals from treated fields at least 3 days before slaughter.

Preharvest: DO NOT apply within 60 days of harvest.

Re-cropping: Barley, canola, field pea*, oat and wheat may be grown the year following an application or the field can be summerfallowed.

* DO NOT seed to field pea for at least 10 months following treatment. Very dry soil conditions following application can result in a risk of injury to field pea grown in rotation. If severe drought conditions are experienced during the months of June to August inclusive in the year of application delay seeding field pea an additional 12 months (22 months following application). Contact your local Dow AgroSciences representative or retailer for more information before seeding field pea following drought conditions in the previous year.

Aerial Application: DO NOT apply by air.

Storage: Store in dry, heated (greater than 5°C) area. *Spectrum A* will freeze at -10°C. If frozen, bring to room temperature and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	5	5	30

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Handheld or backpack applications do not require a buffer.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16. Check the cleanout requirements of pesticides mixed with this product. Additional cleanout measures may need to be integrated into those provided here.

Hazard Rating:



Caution – Poison



May cause eye irritation

For an explanation of the symbols used here see page 11.

Stellar

Herbicide Group
2 - florasulam
4 - fluroxypyr, MCPA
(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

Stellar A (PCP# 29286): 2.5 g/L florasulam and 100 g/L fluroxypyr formulated as a suspension concentrate.
 Container size - 2 x 8 L jug.

Stellar B (PCP# 29165): 600 g/L of MCPA ester formulated as an emulsifiable concentrate.
 Container size - 1 x 9.33 L jug.

Crops and Staging:

Barley, oat and spring wheat (including durum) - 2 to 6 leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Apply when weeds are at the 2 to 4 leaf stage.

Burdock	Russian pigweed
Chickweed (Common)	Shepherd's-purse
Cleavers	Smartweed
Cocklebur	Stinkweed
Flixweed	Stork's-bill*
Hemp-nettle	Sunflower (annual)
Kochia	Vetch
Lamb's-quarters	Volunteer canola
Plantain	Volunteer flax
Prickly lettuce	Wild buckwheat
Ragweed	Wild mustard
Redroot pigweed	Wild radish

* Suppression only

Rates:

Stellar A: 0.4 L per acre

Stellar B: 0.24 L per acre
 (One case treats 40 acres)

Application Information:

Water Volume: Minimum 40 L per acre.

Nozzles & Pressure: For conventional flat fan nozzles use a pressure of 30 to 40 PSI (200 to 275 kPa). Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets. Low drift nozzles may require higher pressures for proper performance.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops or weeds that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury or reduced weed control may result.

Tank Mixes:

Herbicides:

Barley, spring wheat, and durum:

Assert (0.65 L/acre)

Spring wheat (not durum) and barley only:

Axial 100EC (243 mL/acre) plus *Adigor* adjuvant

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Stellar* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Dow AgroSciences also supports the following mixes that are not on the *Stellar* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Simplicity.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Preharvest: Leave 60 days between treatment and harvest.

Grazing: DO NOT graze or harvest for livestock feed within 7 days of treating the crop.

Recropping: Wheat, barley, oat, canola, and pea may be grown the season following application or the field may be fallowed. There are no recropping restrictions the second year after application.

Aerial Application: DO NOT apply by air.

Storage: May be frozen. If frozen, bring to room temperature and agitate before use. This product is combustible. DO NOT store near heat or open flame.

Buffer Zones: Leave 30 metres between the downwind edge of the boom and sensitive terrestrial habitats such as forested areas shelterbelts, woodlots, hedgerows, and shrub lands and 15 metres to sensitive freshwater habitats such as lakes, rivers, sloughs ponds, prairie potholes, creeks marshes streams reservoirs and wetlands.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16. If mixed with another pesticide additional clean-out measures may be necessary.

Hazard Rating:



Warning – Poison



Warning – Eye and Skin Irritant.



Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Tandem

Herbicide Group

2 - pyrosulam

4 - fluroxypyr

(Refer to page 38)

Company:

Dow AgroSciences

Formulation:

The *Tandem* package has 2 components:

Tandem A (PCP# 29985): 30 g/L pyrosulam formulated as an oil dispersion.

Container size - 8 L jug.

Tandem B (PCP# 29965): 333 g ae/L fluroxypyr formulated as an emulsifiable concentrate.

Container size - 4.84 L

Crops and Staging:

Spring wheat (including durum): 3 leaf stage until the first node can be felt in the stem (up to 6 leaf plus 2 tillers).

When tank-mixing always check the tank mix partner recommendations for additional staging restrictions.

Winter wheat: Apply in the spring from the 3 tiller stage to just before the flag leaf stage.

Weeds, Rates and Staging:

Tandem A at 0.15 L per acre plus *Tandem B* at 85 mL per acre (53 acres per case):

Cleavers (up to 6 whorls)

Wild oats (less than 75 plants per sqm)

Tandem A at 0.20 L per acre plus *Tandem B* at 127 mL per acre (40 acres per case):

Grasses:

WEED	STAGE
Wild oat	up to the 4 leaf, 2 tillers
Barnyard grass, Yellow foxtail Green foxtail*	1 to 5 leaf
Japanese brome	1 to 6 leaf

Broadleaves:

WEED	MAXIMUM APPLICATION STAGE
Wild buckwheat	4 leaves
Smartweed	5 leaves
Round-leaved mallow, Volunteer canola**	6 leaves or whorls
Cleavers, Cow cockle, Hemp-nettle†, Kochia†, Redroot pigweed, Stork's-bill*	8 leaves or whorls
Common chickweed, Flixweed	10 cm
Volunteer flax	12 cm
Dandelion*	spring rosettes <20 cm diameter

* Suppression only.

** Not Clearfield varieties

† NOTE Group 2 resistant biotypes only controlled at smaller stage. See label for details.

Application Information:**Water Volume:**

Ground: 20 to 40 L per acre.

Aerial: 20 L per acre.

Nozzles and Pressure: Use 29 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE coarse droplets. Low drift nozzles may require higher pressures for proper performance. See the label for detailed instructions on aerial application.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost before application (3 days) or shortly after (3 days) may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions, (e.g. drought, heat or cold stress, or if weeds have initiated flowering), or if heavy infestations exist.

Tank Mixes:**Herbicides:**

In spring wheat (including durum):

2,4-D Ester 700 (0.24 to 0.32 L/acre)

Curtail M (0.61 to 0.81 L/acre)

MCPA Ester (0.28 to 0.37 L/acre)

Fungicides:

Tilt, Stratego

Note: The above mixes are those listed on the Tandem label only. For other possible registered mixes see the blue fold out chart inside back cover.

Adding ingredients in the correct order is critical for optimum performance.

Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 4 hours may reduce control.

Re-entry: DO NOT re-enter treated fields for 12 hours.

Preharvest Interval: Leave 60 days from treatment to harvest.

Grazing: Must NOT be grazed or fed to livestock for 7 days after treating crop.

Recropping: Barley, canola, flax, lentil, mustard, oat, field pea, spring wheat may be seeded 11 months following treatment or fields may be fallowed.

Aerial Application: May be applied by air.

Storage: Avoid freezing, store above -9°C. Allow product to warm above 7°C before using and thoroughly mix the product prior to use.

Buffer Zones:

Application method	Buffer Zones (metres†)		
	Required for the Protection of:		Terrestrial habitat
	Aquatic Habitats of Depths Less than 1 m	Greater than 1 m	
Ground *	1	1	2
Helicopter	1	1	55
Fixed wing aircraft	1	1	65

See page 29 for an explanation of the different habitats.

* These distances can be reduced by 30% using cones on individual nozzles and by 70% using a full shield (shroud, curtain) that extends to the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Equipment used to apply Tandem should not be used to apply other pesticides to sensitive crops without thorough cleaning. To avoid subsequent injury to crops other

than cereals, all spraying equipment must be thoroughly cleaned both inside and out, as follows:

1. Immediately after spraying drain the sprayer tank. Any contamination on the outside of the spraying equipment should be removed by washing with clean water.
2. Rinse inside of tank with clean water and flush through booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.
3. Add *All Clear* tank cleaner at 0.5 L per 100 L of water or *Clean-Out* tank cleaner at 0.25 L per 100 liters of water while filling the tank ½ full with clean water. Agitate for 15 minutes ensuring the cleaning solution comes in contact with interior surfaces. Flush the boom and hoses with the cleaning solution and thoroughly drain the sprayer.
4. Remove nozzles and screens and clean separately with *All Clear* cleaning solution (50 mL in 10 L water).
5. Rinse the tank with clean water and flush through the

booms and hoses using at least one tenth of the spray tank volume. Drain tank completely.

Refer to page 15 to 16 for additional information on tank cleaning.

Hazard Rating:



Danger – Poison.



Warning – Contains the allergens milk, soy and sulphites.



Caution – Eye and Skin Irritant, Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Tensile

This product is a prepackaged tank mix of *Solo* (page 265) and *Lontrel Dry* (*Lontrel Dry* is only available with *Absolute* and *Tensile*) (page 216). Information listed is restricted to Crop, Weeds and Rates. For other detailed information on the effect of growing conditions, and restrictions for the component products see the product pages listed above.

Herbicide Group

2 - imazamox

4 - clopyralid

(Refer to page 38)

Company:

BASF Canada

Formulation:

Tensile package contains 2 components:

Solo (PCP# 28741): 70% imazamox formulated as a dispersible granule.

Lontrel Dry (PCP# 27306): 75% clopyralid formulated as a dispersible granule.

Container size -

Solo: 4 x 117 g water soluble bags

Lontrel Dry: 2 x 810 g.

Crops and Staging:

CLEARFIELD canola varieties: 2 to 6 leaf stage.

Apply only to CLEARFIELD canola varieties; application to any other variety of canola or any other crop will result in crop death.

Weeds and Staging:

Weeds controlled by *Solo* plus.

Canada thistle (rosette to pre-bud stage)*

Sow-thistle, annual

Sow-thistle, perennial*

Wild buckwheat

* Top growth control for 6 to 8 weeks

Rates:

Solo: 11.7 g per acre.

Lontrel Dry: 40 g per acre.

(One case treats 40 acres)

Merge: 0.5 L per 100 L of spray solution (sold separately). At a spray volume of 40 L per acre one 8.1 L jug of *Merge* will treat 40 acres. *Tensile* MUST be applied with *Merge* adjuvant.

DO NOT apply *Tensile* more than once or follow *Tensile* with any other products containing clopyralid or imazamox in the same year.

Refer to the product label for complete mixing instructions for this product and its mixes. A general guide to mixing can be found on page 13.

Restrictions and Application Information:

See component products for additional information including restrictions. Use the most limiting restrictions across all components for the mix. Particular attention should be paid to the recropping restrictions for both *Solo* and *Lontrel*.

Thifensulfuron/tribenuron

Herbicide Group
2 - thifensulfuron &
tribenuron
(Refer to page 38)

Company:

E. I. duPont Canada (*Refine SG*)
 Arysta LifeSciences (*Deploy WDG*)
 Cheminova Canada (*Nimble*)
 Farmers of North America (*MPower R*)

Formulation:

Refine SG (PCP# 28285): 33.35% thifensulfuron methyl plus 16.65% tribenuron methyl formulated as a water soluble granule.
 Container size - 486 g bottle.

Deploy WDG (PCP# 30846); *MPower R* (PCP# 30945); *Nimble* (PCP# 29467) = 75% WDG formulations: 50% thifensulfuron methyl plus 25% tribenuron methyl formulated as a water dispersible granule.
 Container size - 320 g container.

Crops and Staging:

Apply from 2 leaf to the flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions

Cereals:

Barley	Wheat (including durum
Oat	spring and winter),

Seedling or established forage grasses for forage or seed production:

Bromegrass (meadow, smooth)	Wheatgrass (crested, intermediate, northern,
Fescue (creeping red, tall)	pubescent, slender, stream-
Kentucky bluegrass**	bank, tall, western)
Orchardgrass	

* NOTE: Since the use of this product on forage grasses is registered under the User Requested Minor Use registration system, the manufacturer assumes no responsibility for herbicide performance. Users of this product on forage grass do so at their own risk.

** Established stands only.

Weeds and Staging:

Unless otherwise noted below, apply to young and actively growing weeds that are less than 4 inches (10 cm) in height or width.

Weeds Controlled:

Annual smartweed (green, lady's-thumb)	Redroot pigweed
Ball mustard	Russian thistle
Chickweed (1 to 6 leaf)	Shepherd's-purse
Common groundsel	Stinkweed
Corn spurry	Tartary buckwheat
Cow cockle	Volunteer canola
Flixweed	(CLEARFIELD varieties controlled with 2,4-D or MCPA mixes only)
Hemp-nettle	Volunteer sunflower
Lamb's-quarters	Wild buckwheat*
Narrow-leaved hawk's-beard	Wild mustard

* *Refine SG*: up to 5 leaf stage; 75% WDG formulations: up to 3 leaf stage only.

Weeds Suppressed:

Canada thistle, sow-thistle (less than 6 inches (15 cm) tall or across and prior to budding)
 Cleavers (1 to 3 whorls)
 Round-leaved mallow (2 to 6 leaf)
 Scentless chamomile
 Stork's-bill (2 to 6 leaves)
 Toadflax (less than 6 inches or 15 cm tall)

Rate:

Refine SG: 12 g per acre (one 486 g container treats 40 acres).

75% WDG formulations: 8 g per acre (one 320 g container treats 40 acres).

Maximum of one application of *Thifensulfuron/tribenuron* or other products with the same ingredients per year.

Add *Agral 90*, *Agsurf*, or *Citowett Plus* surfactants at 0.2 L per 100 L of spray solution.

75% WDG formulations may also use *Liberate* and *Nufarm Enhance* surfactants at 0.2 L per 100 L of spray solution.

Thifensulfuron/tribenuron may degrade if left in the sprayer for an extended period. *Apply within 24 hours of mixing*. Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 14.

Application Information:**Water Volume:**

Ground: Minimum 22 L per acre.

Aerial (*Refine SG only*): Minimum 10 L to maximum 20 L per acre.

Nozzles and Pressure: Use 30 to 40 psi (210 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE medium* droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to wheat, barley or oats that are stressed by severe weather conditions (frost, drought or water saturated soil) as crop injury may result.

Under certain conditions (heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) lightening in crop colour and reduction in crop height may occur.

Tank Mixes:

Herbicides:

Tank Mix Partner	CROPS				
	Spring wheat	Winter wheat	Durum	Barley	Oats
2,4-D amine or ester (0.34 to 0.45 L/acre) [†]	✓	✓	✓	✓	
Assert (0.53 to 0.65 L/acre)	✓		✓	✓	
Assert (0.53 to 0.65 L/acre) + MCPA ester (0.28 to 0.45 L/acre)*	✓		✓	✓	
Axial 100EC (243 mL/acre) plus Adigor adjuvant	✓			✓	
Axial 100EC (243 mL/acre) + MCPA Ester (0.23 [◇] or 0.28 L/acre) plus Adigor adjuvant	✓			✓	
Banvel II (44.5 mL/acre to 58.7 mL/acre ^{**}) [†]	✓		✓ [△]	✓	
Curtail M (0.61 L/acre) [†]	✓			✓	
Flurox 2,4 (one case treats 80 acres) [†]	✓ ^{△△}		✓ ^{△△}	✓ ^{△△}	
Clodinafop 240EC (95 to 115 mL/acre) plus Score adjuvant	✓		✓		
Clodinafop 240EC (95 mL/acre) + Banvel II (44.5 mL/acre to 58.7 mL/acre ^{**}) plus Score adjuvant	✓		✓ [△]		
Clodinafop 240EC (95 mL/acre) + MCPA ester (0.23 [◇] or 0.34 to 0.45 L/acre)* plus Score adjuvant	✓		✓		
Lontrel 360 (85 mL/acre) [†]	✓			✓	
Lontrel 360 (85 mL/acre) + 2,4-D ester* or MCPA ester* (0.34 L/acre) [†]	✓			✓	
MCPA amine or ester (0.23 [◇] or 0.28 to 0.45 L/acre)*	✓	✓	✓	✓	✓
Fenoxaprop 120EC (0.16 to 0.31 L/acre)	✓		✓	✓	
Fenoxaprop 120EC (0.16 to 0.31 L/acre) + MCPA ester (0.23 [◇] or 0.34 L/acre)*	✓		✓	✓	
Simplicity (0.15 to 0.20 L/acre) [†]	✓ [△]		✓ [△]		

[†] Marked tank mixes require the addition of a non-ionic surfactant. Unmarked mixes do not require additional adjuvant beyond what is provided for by the tank mix partner.

[△] Refine SG only.

^{△△} Deploy, MPower R and Nimble only.

* 500 g ai/L formulation.

** High rate of Banvel II with Refine SG only.

[◇] Tank mix with 0.23 L/acre to control Clearfield canola at the 2 to 4 leaf stage.

Check the above tank mix partner(s) respective labels for additional staging and varietal restrictions.

Fertilizers: None registered.

Note: The above mixes are those listed on the Thifensulfuron/tribenuron labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

E.I. duPont also supports the following mixes that are not on the Refine SG label. Mixes must be applied according to the most restrictive use limitations for either product:

Herbicides: *Avenge, Simplicity*

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rainfall of 1 inch (25 mm) or more beginning within 1 hour of application of *Refine SG* or 4 hours for 75% WDG formulations may reduce control.

Re-Entry: DO NOT re-enter treated fields for 12 hours.

Grazing: Must NOT be grazed or fed to livestock for 7 days after treatment.

Re-cropping: No restrictions the year after treatment. Canola, flax, lentil and alfalfa may be planted 2 months after application.

Aerial Application: *Refine SG* may be applied by air. DO NOT apply 75% WDG formulations by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	0	15
Fixed wing airplane	1	0	125
Helicopter	1	0	100

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

75% WDG formulations (Ground equipment only):

Leave a 15 m buffer zone between last spray swath and sensitive upland or aquatic habitats such as shelterbelts, wetlands, sloughs, and woodlots.

Tank Cleaning:

Thifensulfuron/tribenuron can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray *Thifensulfuron/tribenuron* should be drained and flushed out immediately after use.

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16. If mixing with another pesticide with different cleaning measures, those measures should be integrated into 'Method A' (e.g. addition of detergent).

Hazard Rating:

75% WDG formulations:



Warning – Eye and Skin Irritant

Refine SG:



Warning – Contains the allergen milk.

For an explanation of the symbols used here see page 11.

Thifensulfuron/tribenuron + MCPA ester

These products are prepackaged tank mix of *Refine SG* (page 272) and *MCPA ester* (page 218). Information listed is restricted to Crop, Weeds and Rates and Tank mixes. For other detailed information on the component products see the product pages listed above.

Herbicide Group
2 - thifensulfuron & tribenuron
4 - MCPA
(Refer to page 38)

Company:

E. I. duPont Canada (*Refine M*)
Loveland Products (formerly Viterra) (*BroadSide*)

Formulation:

Refine SG (PCP# 28285): 33.35% thifensulfuron methyl + 16.65% tribenuron methyl; formulated as a water soluble granule.

Container size - 8 x 121 g water soluble packets.

-plus-

MCPA ester (PCP# 26161): 500g/ L MCPA formulated as an emulsifiable concentrate.

Container size - 2x 9.1 L of *MCPA ester*.

-or-

MCPA ester (PCP# 27803): 600g/ L MCPA formulated as an emulsifiable concentrate.

Container size - 2x 7.6 L of *MCPA ester*.

Crops and Staging:

Barley, wheat (including durum and winter) and oat from full 3 leaf to the flag leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions

Weeds and Staging:

Weeds Controlled or Suppressed by *Refine SG* plus
'Susceptible Weeds' controlled by *MCPA ester*, plus:

Dandelion (rosettes, less than 15 cm in diameter)	Volunteer canola (2 to 4 leaf) (including CLEARFIELD varieties)
---	---

Rate:

Refine SG : 12 g per acre

MCPA Ester: (500 g per L) - 0.23 L per acre, (600 g per L) - 0.19 L per acre

(One case treats 80 acres)

Refer to the product labels for complete mixing instructions. A general guide to mixing can be found on page 13.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Assert (0.54 to 0.67 L/acre)

In spring wheat (NOT durum) and barley:

Axial 100EC (243 mL/acre) plus *Adigor* adjuvant

Lontrel 360 (85 mL/acre)

Check the above tank mix partners respective labels for additional staging and varietal restrictions.

Note: The above mixes are those listed on the *Refine SG* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

See component products for more information on restrictions application details and handling. Use the most limiting restrictions of each component.

Tordon 22K

Herbicide Group
4 - picloram
 (Refer to page 38)

Company:

Dow AgroSciences (PCP#9005)

Formulation:

240 g/L picloram acid present as a potassium salt, formulated as a solution.

Container size - 10 L, 3.6 L

Note: Available only through selected retail outlets.

Crop and Staging:

Apply at any stage of permanent grass pastures, rangeland and non-cropland.

NOTE: It is strongly recommended that this product be applied by a licensed applicator.

Weeds, Rates and Staging:

For the control of biennial and deep-rooted perennial weeds listed below:

Weed	Rate L/Acre	Backpack (mL of Tordon 22K per 100 M ²)*
Scentless chamomile	0.445	11
Knapweed (diffuse, spotted)	0.91	22
Canada thistle, pasture sage, poverty weed, Russian knapweed, perennial sow- thistle	1.8	45
Leafy spurge, field bind- weed, toadflax	3.6†	90†

† NOTE: This rate is only registered for use with hand application equipment (wand or backpack) and only one acre of every two may be treated in this manner at this rate.

* mix with 18 litres of water and the spray solution over 100 square metres.

For best results, applications should be made when perennial weeds have fully developed, green leaves. Application in late summer (or periods of dry weather) when plants are not actively growing may result in unsatisfactory control.

Application Information:

Water Volume: 160 to 325 L per acre without spray running off foliage.

Nozzles and Pressure: Maximum 150 to 350 kPa (20 to 50 psi) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of application equipment and pressure that is designed to deliver an even coverage of coarse droplets that are not prone to drift. Non-target broadleaf plants are very sensitive to Tordon 22K drift.

Avoid conditions that are conducive to drift. (See page 12 for drift control suggestions)

How it Works:

Tordon 22K interferes with cell division, causing leaf cupping, stem distortion and eventual death. Tordon 22K is absorbed through the leaves and roots.

IMPORTANT: Tordon 22K is a very persistent and water-soluble herbicide. Treated soil should NOT be moved from the treated area. DO NOT apply to soils that are permeable, have sinkholes, or lie over limestone bedrock. DO NOT apply to soils whose surfaces are composed of fractured rock or unconsolidated gravel. Application to these sites may allow the movement of herbicide to underlying water sources or aquifers. If shallow aquifers are present, DO NOT apply Tordon 22K. This product is moderately toxic to fish. DO NOT apply to any water bodies or in areas where the runoff from treated areas will reach fish-bearing waters.

Tordon 22K must not be applied on range and pasture acres that are irrigated. DO NOT compost or mulch clippings from grass treated with Tordon 22K.

Effects of Growing Conditions:

Avoid application when pasture and target weeds are under stress from drought, flooding, extreme heat or cold, as injury to grass or unacceptable control may result. Avoid spraying if temperatures exceed 28°C.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Rain within 6 hours of application may cause poor results. Heavy rainfall may dissolve and carry *Tordon 22K* away from the target area, or it may leach dissolved *Tordon 22K* out of the root zone or to undesirable locations.

Grazing: DO NOT graze lactating dairy animals within 6 weeks after treatment. There are no grazing restrictions for other livestock. DO NOT use manure from animals grazing treated forage to fertilize susceptible plants or crops.

Re-cropping: *Tordon 22K* may persist in the soil for up to 5 years. For this reason *Tordon 22K* may only be applied on permanent grass pastures and rangeland unless applied by an authorized pesticide applicator. Avoid the root zone of desirable trees or shrubs.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze.

Buffer Zones:

Hand-held or backpack sprayer and spot treatment DO NOT require a buffer zone from sensitive habitat, but efforts should be made to minimize exposure to sensitive plants and open water or wetlands.

Application method	Buffer Zones (metres [†]) Required for the Protection of:	
	Aquatic Habitats	Terrestrial habitat
Ground only	1	120

See page 29 for an explanation of the different habitats.

* These distances can be reduced by 30% using cones on individual nozzles and by 70% using a full shield (shroud, curtain) that extends to the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Tordon 22K can cause severe injury to sensitive crops (especially pulses and other broadleaf crops) at very low concentrations. Spray equipment should be flushed out immediately after spraying *Tordon 22K*. Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Caution – Poison.



Danger – Eye Irritant.



May Cause Skin Irritation.

For an explanation of the symbols used here see page 11.

Tralkoxydim

Herbicide Group
1 - tralkoxydim
 (Refer to page 38)

Company:

Dow AgroSciences (*Liquid Achieve* - PCP# 28555; *Turbocharge* adjuvant - PCP# 28554)

MANA Canada (*Bison* - PCP# 29256; *Addit* adjuvant - PCP# 29263)

Loveland Products (formerly Viterro) (*Marengo* - PCP# 29289; *Turbocharge B* adjuvant - PCP# 29288)

Formulation:

400 g/L tralkoxydim formulated as a suspension concentrate. Container sizes -

Marengo: 8 L of tralkoxydim plus 4L *Turbocharge*.

Bison: 8 L of tralkoxydim, 8 L *Addit* adjuvant

Liquid Achieve: 8 L of tralkoxydim (*Turbocharge* sold separately).

Contact manufacturers.

Crops and Staging:

No staging restrictions unless otherwise indicated.

Cereals:

Barley	Triticale
Rye (spring & fall)	Wheat (spring, durum, & winter)

Forage legumes: May be used on wheat and barley crops undersown to the following (if not tank mixed with a broadleaf herbicide).

Alfalfa	Clovers
Bird's-foot trefoil	Sanfoin

Forage Grasses (seed production only)*:

Under-seeded with a cereal or grown alone (seedling or established):*

Bromegrass	Wheatgrass
(meadow, smooth)	(crested, intermediate)
Creeping red fescue	

Under-seeded with a cereal or grown alone (seedling only):*

Wheatgrass (northern, slender, western)

* *Liquid Achieve* and *Bison* only. NOTE - Since applications to these crops have been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Applications to these crops is at the risk of the user.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Wild oats - 1 to 6 leaf stage (total leaves including tillers), with a maximum of 2 tillers.

Volunteer tame oats - 1 to 6 leaf stage.

Green and yellow foxtail - 1 to 5 leaf stage (total leaves including tillers), with a maximum of 1 tiller.

Barnyard grass, Persian darnel - 1 to 4 leaf stage (total leaves including tillers).

For forage grasses and perennial cereal rye, apply prior to tillering of the above weeds.

Apply at the 2 to 3 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering.

Rates:

0.2 L per acre. One 8 L jug of *tralkoxydim* treats 40 acres.

Add *Turbocharge* or *Addit* adjuvant at a rate of 0.5 L per 100 L spray solution.

Maximum one application of these products or other products containing tralkoxydim per season.

If water analysis shows bicarbonate levels are 400 ppm or greater, add ammonium sulphate at 0.9 to 1.8 kg per 100 L of spray water prior to mixing.

Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 13

Application Information:

Water Volume:

Ground: 20 to 40 L per acre. Application in less than 20 L per acre water volume may result in mixing problems or unacceptable crop injury.

Aerial: 12 to 18 L per acre.

Nozzles and Pressure:

Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets. See the label for detailed instructions on aerial application.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Cereal crops that have set tillers may incur injury (yellowing and/or stunting) if applications are made within 48 hours of freezing temperatures. Cereal crops that have not set tillers may be injured if exposed to temperatures of 4°C or less up to 48 hours before or after application. Tank mixing with a broadleaf weed herbicide under adverse conditions may increase severity of crop injury. Crops under stress from foliar diseases or low fertility are more

susceptible to injury from application. Temporary crop injury may occur when tralkoxydim tank mixes (particularly dichlorprop/2,4-D ester products, and bromoxynil/MCPA ester products + additional MCPA Ester) are applied under extreme environmental conditions (dry or wet, cool or hot weather) resulting in crop stress. Control of grasses could be reduced when they are stressed due to drought, heat, lack of fertility, flooding or prolonged cool temperatures.

Tank Mixes:

For all tank mixes with tralkoxydim products, add *Turbocharge* or *Addit* adjuvant at a rate of 0.5 L per 100 L of spray solution.

Herbicides:

DO NOT tank mix tralkoxydim products with a broadleaf herbicide when applying to underseeded forage grasses or legumes.

Tank mix Partner	CROPS						
	Spring wheat	Durum	Winter wheat	Barley	Spring rye	Fall rye	Triticale
2,4-D ester (0.36 L/acre) [†]	•	•	•	•	•	•	
Bromoxynil ^{††} (0.4 L/acre)	•	•	•***	•		•***	•***
Bromoxynil/MCPA Ester ^{†††} (0.4 L/acre)*	•	•	•	•		•	
Curtail M (0.81 L/acre)	•	•		•			
Dichlorprop/2,4-D (0.71 L/acre)*	•	•	•	•			
Lontrel (0.11 L/acre) + MCPA ester (0.45 L/acre)	•	•		•			
MCPA ester (0.45 L/acre)	•	•	•	•	•	•	
Thumper (0.40 L/acre)*	•	•		•			
Trophy (20 acres/case)	•	•		•			

[†] 600 g/L formulations.

^{††} *Pardner* only for *Liquid Achieve* and *Marengo*.

^{†††} *Buctril M* only for *Marengo*; *Buctril M* and *Mextrol 450M* only for *Liquid Achieve*.

* Tank mixes may result in some temporary initial injury under adverse environmental conditions.

** Temporary crop injury can occur if applied prior to the 4 leaf stage. A reduction in wild oat control may occur with this mix.

*** *Buctril M* mixed with either *Liquid Achieve* or *Marengo* only in winter wheat, fall rye and triticale.

DO NOT tank mix tralkoxydim products with herbicides or formulations of herbicides not listed above as loss of grass control may result.

When applying broadleaf herbicides not listed above, in the same field, always apply tralkoxydim first. Apply the broadleaf product no sooner than seven days after application of tralkoxydim.

Insecticides:

Matador (49 mL/acre) (*Bison* and *Marengo* only).

Matador tank mixes with *Bison* may also be combined with bromoxynil or bromoxynil/MCPA ester products.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *tralkoxydim* labels only. To check for other possible registered mixes see the blue chart inside the back cover.

Various manufacturers may also support additional mixes that are not on the *tralkoxydim* labels. Check with manufacturers for more details.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour will reduce control.

Re-entry: DO NOT enter treated field for 12 hours.

Grazing: Straw from treated grain crops may be fed to livestock. Immature cereal crops may be grazed or cut for hay 16 days after treatment. DO NOT feed or graze forage crops in year of treatment

Preharvest: Leave 60 days from application to harvest.

Re-cropping: DO NOT replant treated areas to tame oat or corn for at least 4 weeks after application.

Aerial Application: May be applied by air to cereal crops only. DO NOT apply within 50 m of fish bearing waters and wildlife habitat.

Storage: Store in a dry place. DO NOT freeze.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:
	Terrestrial habitat
Ground*	3
Helicopter	80
Fixed wing aircraft	100

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

Caution – Skin and Eye Irritant

For an explanation of the symbols used here see page 11.

Traxos

Herbicide Group
1 - pinoxaden and clodinafop
(Refer to page 38)

Company:

Syngenta Canada

Formulation:

25 g/L pinoxaden and 25 g/L clodinafop propargyl formulated as an emulsifiable concentrate.

Container size - 2 X 10 L, 80 L, 400 L.

Crops and Staging:

Spring wheat (including durum) - prior to the emergence of the 4th tiller.

When tank mixing, check broadleaf product description for additional restrictions.

Weeds, Rates and Staging,

0.5 L per acre, no additional adjuvant required (packages treat 40 and 160 acres). Maximum one application of this product or those with the same ingredients in a season.

For control of:

WEED	STAGE
Barnyard grass, Persian dandel	1 to 5 leaves prior to tillering
Green and yellow foxtail	1 to 5 leaves, maximum 2 tillers
Volunteer canaryseed, Volunteer oats, wild oats, proso (Crown) millet	1 to 6 leaves, maximum 3 tillers

Optimum yield response occurs when weeds are controlled in early stages.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume:

Ground – Minimum 20 L up to 40 L per acre.

*Aerial** – Minimum 12 L/acre.

Nozzles and Pressure: 40 to 45 psi (275 to 310 kPa) when using conventional 80° or 110° flat fan stainless steel nozzles tilted forward at an angle of 45°. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE *coarse* droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance activity. Weeds hardened off by environmental stress such as cold weather, drought or excessive heat may not be adequately controlled.

Tank Mixes:

Herbicides:

*Buctril M** (label rates)

Curtail M (0.6 to 0.81 L/acre)

Infinity (0.33 L/acre)

MCPA 600 ester (0.28 to 0.37 L/acre)

Mextrol 450M (0.5 L/acre)

Pulsar (80 acres / case)

Pulsar (80 acres / case) + *MCPA 600 ester* (0.23 L/acre)

Trophy (20 acres per case)

Refer to the broadleaf herbicide label for crop staging, and other information.

Insecticides: *Matador* (25 to 33 mL/acre).

Fungicides: *Tilt* (0.1 L* to 0.2 L/acre).

Fertilizers: None registered.

* Aerial application approved.

Note: The above mixes are those listed on the *Traxos* label

only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Syngenta also supports the following mixes that are not on the *Traxos* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Attain XC* (low label rate), *Barricade II*, *Broadside*, *Momentum+MCPA ester*, *OctTain XL*, *Prestige XC*, *Pulsar* + *Express SG* (up to 6 g/acre), *thifensulfuron/tribenuron* + *MCPA*, *Retain*, *Stellar*

Fungicides: *Propel*, *Quilt*

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze or harvest treated crops for forage within 7 days of application.

Preharvest: Leave at least 60 days from application to harvest.

Re-cropping: No restrictions in the year following treatment.

Storage: Store in a cool, dry, ventilated area away from food or feed. Avoid ignition sources. If frozen, thaw and shake well before using.

Aerial Application: May be applied by air.

Buffer Zones:

Application method	Buffer Zones (metres†)		
	Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground	1	0	1
Aerial by airplane or helicopter	1	0	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:



Warning – Skin Irritant

For an explanation of the symbols used here see page 11.

Tribenuron

Herbicide Group

2 - tribenuron

(Refer to page 38)

Company:

E. I. duPont Canada (*Express SG*)

Nufarm Agriculture (*Spike* sold as *Spike-Up*)

Cheminova (*Nuance*)

Loveland Products (formerly Viterro) (*FirstStep* sold as *FirstStep Complete*)

Famers of North America (*MPower X*)

Arysta Lifescience (*Inferno WDG*)

Formulation:

Express SG (PCP# 28262): 50% tribenuron methyl, formulated as a water soluble granule (WSG).

75% WDG formulations (*FirstStep* - PCP# 29569;

Nuance - PCP# 29468; *MPower X* - PCP# 30964;

Inferno WDG - PCP# 30838; *Spike*[†] - PCP# 29653): 75% tribenuron methyl, formulated as a water dispersible granule (WDG).

Container sizes -

Express SG^{*}: 486 g.

MPower X^{*}; *Inferno WDG*^{*} and *Nuance*^{*}: 320 g.

FirstStep Complete: *FirstStep* (240 g) + *StartUp* (2 x 10 L)

Spike-Up: *Spike* (160 g, 3.6 kg Bulk) + *Credit 45* (16 L, 360 L Bulk)

[†] *FirstStep* and *Spike*: Sold as a prepackaged mix with the manufacture's glyphosate brand (see 'Company' section).

^{*} *Express SG*, *Inferno WDG*, *MPower X* and *Nuance* are purchased alone but must be used accordingly in combination with other registered tank mix herbicide before use.

Crops and Staging:

Tribenuron + glyphosate - 24 hours prior to seeding or summerfallow:

Alfalfa [†]	Oat [†]
Alsike clover [†]	Pea [†]
Barley	Red clover [†] (forage and seed production)
Canary seed [†]	Summerfallow
Dry bean [†]	Soybean [†]
Fababean [†]	Wheat (spring, durum, winter [†])
Lupin [†]	

NOTE: Injury to pulse crops may occur on coarse-textured

soils, low in organic matter (less than 3%), or in fields with variable soils, gravelly areas, sandy areas or eroded knolls. Avoid planting pulse crops in soils containing more than 50% sand.

Tribenuron + 2,4-D ester:

Summerfallow^{*}

Wheat (spring and durum), barley^{**} - 3 leaf up to emergence of the flag leaf.

Express SG[†] plus non-ionic surfactant:

Post-emergent in rangeland and pasture - stage according to weeds.

Express SG[†] plus *Hasten NT* adjuvant:

Tribenuron tolerant sunflower (eg. *ExpressSun* SU7 variety) - 2 to 8 leaves.

[†] *Express SG* only.

^{*} *Express SG*, *Inferno WDG*, *MPower X* and *Nuance* only.

^{**} *Inferno WDG*, *MPower X* and *Nuance* only.

Weeds, Rates and Staging,

Pre-seeding application and summerfallow mixed with glyphosate[‡]:

Express SG at 6 g per acre or 75% WDG tribenuron formulations at 4 g per acre plus glyphosate (any brand) at 180 g ae per acre (see glyphosate pages for equivalent rates for other formulations.)

Weeds controlled by glyphosate products at the rates above plus:

Canada thistle rosettes ^{**}	Scentless chamomile ^{^**}
Cow cockle [*]	White cockle (rosettes) ^{^**}
Dandelion (up to 6 inches)	Volunteer canola (including glyphosate tolerant varieties) ^{***}
Narrow-leaved hawk's-beard	

Summerfallow[‡]

Express SG 6 g per acre or *Inferno WDG*; *MPower X* and *Nuance* at 4 g per acre plus 2,4-D ester 170 g (6 oz.) ae per acre (eg. 0.24 L per acre LV 700 formulation):

Weeds controlled by 2,4-D ester 170 g (6 oz.) ae per acre plus:

Flixweed ^{♦♦}	Stinkweed ^{♦♦}
------------------------	-------------------------

Post-emergent in barley and spring wheat (including durum):

Nuance, Inferno WDG and MPower X only at 4 g per acre plus 2,4-D ester 170 g (6 oz.) ae per acre (e.g. 0.24 L per acre LV 700 formulation);

Weeds controlled by 2,4-D plus the following weeds up to 4 inches (10 cm) unless otherwise indicated:

Annual sunflower	Redroot pigweed
Canada thistle (top growth)	Wild buckwheat (1 to 3 leaf)**
Cow cockle	

Post emergent for season long control of the weeds below in Rangeland and Pasture only:

*Express SG only at 6 g per acre*** at the early bud – pre-bloom stage;*

Tall buttercup	Narrow-leaved hawk's-beard
----------------	----------------------------

*Express SG only at 12 g per acre****

The weeds listed above plus:

Dandelion	White cockle
Common tansy	

Post-emergent in Tribenuron Tolerant Sunflowers:

Express SG only at 6 g per acre (one 486 g package of Express SG treats 80 acres) plus Hasten NT adjuvant at 0.5L per 100L of spray solution will control;

Lamb's-quarters (up to 9 leaf)	Wild buckwheat** (up to 6 leaf)
--------------------------------	---------------------------------

^A Express SG only.

^{*} Up to the 3 leaf stage

^{**} Suppression only

^{***} Up to 6 inches

[♦] Allow 10 days between treatment and tillage.

^{♦♦} Fall rosettes and spring seedlings.

^{♦♦♦} Addition of a non-ionic surfactant at 0.2L per 100L of spray solution is required.

Tribenuron may degrade if left in the sprayer for an extended period. Apply within 24 hours of mixing.

Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 22 to 40 L per acre.

Nozzles and Pressure: Use appropriate pressure for nozzle. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets or larger. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Warm, moist growing conditions promote active weed growth and enhance the activity of tribenuron. Weeds hardened off by environmental stress such as cold weather, drought or excessive heat may not be adequately controlled.

Tank Mixes:

Herbicides:

Prior to seeding registered crops (all products):

Must be mixed with glyphosate.

Summerfallow (Express SG and 75% WDG formulations only):

Must be mixed with either glyphosate or 2,4-D ester.

In spring wheat (including durum) and barley (Nuance Inferno WDG and MPower X only):

Assert (0.53 to 0.65 L/acre);

*Fenoxaprop** (155 ml/acre).*

In spring wheat (NOT durum) and barley (Nuance; Inferno WDG and MPower X only):

Banvel II (44.5 mL/acre)

Tribenuron Tolerant Sunflowers (Express SG only):

Cletholin (77 mL/acre) plus Amigo adjuvant*

Poast Ultra (0.19 L/acre) plus Hasten adjuvant

** Select, Centurion or Shadow RTM only*

*** Cougar and Cordon only.*

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the tribenuron labels only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check product labels for directions.

General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours will reduce control.

Re-entry: DO NOT enter treated fields for 12 hours.

Grazing:

75% WDG formulations: DO NOT graze treated crops within 30 days of application.

Express SG: Forage may be grazed immediately following application.

Preharvest Interval:

75% WDG formulations: Leave 60 days between spraying and harvest of cereals.

Express SG: Leave 70 days between spraying and harvest of sunflower.

Re-Cropping: There are no restrictions one year after treatment.

75% WDG formulations: Canola, flax, lentil and alfalfa may be planted 2 months after application.

Express SG: Canola, flax, and lentil may be planted 2 months after application.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Fallow, preseed, range and pasture	1	0	4
Tribenuron tolerant sun-flowers	0	0	3

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

Tribenuron can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray *Tribenuron* should be flushed out immediately after use. Refer to 'Method A' in the general section on tank cleaning on page 15 to 16. **This ammonia rinse process should be done twice for the WDG formulations.** See the labels of the various products for specific instructions. The addition of detergent may improve cleanout, especially when mixing with other products.

Hazard Rating:

Express SG, FirstStep and Spike:



Warning – Eye Irritant
Potential Skin Sensitizer

Nuance, MPower X and Inferno WDG



Caution – Eye and skin irritant

All products:



Warning – Contains the allergens milk and sulfites

For an explanation of the symbols used here see page 11.

Trifluralin

Herbicide Group

3 - trifluralin

(Refer to page 38)

Company:

Dow AgroSciences (*Treflan*)

Nufarm Agriculture (*Rival*)

United Agri Products (*Bonanza*)

Formulation:

Bonanza 480 EC (PCP#28289): 480 g/L trifluralin formulated as an emulsifiable concentrate in 9.45 L, 205 L containers.

Bonanza 10G (PCP#22744): 10% trifluralin formulated as a granular in 22.7 kg, 500 kg bags.

Rival EC (PCP#18612): 500 g/L trifluralin formulated as an emulsifiable concentrate in 9 L, 110 L, 200 L, and 900 L containers.

Rival 10G (PCP#18926): 10% trifluralin formulated as a granular in 22.7 kg, 567 kg bags.

Treflan EC (PCP#23933): 480 g/L trifluralin formulated as an emulsifiable concentrate in 9.45 L, 115 L containers.

Crops and Staging:

Certain formulations are not registered for all the crops listed here. Refer to the specific product label for details. All products are for pre-plant incorporated use only.

Summerfallow use in the brown soil zone of Saskatchewan (Granular products only): Spring wheat (including durum).

Not for use in Manitoba.

Apply to summerfallow in May, June, or July for weed control during both years of a fallow-wheat rotation or in the fall (September or October) or spring prior to wheat seeding.

DO NOT apply to stubble when the previous crop was treated with another trifluralin product (*Treflan*, *Rival* or *Bonanza*). This includes application the previous summer or fall. DO NOT apply trifluralin to stubble or fallow when the previous year's crop was an oilseed, barley or pulse crop treated with a deep incorporated, spring or fall applied trifluralin product.

Green and Yellow Foxtail Control in Cereals:

Liquids applied in spring only (after seeding but prior to crop emergence) - spring wheat (including durum), barley.

Granulars applied in fall only (after September 1 but before freeze-up) - spring wheat (including semi-dwarf and durum).

Broadleaf and Grassy Weed Control in other crops:

Spring applied liquid or granular formulations:

Canola, pea, sunflower, safflower (liquid formulations), dry bean, mustard, fababean, alfalfa, sainfoin, sweet clover, soybean, forage legumes (cicer milk-vetch, seedling alsike clover, red clover, bird's-foot trefoil).

Fall applied granular formulations: Canola, pea, sunflower, dry bean, mustard, fababean, soybean, barley, lentil and flax.

Trifluralin liquids only: prior to planting shelterbelt transplants (elm, caragana, green ash, Scots pine).

Weeds:

Summerfallow use in the brown soil zone of Saskatchewan (Granular products only):

Fallow Year:

Barnyard grass	Redroot pigweed
Cow cockle	Russian thistle*
Green foxtail	Wild buckwheat
Lamb's-quarters	Wild oat†
Persian dandelion	

Crop Year:

Green foxtail	Wild buckwheat *
Lamb's-quarters	Wild oat *

* Suppression only

Green and Yellow Foxtail Control in Cereals:

Foxtail (green and yellow)

Broadleaf and Grassy Weed Control in other crops:

Annual brome species (downy, Japanese)	Lamb's-quarters
Barnyard grass	Persian dandelion
Chickweed	Pigweed
Cow cockle	Purslane
Green foxtail	Wild buckwheat*
Knotweed	Wild oats†*
	Yellow foxtail

* Some plants may escape herbicide treatment but are not competitive with the crop.

† Suppression with *Treflan EC* and only.

Rates and Staging:

Summerfallow use in the brown soil zone of Saskatchewan (Granular products only):

DO NOT apply to sandy soils with less than 1% organic matter. Application to severely eroded knolls is not recommended. DO NOT apply to wet soils, soils in poor working condition, soils which contain more than 8 percent organic matter, or soils subject to prolonged periods of flooding.

Granules may be applied to standing or pre-worked stubble, provided trash or green growth does not interfere with cultivation (prevent soil mixing).

Over-application caused by overlapping, improper calibration or non-uniform application may result in reduced crop stand, delayed development or reduced yields.

Soil Organic Matter (%)	RATE (KG/ACRE)	
	1 to 3	4 to 8
May	3.85	4.5
June	3.25	3.85
July	2.65	3.25
	2 to 8% Organic Matter	
September to October	2.23*	

* Control of green foxtail only, on soils between 2 to 8% organic matter.

Broadleaf and Grassy Weed Control in other crops:

For use in canola, pea, sunflower, dry bean, mustard, fababean, seedling alfalfa (spring only), seedling sweet clover (spring only), soybean.

PRODUCT	SOIL TYPE			
	Light soils with less than 6% organic matter		Medium to heavy soils with 6 to 15% organic matter	
	Spring	Fall	Spring	Fall
Rival EC	0.65 L/acre	0.89 L/acre*	0.89 -1.13 L/acre	1.13 -1.37 L/acre*
Rival 10G	3.43 kg/acre**	4.45 kg/acre	4.45 -5.67 kg/acre**	5.67 -6.88 kg/acre
Treflan EC	0.69 L/acre	0.93 L/acre*	0.93 -1.21 L/acre	1.21 -1.37 L/acre*
Bonanza 10G	Not registered	4.45 kg/acre	Not registered	5.67 -6.88 kg/acre
Bonanza 480 EC	0.69 L/acre	0.93 L/acre*	0.93 L/acre	1.17 L/acre*

* Although liquid formulations are registered for fall application, this use is not recommended as tillage requirements before and after application will predispose fields to erosion.

** Spring applications of granular formulations are recommended for Manitoba only.

During the fallow year, susceptible weeds may not be fully controlled until after the second fallow operation has established a uniform layer of treated soil. Control of wild oats in the crop year may be variable depending on wild oat population as well as soil and climatic conditions. Some wild buckwheat may escape but its growth will be retarded and result in limited competition to the wheat crop.

Pre-emergent control of green and yellow foxtail:

Liquids

PRODUCT	RATES (L/ACRE)	
	LIGHT AND MEDIUM SOIL TEXTURE	HEAVY SOIL TEXTURE
Rival EC	0.49 to 0.57 L	0.65 L
Treflan EC, Bonanza 480 EC	0.49 L	0.69 L

Granulars products (wheat only)

Rival 10G/Bonanza 10G at 2.23 kg per acre in all soil textures with 2 to 8 % organic matter.

For use in barley (fall only), apply:

PRODUCT	SOIL TYPE					
	2 to 4% organic matter		4 to 6% organic matter		6 to 10% organic matter	
	Light Soil Texture*	Medium to Heavy Soil Texture**	Light Soil Texture*	Medium to Heavy Soil Texture**	Light Soil Texture*	Medium to Heavy Soil Texture**
<i>Rival 10G, Bonanza 10G</i>	3.44 kg/acre	3.44 kg/acre	4.45 kg/acre	4.45 kg/acre	4.45 kg/acre	5.67 kg/acre

* Light textured soils can be defined as sandy to sandy-loam.

** Medium to Heavy textured soils can be defined as loam to clay.

For use in flax or lentils (fall only), apply:

PRODUCT	SOIL TYPE			
	Soils with 2 to 6% organic matter		Soils with 6 to 15% organic matter	
	Light Soil Texture*	Medium-Heavy Soil Texture**	Light Soil Texture*	Medium-Heavy Soil Texture**
<i>Rival 10G, Bonanza 10G</i>	4.45 kg/acre	4.45 to 5.6 kg/acre***	5.67 kg/acre	5.67 to 6.88 kg/acre
<i>Bonanza 480 EC</i>	0.93 L/acre	0.93 L/acre	1.17 L/acre	1.17 L/acre
<i>Treflan EC</i>	0.93 L/acre	1.21 L/acre	1.21 L/acre	1.21 - 1.38 L/acre
<i>Rival EC</i>	0.89 L/acre	1.13 L/acre	0.89 L/acre	1.13 - 1.38 L/acre

* Light textured soils can be defined as Sandy to Sandy-loam.

** Medium to Heavy textured soils can be defined as loam to clay.

*** Rates vary among products. Refer to product label for specific information.

Application:

Liquid Formulations:

Water Volume: Minimum 40 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets or larger.

Dry Granular Formulations: Use equipment capable of metering granular herbicides and applying in an even layer over the surface of the soil. Close applicator lid after filling to avoid prolonged exposure to direct sunlight.

Incorporation:

Summerfallow use in the brown soil zone of Saskatchewan:

Apply granules to the soil surface and incorporate immediately, in the same operation if possible. DO NOT delay incorporation more than 24 hours after application. Use a

deep tillage cultivator, field cultivator or disc implement set to work 2 to 3 inches (5 to 8 cm) deep, and operating at 8 to 10 km/hr. Granules should not be incorporated when soil is crusted, lumpy or too wet for good mixing action.

May - July: A second incorporation at the same depth and at an angle to the first should be done when weed growth requires it. Wait at least one week before making the second incorporation. After completing two fallow incorporations, additional operations with a rod weeder, shallow tillage or fall 2,4-D application may be required to control remaining weed growth.

September - October: A second incorporation may be done in the fall a minimum of 3 days later. Alternatively, to conserve trash cover through the winter, the second incorporation can be completed in the spring at the same depth and at an angle to the first incorporation. When both incorporations take place in the fall, shallow spring tillage should be completed in the spring. If a discer or air seeder is used for seeding, separate spring tillage may not be necessary.

NOTE: Fall application is not recommended on soils where a lack of trash cover combined with the required incorporation would leave the soil vulnerable to erosion.

Spring (In the year of seeding): Apply granules and incorporate immediately, in the same operation if possible. DO NOT delay the first incorporation longer than 24 hours after application. The second incorporation must be delayed a minimum of 3 days following the first incorporation. When applied to cold soils, wait 14 days before making second incorporation.

The second incorporation should be done at an angle to the first incorporation, and at the same depth. If a discer or air seeder is used for seeding, the seeding operation can be used as the second incorporation.

Green and Yellow Foxtail control in Cereals:

Liquid formulations: Apply and incorporate in spring just after seeding. Incorporate to a depth of 1 to 1.5 inches (2 to 4 cm) into a trash free soil (80 percent black when viewed from above) using diamond or tine type harrows operated at a speed of 6 mph (9 km/h). Incorporate twice, with the second incorporation at right angles to the first. The first incorporation should be performed immediately in the same direction of application. Both incorporations should be done within 24 hours of application. When tank mixing liquid formulations with *Avadex BW*, follow the same incorporation procedure.

Granular formulations: May be applied to standing or pre-worked stubble. Very heavy trash fields should be worked prior to application to allow product penetration to the soil surface. Incorporate with cultivators or disc implements only. Perform the first tillage operation within 24 hours of application. Incorporate at a working speed of 5 to 8 mph (8 to 13 km/hr) and to a depth of 2 to 3 inches (5 to 8 cm). Wait a minimum of 5 days, then incorporate a second time at right angles to the first. This second incorporation may be delayed until the following spring. Subsequent working should be no deeper than 2 to 3 inches (5 to 8 cm).

Broadleaf and Grassy Weed Control in other crops:

Granular formulations are recommended for use in fall or spring as a pre-plant incorporated treatment on broadleaf crops listed on the product label. The liquid formulations should be used only on soils free of lumps and relatively trash-free (75% black) and are recommended only for spring use. Granular formulations may be applied to standing or pre-worked stubble. Very heavy trash fields should be worked prior to application to allow product penetration to the soil surface. DO NOT use liquid or formulations of trifluralin as a pre-plant incorporated treatment in barley, as severe injury will result. Only the fall applications of *Rival 10G*, and *Bonanza 10G* are registered for use as pre-plant incorporated treatments in barley. For fall application of granular formulations, work the chemical into the soil between September 1 and freeze-up. Use a discer or field cultivator (vibrating shank-type). Disc implements are preferred on stubble. Set equipment to cut at 3 to 4 inches (8 to 10 cm) depth. The initial incorporation should be done within 24 hours of application.

The second incorporation should be done at right angles to the first. The second incorporation may be delayed

until spring, except when planting barley, flax or lentils; for these crops both incorporations must be done in fall. Delay the second incorporation 5 days for better weed control. This will allow greater release of the chemical onto soil particles and assure more even distribution. Fall application of granular trifluralin on flax, lentils or barley is not recommended on soils prone to erosion, as the 2 fall incorporations necessary in these crops may leave soils vulnerable to wind or water erosion.

For spring application of liquid and granular formulations, work the chemical into the soil prior to seeding by setting the implement at 3 to 4 inches (8 to 10 cm) cutting depth. The first incorporation must be done within 24 hours of application. The second incorporation must be done at right angles to the first. If incorporating granular trifluralin, delay the second incorporation for 3 days after the first to achieve better weed control.

Seeding:

Summerfallow use in the brown soil zone of Saskatchewan: Allow soil to warm before seeding to reduce risk of injury to crop. Place seed 1.25 to 2.5 inches (3 to 6 cm) deep. If spring seedbed preparation is required, set cultivator 2 inches (5 cm) deep. To reduce the risk of wheat injury, use good quality seed and agronomic practices that will promote good growing conditions. Avoid deep seeding, loose seedbeds and seeding into cold soils. If extended dry periods were present after a fallow application, a 10 percent increase in seeding rate is recommended.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Prolonged drought conditions after a May-July application to fallow may result in higher levels of trifluralin in the soil at the time of seeding.

Injury to flax, barley, wheat or lentil may occur if soil and weather conditions are not conducive to rapid crop emergence (cold or dry soils at the time of seeding and crop emergence).

To minimize crop injury, seed into a firm, moist seed bed using a seeder with good depth control and on row packing. Plant barley no deeper than 2 inches (5 cm). Plant cereals, lentil and flax no deeper than 1.5 inches (4 cm).

Less than acceptable weed control will result if dry conditions prevail at the time of weed emergence.

Rainfall has no direct effect on products' activity. Flooding (3 to 5 days) will cause rapid breakdown of the product resulting in reduced weed control. Flooding for 3 weeks or more will result in total breakdown of the product resulting in loss of weed control.

Tank Mixes:

Herbicides:

Soybeans:

Sencor (Treflan EC only).

Fertilizers: Liquid product may be applied with liquid fertilizer as a carrier. Before the herbicide is added to the tank, compatibility of the herbicide to liquid fertilizer should be tested following instructions on the herbicide container. *Trifluralin* liquids may be blended with dry bulk fertilizers (DO NOT mix with nitrate fertilizers). Check label for blending instructions.

Insecticides: None registered.

Fungicides: None registered.

Note: The above mixes are those listed on the *Trifluralin* labels only. To check for other possible mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No restriction. Flooding may reduce weed control.

Re-entry: Wait at least 12 hours before entering treated fields.

Grazing: DO NOT graze the treated crops or cut for feed prior to crop maturity.

Re-cropping: Oat, canaryseed, and small-seeded grasses may be affected the year after treatment. Corn is sensitive at higher rates of application. Damage to wheat can occur if the crop is seeded into land that has been treated during the previous 21 months with trifluralin products and has received abnormally low amounts of precipitation. Damage is worse if conditions are not conducive to rapid emergence of the wheat (for example, if the crop is seeded deep or if soil conditions remain cool during emergence). Damage tends to be greater on fields treated with granular formulations.

Aerial Application: DO NOT apply by air.

Storage: Granular formulations must be stored in a cool, dry location, out of sunlight.

Rival EC: DO NOT store below 5°C.

Treflan EC: DO NOT freeze.

Bonanza 480 EC: DO NOT freeze.

Recommendations for liquid formulations: Crystallization of the active ingredient may occur at less than 5°C. To reconstitute, bring temperature to 15°C and shake well until no crystals are visible. This should be done before adding to the spray tank.

Buffer Zones: (liquid formulations only)

CROP	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Field crops	80	10	1
Shelterbelts, woody crops	120	20	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Tank Cleaning:

Refer to 'Method B' in the general section on tank cleaning on page 15 to 16.

Hazard Rating:

Bonanza 480 EC:



Warning – Poison



Warning – Eye and Skin Irritant

All products:

Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Triton C

Herbicide Group
2 - thifensulfuron & tribenuron
4 - quinclorac
(Refer to page 38)

Company:

E. I. du Pont Canada Inc. (PCP# 28622)

Formulation:

51.55 % quinclorac + 10.30 % thifensulfuron methyl + 5.15 % tribenuron methyl formulated as a water dispersible granule.

Container size - 1.566 kg container.

Crops and Staging:

Spring barley, wheat (including durum, and spring) - 2 to 5 leaf stage.

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions

Weeds and Staging:

Unless otherwise noted below, apply to young and actively growing weeds that are less than 4 inches (10 cm) in height or width.

Weeds Controlled:

Annual smartweed (green, lady's-thumb)	Redroot pigweed
Ball mustard	Round-leaf mallow (2 to 6 leaf)
Chickweed (1 to 6 leaf)	Russian thistle
Cleavers (1 to 4 whorls)	Shepherd's-purse
Common groundsel	Sow-thistle, annual
Corn spurry	Stinkweed
Cow cockle	Stork's-bill (2 to 6 leaves)
Flixweed	Tartary buckwheat
Hemp-nettle	Volunteer canola (not CLEARFIELD varieties)
Lamb's-quarters	Volunteer sunflowers
Narrow-leaved hawk's- beard	Wild buckwheat (1 to 5 leaf)
	Wild mustard

Weeds Suppressed:

Canada thistle, perennial sow-thistle (less than 6 inches (15 cm) tall or across and prior to budding)

Scentless chamomile

Toadflax (less than 6 inches or 15 cm tall)

Volunteer flax

Rate:

39.25 g per acre. Limit to one application of this product or other products containing the same ingredients per year. (One 1.566 kg container treats 40 acres)

Merge adjuvant (not included) must be added at 1.0 L per 100 L of spray solution.

Refer to the product label for complete mixing instructions. A general guide to mixing can be found on page 13.

Application Information:

Water Volume: Minimum 22 L per acre.

Nozzles and Pressure: 30 to 40 psi (210 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of ASABE medium droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to wheat, or barley that are stressed by severe weather conditions (frost, drought or water saturated soil) as crop injury may result. Under certain conditions (heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) lightening in crop colour and reduction in crop height may occur.

Group 2 susceptible kochia control may be reduced during stress conditions or if extremely heavy infestations exist.

Tank Mixes:

DO NOT mix with substances that contain boron or that release chlorine.

Herbicides:

In spring wheat (NOT including durum) and barley only:
Axial 100EC (0.24 L/acre) plus *Adigor* adjuvant*.
Axial 100EC (0.24 L/acre) + *MCPA ester* (0.19 L/acre)** plus *Adigor**

* Merge adjuvant is NOT required with this mix.

** 600 g/L formulation

Fertilizers: None registered.

Note: The above mixes are those listed on the *Triton C* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

E.I.duPont also supports the following mixes that are not on the *Refine SG* label. Mixes must be applied according to the most restrictive use limitations for either product:

Herbicides: *Avenge*, *Axial BIA*, *Puma Advance*

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 6 hours may reduce control in general.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: Must not be grazed or fed to livestock for 77 days after treatment.

Preharvest Interval: Leave 77 days between treatment and harvest for wheat and durum and 80 days for barley.

Re-cropping: Spring wheat (including durum) and spring barley may be reseeded immediately following application. Wheat, barley, oat, canola, field pea, flax, lentil and sunflower may be grown the year after application. On low organic matter soils or under dry conditions, flax and lentils should NOT be grown until the second year after application. DO NOT use *Triton C* on land where potato or vegetables are grown. A field bioassay (a test strip grown to maturity) must be conducted the year before growing any crops other than those listed above.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	0	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Triton C can cause severe injury to sensitive crops at very low concentrations. Sprayers should be flushed out immediately if application is to be stopped for an extended period. The manufacturer recommends a cleanout procedure similar to 'Method A' in the general tank cleaning section on page 15 to 16. See label for specific process.

Hazard Rating:



Caution – Poison



Warning – Eye and Skin Irritant
 Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Triton K

Herbicide Group

2 - tribenuron

4 - dicamba, 2,4-D

(Refer to page 38)

Company:

E. I. duPont Canada

Formulation:

Triton K package contains the following components:

Express SG (PCP# 28606)*: 50% tribenuron methyl formulated as a water soluble granule.

Container size - 243 g bottle.

Plus

Banvel II (PCP# 23957)*: 480 g/L dicamba diglycolamine salt, formulated as a solution.

Container size - 1.8 L

-or-

DB-858 (PCP# 28872): Triton Broadleaf (PCP# 29989): 58.45% dicamba sodium salt, and 8.25% tribenuron methyl formulated as a water dispersible granule.

Container size - 1.47 kg

-Plus-

Both versions contain:

2,4-D LV 700 (PCP# 23192): 660 g/L 2,4-D ester formulated as an emulsifiable concentrate.

* *The original formulation of Triton K is a tank mix of Express SG (page 271), 2,4-D ester (page 62) and Banvel II (page 121). This formulation is no longer being manufactured, but supplies still remain in the distribution system. The older formulation will be removed from future editions of this publication when supplies are exhausted.*

Crops and Staging:

Spring wheat (including durum), winter wheat and barley:

3 leaves fully expanded to 6 leaves plus 3 tillers. Application outside of this stage range can result in injury to the crop.

Summer-fallow: Stage according to weeds.

* *DB-858 / Triton Broadleaf only.*

Weeds and Staging:

Weeds controlled up to 10 cm tall or across:

Annual sunflower	Prickly lettuce
Ball mustard	Redroot pigweed
Canada thistle (top growth control)	Russian pigweed
Cow cockle	Russian thistle
Dandelion ***	Shepherd's-purse**
Flixweed**	Stinkweed**
Hare's-ear mustard	Sweet clover
Indian mustard	Thyme-leaved spurge
Kochia (2 to 10 leaf)†	Tumble mustard
Lamb's-quarters	Wild buckwheat*
Narrow-leaved hawk's-beard**	Wild mustard
	Wild radish
	Wormseed mustard

* 1 to 4 leaf stage

** Fall rosettes and spring seedlings only.

*** Spring or fall rosettes up to 15 cm in diameter.

Rate:

Depending on formulation available add either:

Express SG: 6 g per acre, and;

Banvel II: 44.5 mL per acre

-or-

DB-858/Triton Broadleaf: 36.8 g per acre

Plus with either of the options above add:

2,4-D 700 LV ester: 243 mL per acre

(One package treats 40 acres or 16 ha)

Apply this product or other products containing the same ingredients only once per season.

Triton K may degrade if left in the sprayer for an extended period of time. Apply within 24 hours of first mixing. Refer to the product label for complete mixing instructions.

A general guide to mixing can be found on page 14.

Application Information:

DO NOT apply if temperatures are greater than 30°C, if humidity is high, or wind is blowing toward non-target plants as injury from drift may result.

Water Volume: Minimum 22 L per acre.

Nozzles and Pressure: Maximum 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE coarse* droplets. Use a 50 mesh or coarser screen and filter system.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to wheat, or barley that are stressed by severe weather conditions (frost, drought or water saturated soil) as crop injury may result. Under certain conditions (heavy rainfall, prolonged cool weather, frost conditions, wide fluctuations in day/night temperatures) lightening in crop colour and reduction in crop height may occur.

Kochia control may be reduced during stress conditions or if extremely heavy infestations exist.

Tank Mixes:

None registered.

Note: The above mixes are those listed on the *Triton K* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 4 to 6 hours may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: Lactating dairy animals MUST NOT graze fields with 7 days of treatment.

Preharvest Interval: Leave 30 days between application and harvest.

Re-cropping: No restrictions the year following application.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. May be frozen.

Buffer Zones:

CROP	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Cereals	1	0	4
Fallow	1	1	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance is measured from the downwind edge of the boom to sensitive areas.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

Triton K can cause severe injury to sensitive crops at very low concentrations. Sprayers used to spray this product should be flushed out immediately after use. The manufacturer recommends a process similar to 'Method A' in the general section on tank cleaning on page 15 to 16. see label for specific instructions.

Hazard Rating:



Caution – Poison



Warning – Eye and Skin Irritant
Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Trophy

Herbicide Group
4 - fluroxypyr & MCPA
(Refer to page 38)

Weed Control

Company:

Nufarm Agriculture

Formulation:

The *Trophy* package has 2 components:

Nufarm fluroxypyr herbicide (PCP#30194): 180 g/L fluroxypyr.

MCPA Ester 600 (PCP# 27803): 600 g/L MCPA ester.

Container size - *Trophy A* - 4.8 L, *Trophy B* - 7.5 L

All components above are formulated as emulsifiable concentrates.

All components above are formulated as emulsifiable concentrates.

Crops and Staging:

Spring wheat (including durum), canaryseed* & barley: - 3 leaf up to to full emergence of the flag leaf.

**Since the use of this product on canaryseed is registered under the User Requested Minor Use registration system, the manufacturer assumes no responsibility for herbicide performance. Users of this product on canaryseed do so at their own risk.*

When tank mixing, always check the tank mix partner recommendations for additional staging restrictions.

Weeds and Staging:

Weeds controlled at the 2 to 4 leaf stage, unless specified include:

Burdock	Ragweed (common)
Cleavers (1 to 4 whorls)	Redroot pigweed
Cocklebur	Shepherd's-purse
Flixweed	Stinkweed
Hemp nettle (2 to 6 leaf)	Sunflower (annual)
Kochia	Vetch
Lamb's-quarters	Volunteer canola
Mustards (except dog and tansy)	Volunteer flax (1 to 12 cm)
Prickly lettuce	Wild mustard
	Wild radish

Weeds suppressed include:

Smartweed (green)

Wild buckwheat (1 to 4 leaf)

Stork's-bill (1 to 8 leaf)

Rate:

Nufarm fluroxypyr herbicide: 0.24 L per acre

MCPA Ester 600: 0.38 L per acre.

(One case treats 20 acres)

Make only one application of this product or other products containing fluroxypyr per year.

Application Information:

Water Volume: Minimum 40 L per acre.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with *ASABE coarse* droplets

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Trophy activity is influenced by weather conditions. The temperature range for optimum activity is 12°C to 24°C. Reduced activity will occur when temperatures are below 8°C or above 27°C. Frost 3 days before or after application may reduce weed control and crop tolerance. Weed control may be reduced during stress conditions (drought, heat or cold stress) or if extremely heavy infestations exist.

Tank Mixes:

Herbicides:

In spring wheat (including durum) and barley:

Achieve Liquid (0.2 L/acre) plus *Turbocharge* adjuvant.

Assert (0.53 to 0.65 L/acre) plus pH adjuster.

Insecticides: None registered.

Fungicides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Trophy* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: No rainfast period is specified on the label; required interval may be up to 8 hours. Contact manufacturer for more information.

Re-Entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze, harvest forage or cut hay within 7 days of application. Remove meat animals from treated fields at least 3 days before slaughter.

Preharvest: Leave 60 days from application to harvest.

Re-cropping: Wheat, barley, oat, rye, forage grasses, flax, canola, mustard, lentil and pea may be grown the year after application. There are no re-cropping restrictions the second year after application.

Aerial Application: DO NOT apply by air.

Storage: May be frozen. If frozen, bring to room temperatures and agitate before use.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	15	15	15

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Hand-held or backpack sprayers, inter-row hooded sprayers and spot treatments are exempt from buffer zone requirements.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack applications do not require a buffer.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Danger – Poison.



Warning – Eye Irritant.



Caution – Skin Irritant.

For an explanation of the symbols used here see page 11.

Tundra

Herbicide Group

1 - fenoxaprop

6 - bromoxynil

27 - pyrasulfotole

(Refer to page 38)

Company:

Bayer CropScience (PCP#29367)

Formulation:

46 g/L of fenoxaprop-p-ethyl, 87.5 g/L of bromoxynil and 15.5 g/L of pyrasulfotole formulated as an emulsifiable concentrate.

Container size - 8.1 L, 129.6 L and 405 L.

Crops and Staging:

Application beyond the maximum rates provided below may result in crop injury.

CROP	STAGE
Barley, Spring wheat (including durum)	1 to 6 leaves on the main stem plus 3 tillers

Weeds and Staging:

Apply 0.81 L per acre (one 8.1 L container treats 10 acres) to control;

Grass weeds from the 1 to 6 leaf stage up to emergence of 3rd tiller:

Barnyard grass Wild oat
Foxtail (green and yellow)

Apply at the 3 to 4 leaf stage for optimum control. Optimum weed control and yield response occurs when weeds are removed before tillering.

Broadleaf weeds from the 1 to 6 leaf stage unless otherwise indicated:

Annual sow-thistle	Perennial sow-thistle*
Chickweed	Ragweed (Common)
Canada thistle* (up to 30 cm)	Redroot pigweed
Cleavers (1 to 3 whorls)	Round leaf mallow*
Cleavers (4 to 6 whorls)**	Russian thistle (up to 10 cm)
Dandelion*	Shepherd's-purse
(up to 25 cm across†)	Stinkweed
Flixweed (up to 10 cm)	Volunteer canola***
Hemp-nettle	Wild buckwheat
Kochia (up to 10 cm)	Wild mustard
Lamb's-quarters	
Pale smartweed	

* Suppression only

** To control cleavers at 4 to 6 whorls add ammonium sulfate at 202 g per acre (99% dry) or 0.4 L per acre (49% solution). If using an ammonium sulphate product with a different concentration, adjust the rate accordingly.

*** All herbicide tolerant varieties.

† Spring seedlings and over-wintered rosettes.

DO NOT apply *Tundra* or other products containing fenoxa-prop, pyrasulfotole or bromoxynil more than once in the same year.

Application Information:

Water Volume:

Ground: 18.9 L per acre. Use higher water volumes for dense crop/weed canopies.

Aerial: 11.4 L per acre.

Nozzles and Pressure: Use a combination of nozzles and pressure designed to deliver thorough, even coverage with

ASABE medium classification droplets. Low drift nozzles may require higher pressures for proper performance.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Crop injury may result if applied to a crop that is stressed by severe weather conditions, frost, low fertility, drought, water-saturated soil, disease or insect damage. Weeds growing under adverse environmental conditions such as drought will be less susceptible to *Tundra*. Under stressed conditions and/or heavy crop canopy, early application will result in improved weed control.

Tank Mixes:

None registered.

Bayer also supports the following mixes that are not on the *Tundra* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: 2,4-D Ester, Lontrel, MCPA Ester.

Fungicides: Tilt

Insecticides: Decis, Sevin XLR.

Adding ingredients in the correct order is critical for optimum performance. Check labels of both products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-Entry: DO NOT enter treated areas for 24 hours.

Grazing: DO NOT graze or cut cereal crops for hay, within 25 days of application.

Preharvest Interval: Leave 65 days from application to harvest.

Re-cropping: Alfalfa, barley, canaryseed, canola, corn (Manitoba only), flax, oat, potato, soybean (Manitoba only), sunflower, tomato (Manitoba only), and wheat (spring, and durum) may be planted the season following application. Field pea may be grown the following year in all black, grey-wooded and dark brown soil zones. DO NOT plant field pea the season following *Tundra* use in the brown soil zone where organic matter content is below 2.5 % and where soil pH is above 7.5. Lentil may be seeded the second season following application.

Aerial Application: May be applied by air.

Storage: Store in a dry controlled temperature facility. DO NOT freeze. Shake before using if stored for longer than one year.

Buffer Zones:

Application method	Buffer Zones (metres †) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	3	1	10
Fixed wing aircraft	20	5	375
Helicopter	20	3	225

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general section on tank mixing on page 15 to 16.

Hazard Rating:

Caution – Poison



Danger – Corrosive to eyes and skin.

Potential skin sensitizer.



Warning – Eye Irritant.

For an explanation of the symbols used here see page 11.

Ultim

(For use only in Manitoba)

Herbicide Group
2 - rimsulfuron & nicosulfuron
(Refer to page 38)

Company:

E. I. duPont Canada (PCP#24736)

Formulation:

37.5% rimsulfuron + 37.5% nicosulfuron formulated as a water dispersible granule.
 Container sizes - 134.8 g (4 x 33.7 g water soluble bags).

Crops and Staging*:

The following field corn hybrids at the 1 to 4 leaf stage:

Pioneer Brands 39K72, 39K73, 39W54, 39M27. For use in Manitoba only.

**NOTE - Since applications to corn in Manitoba has been registered under the User Requested Minor Use program, the manufacturer assumes no responsibility for herbicide performance. Application to corn is at the risk of the user.*

Weeds and Staging:

No information is provided on the label for leaf staging. The manufacturer recommends the following staging:

CROP	STAGE
Wild oats	3 to 6 leaf
Foxtail (green and yellow), barnyard grass, volunteer cereals	1 to 6 leaf (up to 2 tillers)
Quackgrass	3 to 6 leaf stage (with extended leaf 4 to 8 inches long)
Redroot pigweed	2 to 6 leaf

Contact the manufacturer for additional weeds not listed on the label.

Rates:

13.5 g per acre.

One water soluble bag of *Ultim* will treat 2.5 acres (10 acres per container).

Add a non-ionic surfactant (*AgSurf II*, *Agral 90*, *Citowett Plus*) at 0.2 L per 100 L of spray solution.

Apply *Ultim* within 24 hours of mixing, as product degradation may occur resulting in reduced weed control. Refer to the product label for complete mixing instructions.

Application Information:

Water Volume: Minimum 40 L per acre; for best results apply 56 to 77 L per acre.

Nozzles and Pressure: No pressures listed on label when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage of *ASABE medium* droplets. Use a 50 mesh or coarser screen and filter system.

Effects of Growing Conditions:

Rapid fluctuations in temperature (greater than 20°C difference within 24 to 36 hours) will stress the corn crop. For maximum crop safety, allow 48 to 72 hours for the corn to acclimatize before applying *Ultim*.

Apply **ONLY** when the temperature in the 24 hours before AND after application is between 5°C and 28°C. Temperatures beyond this range increase the potential for crop injury. Separate applications of *Ultim* herbicide followed by a broadleaf herbicide (minimum 12 hours later) will reduce the potential for injury.

WARNING: Crop injury may result if application is made to corn that has been stressed by abnormally hot, humid or cold weather conditions, frost, low fertility, drought, water saturated soil, compacted soil, previous pesticide applications, disease or insect damage. If corn has been injured by frost, wait 48 to 72 hours before applying *Ultim*.

Tank Mixes:

Herbicides: None registered.

Insecticides: None registered. *Ultim* **should NOT** be applied to corn that has been treated with *Thimet*, or *Lorsban*. Leave 7 days between the application of *Ultim* and that of a foliar organophosphate insecticide.

Fungicides: None registered.

Note: The above mixes are those listed on the *Ultim* label only. To check for other possible registered mixes see the blue fold out chart inside the back cover.

Restrictions:

Rainfall: Within 2 to 4 hours may reduce control.

Re-entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze treated crops or cut for hay.

Preharvest: Leave 30 days from application to harvest.

Re-cropping: Field corn, winter wheat and spring barley may be planted the year following application. Perform a field bioassay before planting any other crops, or where *Ultim* is more persistent (sandy soils, with low organic matter and pH greater than 7).

Aerial Application: DO NOT apply by air.

Storage: Store product in original containers in a secure, dry area, away from other pesticides, food, or feed.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground only*	1	1	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.


† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.


Tank Cleaning:

The manufacturer recommends a cleanout process similar to 'Method A' in the general section on tank cleaning on page 15 to 16. See the label for specific instructions.

For additional information, Refer to page 15.

Hazard Rating:

 Warning – Eye Irritant.

 Caution – Skin Irritant.

For an explanation of the symbols used here see page 11.



Herbicide Group
2 - thien carbazone
(Refer to page 38)

Company:

Bayer CropScience (PCP# 29070)

Formulation:

10 g/L thien carbazone-methyl formulated as a suspension concentrate.

Container size - 2 x 8 L jug.

Crops and Staging:

Spring wheat (including durum):

1 to 6 main stem leaf stage to a maximum of 3 tillers, and before the first node can be felt in the stem. DO NOT apply beyond 35 days of emergence.

Winter wheat:

Spring or fall application from 1 to 6 main stem leaf stage and before the first node can be felt in the stem.

DO NOT apply after the presence of the first node as crop injury may occur.

Weeds and Staging:

Grass weeds controlled from 1 to 6 main stem leaves and prior to the emergence of the 3rd tiller unless otherwise indicated:

Barnyard grass	Persian darnel†
Foxtail (green and yellow†)	Volunteer Canaryseed *
Japanese brome† **	Wild oats

Broadleaf weeds controlled at the 1 to 6 leaf stage unless otherwise indicated:

Cleavers (1 to 6 whorls)	Shepherd's-purse
Hemp-nettle	Stinkweed
Lamb's-quarters†	Volunteer canola
Pale smartweed	(except Clearfield varieties)
Pigweed, redroot	Wild buckwheat
Round-leaved mallow†	Wild mustard
Russian thistle (up to 10 cm)†	

* Up to the emergence of the 2nd tiller.

** Prior to tillering.

† Suppression only.

Rates:

0.2 L per acre

(One 8 L container will treat 40 acres)

Add either Agral 90, AgSurf II, Surf 92, at 0.2 L per 100 L on spring wheat or durum only;

or,

Add ammonium sulphate on spring wheat only for improved weed control. Add ammonium sulphate to the tank first at 202 g/acre (99% solution) or 0.4 L/acre (49% solution).

Application Information:

Water Volume:

Ground: 20 to 40 L per acre. Use higher water volumes for dense canopies.

Aerial: Minimum 11.3 L per acre.

Nozzles and Pressure:

Ground: For conventional flat fan nozzles use a pressure of 30 to 50 PSI (207 to 345 kPa). Angle nozzles forward 45 degrees for better coverage. Low drift nozzles may require higher pressures for proper performance.

Aerial: Minimum 43 PSI (300 kPa).

For either ground or aerial, use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops or weeds that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury or reduced weed control may result. Under drought conditions DO NOT apply to spring or durum wheat if the time from seeding to spraying exceeds 35 days or if temperatures will be 3°C or lower within 3 days of application (before or after).

Tank Mixes:

Add ammonium sulphate to the tank first then *Varro* then the tank mix partner.

Herbicides:

Wheat (including Spring, durum, winter)

Infinity (0.33 L/acre)

Thumper (0.4 L/acre)

Spring Wheat (including durum)

2,4-D ester (0.23 L/acre – 600 g/L form)

Buctril M (0.4 L/acre)

MCPA ester (0.23 L/acre – 600 g/L form)

Spring Wheat (NOT including durum)

Curtail M (0.61 L/acre)

Refine SG (12 g/acre)

Refine SG + 2,4-D ester (rates above)

Refine SG + MCPA ester (rates above)

Fungicides: None registered.

Insecticides: None registered.

Fertilizers: None registered.

Note: The above mixes are those listed on the *Varro* label only. To check for other possible mixes see the blue fold out chart in the back inside cover of the guide.

Bayer CropScience also supports the following mixes that are not on the *Varro* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Attain XC*, *Barricade II*, *Momentum*, *OcTTain*, *Prestige XC*, *Retain SG*, *Stellar*

Fungicides: *Tilt*.

All tank-mixes are applicable to spring wheat and durum. When tank-mixing *Varro* with *Prestige XC* or *Curtail M* in spring wheat always add ammonium sulphate.

Adding ingredients in the correct order is critical for optimum performance. Check labels of products to be mixed for directions. General guidelines can be found on page 14.

Restrictions:

Rainfall: Rain within 1 hour of application may reduce control.

Re-Entry: DO NOT enter treated field for 12 hours.

Grazing: Must not be grazed within 7 days or cut for live-stock feed within 30 days of treatment.

Preharvest Interval: DO NOT harvest grain or straw within 60 days of application for spring and durum wheat or within 72 days of application to winter wheat.

Re-cropping: Alfalfa, barley, canaryseed, canola, chickpea, dry bean, field corn, flax, lentil, mustard, oat, pea, soybean, sunflower, timothy, and wheat (durum, spring) may be seeded the year following application.

Aerial Application: May be applied by air.

Storage: Store in a cool, dry place. Keep from freezing. Shake well before using.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	0	1
Fixed wing aircraft	1	0	30
Helicopter	1	0	30

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general section on tank cleaning on page 15. If mixing with other pesticides, combine this method with the method indicated for the tank mix partner.

Hazard Rating:



Warning – Eye and Skin Irritant.

For an explanation of the symbols used here see page 11.

Velocity m3

Herbicide Group

2 - thien carbazone

6 - bromoxynil

27 - pyrasulfotole

(Refer to page 38)

Company:

Bayer CropScience (PCP# 29584)

Formulation:

5g/L thien carbazone-methyl, 31.3g/L pyrasulfotole and 175 g/L bromoxynil formulated as a suspension concentrate

Container size - 8.1 L and 129.6 L.

Crops and Staging*:

Spring wheat (including durum):

1 to 6 main stem leaf stage to a maximum of 3 tillers, and before the first node can be felt in the stem. DO NOT apply beyond 35 days of emergence.

Winter wheat:

Spring or fall from 1 to 6 leaf stage and before the first node can be felt in the stem.

* DO NOT apply after the first node as crop injury may occur.

Weeds and Staging:

Grass weeds controlled from 1 to 6 main stem leaves and prior to the emergence of the 3rd tiller:

Barnyard grass	Persian darnel†
Foxtail (green and yellow†)	Wild oat
Canaryseed	Japanese brome†

Broadleaf weeds controlled at the 1 to 6 leaf stage unless otherwise indicated:

Canada thistle (up to 30 cm)†	Ragweed, Common
Common chickweed	Round-leaved mallow
Cleavers (1 to 6 whorls)*	Russian thistle (up to 10 cm)
Dandelion (up to 25 cm diameter)†	Shepherd's-purse
Flixweed (up to 10 cm)	Sow-thistle
Hemp-nettle	(annual, perennial†)
Kochia (up to 10 cm)	Stinkweed
Lamb's-quarters	Volunteer canola
Pale smartweed	(all varieties)
Pigweed, redroot	Wild buckwheat
	Wild mustard

† Suppression only.

* Group 2 resistant cleavers from the 4 to 6 leaf stage add ammonium sulphate as per the Rates: section below.

Rates:

0.405 L per acre

(One 8.1 L container treats 20 acres, 129.6 L drum will treat 320 acres)

To control Group 2 resistant cleavers from the 4 to 6 leaf stage or when applying to high weed populations, advanced weed staging, in cool conditions, or when tank-mixing MCPA or 2,4-D, add ammonium sulphate at 202 g/acre (99% dry) or 0.4 L/acre (49% solution) at 4 to 6 whorls. If using an ammonium sulphate product with a different concentration, adjust the rate accordingly.

DO NOT apply Velocity m3 or other products containing thien carbazone, pyrasulfotole or bromoxynil more than once in the same year.

Application Information:

Water Volume:

Ground: 20 to 40 L per acre. Use higher water volumes for dense canopies.

Aerial: Minimum 11.4 L per acre.

Nozzles and Pressure:

Ground: For conventional flat fan nozzles use a pressure of 30 to 50 PSI (207 to 345 kPa). Angle nozzles forward 45 degrees for better coverage. Low drift nozzles may require higher pressures for proper performance.

Aerial: Minimum 43 PSI (300 kPa).

For either ground or aerial, use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

DO NOT apply to crops or weeds that are stressed (frost, low fertility, drought or flooding, disease or insect damage) as crop injury or reduced weed control may result.

DO NOT apply to spring or durum wheat under conditions where the time from seeding to spraying exceeds 35 days or if temperatures will be 3°C or lower within 3 days of application (before or after).

Tank Mixes:

None registered.

Bayer also supports the following mixes that are not on the *Velocity m3* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: 2,4-D Ester* (56-112 g ai/acre), Lontrel, MCPA Ester* (56-112 g ai/acre).

*When adding 2,4-D Ester or MCPA Ester to *Velocity m3* tank-mix, ammonium sulphate must be added as per the Rates: section above.

Fungicides: Tilt

Insecticides: Decis, Sevin XLR.

Restrictions:

Rainfall: Within 1 hour may reduce control.

Re-Entry: DO NOT enter treated field for 24 hours.

Preharvest: DO NOT harvest grain or straw within 60 days of application to spring and durum wheat or within 72 days of application to winter wheat.

Grazing: Must not be cut for livestock feed within 30 days or grazed by livestock within 25 days of treating the crop.

Re-cropping: Alfalfa, barley, canaryseed, canola, field corn (Manitoba only), flax, soybean (Manitoba only), tame oat, and wheat (durum, spring) may be seeded the year following application. Field pea may be grown the following year in all black, grey-wooded and dark brown soil zones. DO NOT plant field pea the season following *Velocity m3* use in the brown soil zone where organic matter content is below 2.5 % and where soil pH is above 7.5. Lentil may be seeded the second season after application.

Aerial Application: May be applied by air.

Storage: Store in a cool, dry place. Keep from freezing. This product is combustible. DO NOT store near heat or open flame.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	1	5
Fixed wing airplane	10	1	375
Helicopter	10	1	225

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to 'Method A' in the general tank cleaning section on page 15 to 16.

Hazard Rating:

Velocity m3 All-In-One:



Warning – Poison



Danger – Corrosive to eyes.

Skin Irritant

Potential Skin Sensitizer

For an explanation of the symbols used here see page 11.

Velpar DF

Herbicide Group

5 - hexazinone

(Refer to page 38)

Company:

E. I. duPont Canada (PCP#25225)

Formulation:

75% hexazinone formulated as a water dispersible granule
Container size - 2 kg.

Crops and Staging:

Established alfalfa for forage and seed. Apply in late fall prior to freeze-up when alfalfa is dormant or in early spring before alfalfa growth resumes. Apply only on alfalfa that has been established for 18 months or longer. If burning or irrigation is to be carried out, do not apply until these operations have been completed.

NOTE: DO NOT apply to frozen ground. DO NOT apply to soils with less than 1% organic matter. DO NOT apply to gravelly or rocky soils, exposed subsoils or sand. Crop injury may occur in fields where alfalfa root growth has been restricted by hard pans or other physical barriers to root growth.

Weeds, Rates and Staging:

Application stage is dictated by the crop above.

Apply a minimum of 0.272 kg per acre to control:

Dandelion

Quackgrass

Sow-thistle

Apply 0.544 kg per acre to control:

The weeds above plus:

Narrow-leaved hawk's-beard Scentless chamomile

Use the lower rate on medium-textured soils with low organic matter.

Application Information:

Water Volume: 81 L per acre.

Nozzles and Pressure: 30 to 40 psi (200 to 275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage with ASABE medium droplets.

How it Works:

Refer to Table 2 on page 40.

Effects of Growing Conditions:

Adequate soil moisture is required for activation of the product.

Tank Mixes:

None registered.

Restrictions:

Rainfall: Rainfall is beneficial for activation of the product.

Re-Entry: DO NOT re-enter treated fields for 48 hours.

Grazing: Leave 30 days between application and grazing harvesting for feed (hay or greenfeed).

Re-cropping: Leave 2 years of between treating alfalfa and the seeding of a crop. A field bioassay is required after 2 years to determine which crops are safe to grow.

Aerial Application: DO NOT apply by air.

Storage: May be frozen.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	1	1	5

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Handheld or backpack sprayers do not require a buffer zone.

Tank Cleaning:

No specific cleaning procedures are indicated on the label. Based on products with similar chemistry, 'Method B' found in the general tank cleaning section on page 15 to 16 or a commercial spray tank cleaning product, may provide adequate cleaning. Contact the manufacturer for more information.

Hazard Rating:



Danger - Corrosive to eyes.



Caution - Poison



Caution - Skin Irritant

For an explanation of the symbols used here see page 11.

Viper ADV

Herbicide Group

2 - imazamox

6 - bentazon

(Refer to page 38)

Company:

BASF Canada (PCP#30626)

Formulation:

20 g/L imazamox and 429 g/L bentazon formulated as a solution.

Container size - 2 x 8.1 L.

Requires the addition of:

BASF 28% UAN (28-0-0) is required, but sold separately.

Container size - 2 x (2 x 8 Liters); 128 L drums.

Crops and Staging:

Field pea and soybean: 3 to 6 above-ground nodes (3 to 6 true leaves).

Dry bean (black, cranberry, great northern, navy, pinto, pink, red Mexican): *Viper ADV* plus additional Basagran Forte (see tank mix section) from the fully expanded first trifoliate leaf to the second trifoliate fully expanded.

Even though *Viper ADV* is registered for all the dry bean types above, tolerance may vary between varieties (esp. navy). Test new varieties on a small area for tolerance before widespread use.

Note: Applications under hot, humid conditions may result in temporary leaf yellowing, leaf flecking, bronzing or burning. The crop usually outgrows this condition within 10 days and new tissues will not be affected.

Weeds and Staging:

Grasses - 1 to 4 main stem leaves or until early tillering.

Barnyard grass
Green foxtail
Japanese brome*
Persian dandel
Volunteer barley
Volunteer canaryseed

Tame oat
Volunteer wheat (including durum, not CLEARFIELD varieties)
Wild oat
Yellow foxtail

Broadleaf Weeds - cotyledon to 4 leaf stage.

Cleavers* †	Shepherd's-purse
Cow cockle	Stinkweed
Green smartweed	Sow-thistle (spiny annual)††*
Kochia*†	Stork's-bill††*
Lamb's-quarters	Volunteer canola (including CLEARFIELD varieties)
Pigweed (prostrate††*, redroot)	Volunteer lentils (including CLEARFIELD lentils)
Pigweed (prostrate††*, redroot)	Wild buckwheat*
Round leaf mallow*	Wild mustard†
Russian Thistle	

* Suppression only.

† Including Group 2 resistant biotypes.

†† *Viper ADV* + Basagran Forte in dry beans only.

Rates:

400 ml per acre

(One case of *Viper ADV* treats 40 acres)

Add 28 % **BASF UAN** (sold separately) at 0.81 L per acre.

Failure to include UAN will result in significantly reduced product performance.

DO NOT use any other adjuvants as injury may result.

DO NOT apply *Viper ADV* more than once or follow *Viper ADV* with any related products (*Basagran*, *Odyssey*, *Solo*) in the same year.

DO NOT apply to any crop other than those registered as severe injury will result.

Refer to the product label for complete mixing instructions for this product.

A general guide to mixing can be found on page 13.

Application Information:

Water Volume: 40 L per acre. High water volumes are required for adequate coverage, particularly when weed densities are high or weed staging is large.

Nozzles and Pressure: Use 40 psi (275 kPa) when using conventional flat fan nozzles. Low drift nozzles may require higher pressures for proper performance. Use a combination of nozzles and pressure designed to deliver thorough, even coverage a of ASABE medium droplets. Use 50 mesh (or coarser) filter screens.

Effects of Growing Conditions:

DO NOT spray if temperatures of +5°C or less are forecast within 3 days of application. Under cool or dry conditions, control of some weeds may be severely reduced. DO NOT apply to crops stressed from hail damage, flooding, drought, hot, humid weather, widely fluctuating temperatures, prolonged cold or injury from previous herbicides, as crop injury may result.

Tank Mixes:

Dry bean (types above):

Basagran Forte (145 mL/acre) plus UAN as above.

Restrictions:

Rainfall: Rain within 6 hours may reduce control.

Re-Entry: DO NOT enter treated fields for at least 12 hours.

Grazing: DO NOT graze or cut for feed.

Preharvest Interval: DO NOT apply within 60 days of harvest.

Re-cropping: Winter wheat may be seeded 3 months after application. Barley, canaryseed, canola, chickpea, field corn, field pea, flax, lentil, oat, sunflower, and spring wheat (including durum) may be seeded the first season after application and tame mustard (condiment types only) the second season after application. The company recommends that a field bioassay (a test strip grown to maturity) be conducted the year before growing any crops other than those listed above.

Contact manufacturer for additional information on recropping intervals.

Aerial Application: DO NOT apply by air.

Storage: DO NOT freeze. Store in a cool, dry place above 5°C.

Buffer Zones: Avoid spraying in situations where drift may occur. Leave at least 11 metres between the outside edge of the sprayed area and sensitive non-target areas such as shelterbelts, hedgerows, wetlands, woodlots, vegetated ditch banks, ponds, streams, and sloughs.

Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

Tank Cleaning:

Refer to 'Method B' in the general tank cleaning section on page 15.

Hazard Rating:



Warning – Poison



Danger – Corrosive to eyes (*Viper B* only).



Warning – Contains the allergen soy.



Warning – Eye and Skin Irritant.

For an explanation of the symbols used here see page 11.

Plant Disease Control

Integrated Plant Disease Management

Cereal Leaf Diseases

Cereal leaf diseases affect both the yield and quality of cereals. The following management practices are recommended for effective control of leaf diseases in all cereal crops.

Scouting: Scout fields prior to, during, and following flag leaf emergence to check for disease levels. Flag and upper leaves are responsible for 50 per cent or more of grain-fill.

Crop Rotation: Rotate crop types [e.g. cereal (wheat)/oilseed/cereal(barley)/pulse] to reduce the build-up of disease inoculum in crop residue. If at all possible, do not seed the same crop in back to back years. When a short rotation is absolutely necessary, seed in the second year a variety that is more resistant to an anticipated disease problem.

Resistant Varieties: Provincial crop/seed guides provide a comprehensive listing of the performance of adapted varieties, including their resistance status to specific diseases.

Foliar Fungicides: Foliar fungicides, applied at the proper time in accordance to manufacturers' Pesticide Product labels, can control cereal leaf diseases and help to attain target yields. The greatest benefit occurs when disease pressure is high or with varieties that have poor resistance.

Spraying Practices: Foliar fungicides should be applied preventively, before disease is well-established in a crop and already causing crop loss. Good spray coverage with minimal drift is essential. Ideally, the best time to spray is when the wind is light, humidity is above 60 per cent and air temperature is between 10 and 25°C.

Ergot of Cereals

Ergot is a fungal disease that affects most cereals and grasses in Canada. Ergot bodies contain toxic alkaloids; they should never be consumed by humans or fed to animals. Ergot is a particularly damaging disease of rye, and has also been observed sporadically over the years on wheat, durum, and triticale in the prairie provinces, with elevated levels reported in 1999, 2005, 2008, and 2011. After an ergot outbreak, crop residue and soil become contaminated with a higher load of ergot bodies, placing nearby grasses and cereal crops at greater risk of infection in the following seasons. This risk increases further when cool, moist weather conditions promote ergot spore production and/or when cereals experience an extended period of flowering or an induction of floret sterility due to any of a variety of agronomic or environmental factors. Once ergot is present, little can be done to control the disease in the field, so prevention is important. Planting seed contaminated with ergot bodies can potentially spread disease to previously clean fields and there are no seed treatments registered; therefore only

clean, healthy seed should be used. During the field season, nearby grasses may be mowed to remove additional hosts. Prior to harvest, fields should be scouted to determine where ergot has developed, such as headlands, and those areas should be harvested separately. Viability of ergot bodies decreases after one to two years.

Fusarium Head Blight of Cereals

Fusarium head blight (FHB) causes a reduction in yield as a result of floret sterility and the loss of light weight cereal kernels during combining. More important is the effect on grain quality and food safety due to production of mycotoxins, including deoxynivalenol (DON) and vomitoxins. In Manitoba, FHB occurs throughout all crop regions and will damage wheat crops whenever environmental conditions favour the disease. In Saskatchewan, FHB has been established in eastern regions for several years, but occurs across the province, particularly in wet years.

Field Management of FHB: Weather is by far the greatest factor in development of FHB. The disease is most likely to develop when the plants are flowering, temperatures range from 15 to 30°C and high moisture is continuous for 48 to 60 hours. If conditions remain warm and moist, the pathogen can continue to sporulate and spread to other kernels or heads. Under these optimum conditions, crop management has little impact on FHB outbreaks. Production practices, which lead to reduced tillering and shortened flowering duration, could reduce the risk period of FHB infection.

Crop Rotation and Crop Selection: A break of at least one year – preferably two years – is advised between cereal, grass and corn production. In fields of wheat on wheat stubble, the incidence of FHB was about one and a half times higher than in fields of wheat planted into pulse crop residue. Regardless of the rotation, producers should consider planting cereals that are less susceptible to FHB. Results from previous years show that durum and soft white wheat varieties are more susceptible than hard red spring wheat varieties. Barley is more resistant than wheat, and oat is more resistant than either wheat or barley. Refer to provincial seed guides for FHB disease ratings for each variety. Avoid planting the more susceptible types or varieties of wheat in high risk areas such as the southern part of the Red River Valley and southeastern Saskatchewan. Planting two or more varieties of wheat with differing flowering times or varying planting dates will help reduce the risk of infection. Susceptible crops should not be planted on infected corn stubble. Corn trash is slower to decompose than cereal trash, and acts as a source of inoculum for a much longer time period.

Late Blight of Potatoes

One of the major threats to Manitoba and Saskatchewan's potato industry is the fungal disease late blight. At present, there is no fungicide registered for use on potatoes that is capable of eradicating the fungus from infected plants. As a result, producers are forced to adopt preventive management to control this disease. One of the main components of this strategy is the application of fungicides at specified spray intervals. This interval varies with the type of fungicide used. Shortening or lengthening of this interval should be based on current weather conditions and the status of the disease in the crop.

In Manitoba, potato producers can make use of a weather-based late blight risk forecasting program. The purpose of this program is to predict when environmental conditions are most conducive to disease development and issue warnings based on those parameters.

Accurate weather monitoring and scouting techniques are very important for achieving the most effective use of fungicides. Combining precise weather forecasting with spray interval scheduling may lower input costs for the farmer and lead to a more productive, higher quality crop. These weather monitoring systems monitor key environmental variables, such as relative humidity, temperature, leaf wetness and precipitation.

The following practices are recommended for effective disease management:

Scouting: To effectively schedule preventative fungicide applications and eliminate unnecessary fungicide use, local weather forecasts should be used to identify conditions conducive to disease development. Scout fields regularly to identify diseases and pests that may be developing. Low areas in rolling or hilly fields and in wind-protected area near trees lines should be specially checked.

Crop Varieties: *(There are no known commercial resistant varieties currently available in Canada).* Where practical, the use of short season varieties may help reduce the period of use for fungicides.

Healthy Seed: Obtain seed from sources with effective disease management practices. The use of certified seed is legislated in Manitoba and Saskatchewan. Grade seed carefully while cutting and discard suspicious looking tubers and seed pieces.

Cull Clean-up: Avoid leaving tubers, including debris or slivers from seed cutting, in cull piles for any length of time. Follow a program of sanitation for storage facilities and equipment to eliminate sources of the disease. Dispose of cull piles in an approved manner so they do not serve as a source of disease inoculum for future infections. Dispose by burying, using a cover, spreading out on the field over winter, or feeding to livestock.

Sanitation: Follow a program of sanitation for storage facilities and equipment to eliminate sources of disease. Sanitation consists of cleaning and disinfecting all equipment, storage, and tools that contact potatoes from seeding

through harvest and storage. Since most disinfectants are inactivated by soil and plant debris, it is essential that equipment and storage is thoroughly cleaned with a pressure washer or steam cleaner with detergent before disinfectant is applied. Treated surfaces should remain wet for at least 20 minutes for the disinfectant to destroy disease organisms.

Cultural Practices and Rotation: Use proper cultural practices including a one in four year potato crop rotation; proper hilling to reduce disease and greening in tubers; manage irrigation to avoid an excess or deficit of soil moisture; schedule irrigation throughout the day so it is not extending the natural dew period and prolonging leaf wetness; if late blight is discovered destroy hot spots of infected fields; control weed hosts (especially nightshades) and remove and destroy volunteer potatoes. Use appropriate weed control practices in rotational crops to control those weeds that may be hosts of diseases in potatoes.

Foliar Fungicides: Preventative fungicide applications are most effective in controlling late blight. Follow product label guidelines for most efficient and safe use of products. Labels of newly registered products also provide information on resistance management. In this context – medium to high risk of resistance fungicides (e.g. Group 7 – boscalid and Group 11 – strobilurins) should be rotated or mixed with low risk fungicides (e.g. mancozeb (M3) and chlorothalonil (M5)).

Farm Visits: The following recommendations are provided to prevent the spread of potato diseases from field to field or between farms. All people serving the potato industry should use these sanitary practices.

1. Contact the grower for permission to enter fields and other facilities on the farm.
2. Keep your vehicle clean and whenever possible, avoid driving your vehicle into fields or potato handling areas.
3. Carry a boot brush and a supply of disinfectant in your vehicle at all times. Quaternary ammonia (General Storage Disinfectant) is recommended as it is also registered for bacterial ring rot disinfection.
4. Wear coveralls or other protective outerwear that can be discarded or disinfected regularly.
5. Clean, washable, footwear is recommended and rubber boots are preferred.
6. Clean, wash, and disinfect your boots thoroughly on arrival at each field / farm / storage shed and before leaving.
7. Remove dirty outerwear, including boots before entering your vehicle.
8. Any tools to be used during the farm call (potato forks, shovels, soil probes, knives, etc) should be cleaned and disinfected before and after use.
9. Maintain a detailed logbook of field / farm / storage shed visits.

Canola Diseases

Sclerotinia stem rot has been one of the most prominent diseases affecting canola in Manitoba and Saskatchewan for the past 25 years. An important factor for disease development is environmental conditions. The disease is much more widespread and severe during wet years. Fungicide applications are an important element in controlling the development and spread of sclerotinia. Fungicide spray decisions are based on soil moisture, weather conditions, crop stage and density, and disease history. The sclerotinia resting bodies (sclerotia) require moist soil conditions for up to 10 days for germination to occur and the spore-bearing structures (apothecia) to form. Usually these conditions do not occur until the crop canopy closes. The spores released from the apothecia utilize the canola petals as a food source and fall into the canola canopy where they infect plants. Lesions form up and down the stem, wilting leaves and eventually killing the plant. Fungicide should be applied between the 20 to 50 per cent flower stages to protect the petals from being colonized by the spores.

Blackleg caused by *Leptosphaeria maculans* affects canola and most crucifer field and vegetable crops. After many years of low incidences, due to resistant canola varieties grown in the prairie provinces, the disease is gaining importance again. High frequency of canola in crop rotations, accompanied by changes in the pathogen populations, has led to higher incidences and severities in some fields. For an effective control, a 4 year crop rotation is highly recommended.

Clubroot is a soil-borne disease caused by a microbe, *Plasmodiophora brassicae*. Clubroot affects the roots of cruciferous field crops such as canola, mustard, and camelina, as well as cruciferous vegetables and weeds. Clubroot has become a significant problem for canola growers in some areas of Alberta and the pathogen has been detected in Saskatchewan and reported once in Manitoba. Clubroot is a regulated pest in Saskatchewan under *The Pest Control Act*. Currently there is no provincial legislation that regulates clubroot in Manitoba.

Invasion of the interior of the host roots alters hormone balance and leads to increased cell division and growth, resulting in clubroot galls. These deformed roots have a reduced ability to absorb water and nutrients leading to stunting, wilting, yellowing, premature ripening and shrivelling of seeds. The cause of these above-ground symptoms can be confirmed by digging up suspect plants to check roots for gall formation. Clubroot affects canola yield and quality to a similar degree as other diseases affecting water and nutrient uptake, and its impact depends on soil conditions and the growth stage of the crop when infection occurs. Spore germination, infection and disease development are favoured by warm soils, high soil moisture and low soil pH; however, the disease can still occur under conditions outside of the optimum parameters. Infected roots will eventually disintegrate, releasing resting spores

into the soil, which may then be transported by wind, water erosion, animals/manure, shoes/clothing, vehicles/tires or earth tag on agricultural or industrial field equipment. Resting spore numbers will decline over time when non-host crops are grown, but a small proportion can survive in soil for up to 20 years. Clubroot is primarily a soil-borne disease; it does not infect seed but it may be found in soil attached to seed or other plant parts. There are currently no seed treatments or foliar fungicides registered for control of clubroot on canola. The following best practices are recommended for prevention and management of clubroot:

1. Plant susceptible crops, including resistant varieties, no more than once every four years. Although crop rotation will not prevent the introduction of clubroot to fields that are free of the pathogen, it will restrict clubroot development by limiting the increase of clubroot resting spores and preventing the increase of clubroot inoculum, as well as help alleviate the impact of other plant pathogens.
2. Scout crops regularly and carefully.
 - Identify suspicious above-ground symptoms including wilting, stunting, yellowing and premature ripening of canola or other susceptible crops.
 - Field entrances and approaches are likely to be contaminated with clubroot spores first. Therefore, symptoms will often appear there first.
 - Confirm cause of above ground symptoms by checking the roots for galls.
 - Send sample of symptomatic plants into a commercial lab for confirmation of diagnosis.
3. Practice good sanitation by restricting movement of potentially contaminated soil to non-contaminated regions.
 - For Saskatchewan and Manitoba producers, this means restricting entry into their fields of vehicles, field machinery or oil rig equipment with earth tag from infested regions unless it has been properly sanitized. Ask questions about where the equipment is from and what sanitation measures have been used before the equipment left the infested area, dealer or auction site.
 - Cleaning steps may include: removal of crop debris and soil, washing of equipment with a power washer using hot water or steam and misting with disinfectant (1-2 per cent bleach solution), followed by an additional rinse with water.
 - Other agricultural products, which could carry soil, should be carefully checked for excess soil and if possible be from clubroot free areas.

For more information on clubroot, visit www.clubroot.ca, www.agriculture.gov.sk.ca, or www.gov.mb.ca/agriculture.

Pulse Crop Diseases

There are a variety of pulse crops produced in Manitoba and Saskatchewan including field pea, field bean, lentil, chickpea and soybean. Pulse crops are adapted to different regions and will require unique agronomic and disease management practices. Some diseases will attack all pulse crops, e.g. sclerotinia (white mould) and seedling/root rots caused by *Pythium*, *Rhizoctonia*, *Fusarium* and *Botrytis* species. Some diseases may occur on more than one type of pulse crop, but the pathogen species infecting each is often specific to that crop. This is the case for the ascochyta blights, powdery mildews and anthracnose. It is important to source information on pulse disease control from grower organizations such as the Saskatchewan Pulse Growers (www.saskpulse.com), Manitoba Pulse Growers (www.manitobapulse.ca), provincial specialists, and field diagnostic guides. Most foliar diseases are favoured by warm, moist conditions and lush crop canopies, but root rots and powdery mildew can be present in dry years as well. In general, pulse disease management will need to include the following practices:

Use of clean seed and seed treatments: Plant certified seed or seed that has been tested at an accredited lab and known to have high germination and zero or acceptable levels of

seed-borne disease. Seed treatments will help protect seed and seedlings from low levels of seedborne and soilborne pathogens

Crop Rotation: It is important to keep at least three years between the same type of pulse crop to allow for the breakdown of crop residue on which disease pathogens survive. Since there are diseases that affect more than one type of pulse crop, it is still important to maintain at least two years between different pulse crops.

Crop Varieties with Disease Resistance: Refer to provincial seed guides for varieties adapted to your region. When available, choose varieties with disease resistance.

Scouting and Foliar Fungicide Application: Begin crop scouting at the vegetative stages for aggressive diseases such as ascochyta blight in chickpea. Scout for other foliar diseases at early bloom, e.g. ascochyta blight and anthracnose in lentil. It is too late to apply fungicide to control sclerotinia (white mould) once symptoms are observed, and/or the canopy has closed, so forecasting to determine risk is necessary.

Use foliar fungicides only when disease risk and potential loss are significant. Rotate fungicides or use tank mixes from different fungicide groups to prevent development of resistant pathogen populations.

Effects of Weather

Do not apply foliar fungicides during periods of dead calm or when winds are gusty. Avoid application immediately after a rainfall and delay spraying if rainfall is imminent. Contact fungicides are always more sensitive to wash-off by rainfall than systemic fungicides, because their mode of action relies on drying on the leaf surface. Failure of a

contact fungicide to dry on the leaf surface may result in a loss in efficacy. Systemic fungicides are less sensitive than contact fungicides, but still need sufficient drying time and fully absorbed by plants prior to rainfall. Consult the label or product manufacturers for rainfall period for individual products.

Pathogen Resistance (Insensitivity) Management

Any fungal pathogen population may contain strains naturally insensitive to a fungicide and other fungicides within the same Group. A gradual or total loss of disease control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist.

To delay fungicide resistance/insensitivity:

- Where possible, rotate the use of a fungicide, (and others within the same Group) with different Groups that control the same pathogens.
- Where possible, tank mix fungicides with a high risk of developing insensitivity with other fungicides from a different Group.
- DO NOT apply more than the maximum number of applications listed on the label. Avoid consecutive

sprays of a fungicide, or other fungicides in the same Group, in a season.

- Fungicide use should be based on an integrated pest management (IPM) program that includes scouting and accurate recording related to pesticide use and crop rotation. An IPM program also considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for signs of fungicide insensitivity. If disease continues to progress after treatment with a product, DO NOT increase the use rate. Discontinue use of the product and switch to another fungicide with a different target site of action.
- Contact your local regional crops specialist or certified crop advisor for any additional pesticide management and/or IPM recommendations for specific crops and disease problems in your area.

Fungicide Modes of Action

Why are fungicides needed?

- Control of disease during crop establishment.
- Increase productivity of crop (photosynthesis) and/or reduce blemishes.
 - Maintain yield and/or market value.
- Improve storage life and quality of harvested plants / grain / produce.
 - Prevent spoilage and/or production of mycotoxins.

How do fungicides work?

There are several ways to define 'mode of action':

Timing:

- Preventative: fungicide must be present on plant surface before the pathogen and repeated applications are required to protect new growth.
- Curative: pathogen may already be present (post-infection, pre-symptom kick-back activity).
- Eradicant: (post-symptomatic activity).
- Inhibitive: prevents spore germination or sporulation.

Placement:

- Contact (AKA protectant): immobile – must come in direct contact with the pathogen.
- Systemic (AKA penetrant): mobile – can move within plant.

Movement:

- Intra-plant Movement: within crop via vapour phase or redistribution by rain.
- Passive Absorption – by diffusion.
- Apoplastic Movement: xylem-mobile; move within free space and cell walls, upward through the transpiration stream (with water).

- Symplastic Movement: phloem-mobile (common characteristic of herbicides and insecticides but very few fungicides).

Spectrum:

- General, Non-specific, or Broad Spectrum: fungicide affects pathogen in multiple ways.
- Specific or Narrow Spectrum: fungicide targets a specific metabolic site in pathogen or against critical enzyme or protein. Genetic changes or naturally insensitive fungi have a greater chance to overcome the fungicidal effect (resistance/insensitivity).

Composition:

- Inorganic Fungicides: sulfur or metal ions such as copper.
- Organic Fungicides: contain carbon atoms.
- Biopesticides: suppressing pest populations using naturally occurring organisms or natural products derived from plants.

Biochemistry:

- Primary basis to classify fungicides, developed by Fungicide Resistance Action Committee (FRAC) using their general Mode of Action on fungi and their chemistry.
 - All fungicides within a group share a common mode of action and resistance mechanism.
 - Fungicides within a group may have different chemical structures.
 - Resistance management strategies required wherever resistance is known or there is a risk of resistance development
 - See Table 1.

Table 1. Fungicide Groups Based on Biochemical Mode of Action (FRAC)

Mode of Action Target	Chemical Group & Chemical Name	Resistance Risk	Foliar Fungicide Products Registered in Saskatchewan/Manitoba	Seed Treatment Registered in Saskatchewan/Manitoba	Products
A. Nucleic Acid Synthesis	4. Phenyl Amides	High	Ridomil Gold/Bravo [®] , Ridomil Gold SL/Bravo [®] , Ridomil Gold 480EC, Ridomil Gold 480SL	Acceleron Seed Treatment Package [®] , Allegiance II, Apron Advance [®] , Apron Mxxx RIC [®] , Apron Mxxx RTA [®] , Belmont 2.7FS, Cruiser Mxxx Beans [®] , Cruiser Mxxx Cereals [®] , Cruiser Mxxx Pulses [®] , Cruiser Mxxx Vibrance Beans [®] , Cruiser Mxxx Vibrance Cereals [®] , Dividend XL RTA [®] , EverGol Energy [®] , Helix Vibrance [®] , Helix Xtra [®] , Insure Cereal [®] , Maxim Quattro [®] , Prosper Evergol [®] , Prosper FX [®] , Raxil MID [®] , Raxil PRO [®] , Raxil PRO Shield [®] , Raxil WW, Trilex AL [®] , Trilex EverGol [®] , Vibrance Mxxx RTA/RIC [®] , Vibrance XL [®]	
B. Mitosis & Cell Division	1. Methyl Benzimidazole Carbamates	High	Senator 70WP	Apron Advance [®] , Crown [®] , DCI [®] , Maxim Quattro [®] , Mertect SC, Senator PSPT	
	22. Benzamide	Low to Medium	Gate 75DF [®] , Torrent	None	
C. Respiration	7. Carboxamides	Medium	Cantus WDG Fungicide, Fontelis, Headline Duo [®] , Lance, WDG Fungicide, Luna Tranquility [®] , Priaxor DS [®] , Propulse [®] , Vertisan	Crown [®] , Cruiser Mxxx Vibrance Beans [®] , Cruiser Mxxx Vibrance Cereals [®] , EverGol Energy [®] , Gaucho CS FL [®] , Helix Vibrance [®] , Prosper Evergol [®] , Prosper FX [®] , Rancona RS, Trilex EverGol [®] , Vibrance Mxxx RTA/RIC [®] , Vibrance 500S, Vibrance XL [®] , Vitaflo 220 [®] , Vitaflo 280 [®] , Vitaflo Fungicide [®] , Vitaflo SP Fungicide [®]	
	11 Strobilurins	High	Acapela, Blanket [®] , Cabrio Plus [®] , Headline EC, Headline Duo [®] , MANA Azoxystrobin, Quadris, Quilt [®] , Priaxor DS [®] , Quadris Top [®] , Reason 500SC, Stratego 250EC [®] , Tanos 50DF [®] , Twinline [®]	Insure Cereal [®] , Maxim Quattro [®] , Prosper Evergol [®] , Prosper FX [®] , Trilex AL [®] , Trilex EverGol [®]	
	21. Cyanoimidazole	Medium to High	Ranman 400SC, Torrent 400SC	None	
	29. 2,6-Dinitroanilines	Low	Allegro 500F	None	
	45. Triazolopyrimidyl-amine	Medium to High	Zampro [®]	None	
	9. Anilinopyrimidine	Medium	Astound [®] , Luna Tranquility [®] , Scala SC	None	
D. Amino Acid & Protein Synthesis					
E. Signal Transduction	2. Dicarboximides	Medium to High	Rovral Flo, Rovral RX, Overall 240SC	None	

Continued...

Table 1. Fungicide Groups Based on Biochemical Mode of Action (FRAC) *continued*

Mode of Action Target	Chemical Group & Chemical Name	Resistance Risk	Foliar Fungicide Products Registered in Saskatchewan/Manitoba	Seed Treatment Registered in Saskatchewan/Manitoba
E. Signal Transduction <i>Continued</i>	12. Phenylpyrroles	Low to Medium	<i>Astound*</i>	<i>Acceleron Seed Treatment Package*, Apron Advance*, Apron Maxx RFC*, Apron Maxx RTA*, Cruiser Maxx Beans*, Cruiser Maxx D Potatoes*, Cruiser Maxx Pulses*, Cruiser Maxx Vibrance Beans*, Helix Vibrance*, Helix XTra*, Maxim D*, Maxim Liquid PSP, Maxim MZ PSP*, Maxim PSP, Maxim Quattro*, Proseed, Vibrance Maxx RTA/RFC*</i>
F. Lipid / Membrane Synthesis & Cell Wall Degradation	28. Carbamates	Low to Medium	<i>Tattoo C*</i>	None
	40. Carboxylic Acid Amides (CAA)	Low to Medium	<i>Acrobat 50WP, Revus, Zampro*</i>	None
	44. <i>Bacillus</i> strain QST 713	Low	<i>Serenade CPB, Serenade Max</i>	None
G. Sterol Biosynthesis	3. Demethylation Inhibitors	Medium	<i>Blanket*, Bumper 418EC, Caramba, Folicur 432F, Folicur 250EW, Fuse, Inspire, Palliser, Pivot 418EC, Proline 480SC, Propel, Propiconazole Fungicide, Propulse*, Prosaro 250EC, Quadris Top*, Quash, Quilt*, Stratego 250EC*, Tilt 250E, Twinline*</i>	<i>Acceleron Seed Treatment Package*, Armour, Armour RTU, Charter RTU, Cruiser Maxx Cereals*, Cruiser Maxx D Potatoes*, Cruiser Maxx Vibrance Cereals*, Dividend XL RTA*, EverGol Energy*, Gemini*, Helix Vibrance*, Helix XTra*, Insure Cereal*, Maxim D*, Ramona Apex, Ramona RS, Raxil MD*, Raxil PRO*, Raxil PRO Shield*, Raxil WW*, Vibrance XL*</i>
U. Unknown	27. Cyanoacetamide-oximes	Low to Medium	<i>Curzate 60DF, Tanos 50DF*</i>	None
	33 Phosphonates	Low	<i>Phostrol</i>	<i>Confine</i>
	NC (Not classified) and diverse	Not known	<i>Contans WG, Regalia Maxx</i>	<i>Acceleron DX-910, Acceleron Seed Treatment Package*, Heads Up Plant Protectant, StorOx</i>
M. Multi-Site Contact Activity	M1. Inorganic copper	Low	<i>Copper products, Kocide 2000</i>	<i>Kocide 2000</i>
	M2. Inorganic sulphur		<i>Kumulus DF</i>	None
	M3. Dithiocarbamates		<i>Cabrio Plus*, Dithane Rainshield, Gavel 75DF*, Manzata Pro-Stick, Penncozeb 75DF, Polyram DF</i>	<i>DB-Red I, Gaucho CS FL*, Polyram 16D, Potato ST16, Solan MZ, Thiram 75 WP, Tuberscal, Vitaflo 220*, Vitaflo 280*, Vitaflo Fungicide*, Vitaflo SP Fungicide*</i>
	M4. Phthalimides		None	<i>Agrox B-2, Agrox CD, Agrox FL, DCT*, Gemini*, Maxim MZ PSP*</i>
	M5. Chloronitriles		<i>Bravo 500, Bravo Zn, Echo 720, Echo 90DF, Ridomil Gold/Bravo*, Ridomil Gold SL/Bravo*, Tattoo C*</i>	None

*Products contain more than one active ingredient and appear in more than one group.

Foliar Fungicide Tables

Table 2. Foliar Fungicides for Disease Control in Potatoes

FUNGICIDES	Page	Black Dot	Botrytis Gray Mould / Botrytis Vine Rot	Brown Leaf Spot	Early Blight	Late Blight	Late Blight Tuber Rot	Pythium Leak	Pink Rot	Rhizoctonia Canker, Black Scurf, Stolon Canker, and Stem Rot	Silver Scurf	White mould / Sclerotinia rot
Acrobat WP	321					1	1					
Allegro 500F	322			•		•						
Bravo 500	330		•		•	•						
Bravo Zn	330				•	•						
Cabrio Plus	327				•	•						
Cantus WDG	328				•							
Copper 53W	334				•	•						
Copper Spray	334				•	•						
Curzate 60 DF	336					1						
Dithane Rainshield	349				•	•						
Echo 90DF / Echo 720	330		•		•	•						
Gavel 75 DF	339				•	•						
Headline EC	341				•	•						
Inspire	344				•							
Kocide 2000	334				•	•						
Luna Tranquility	348	4		•	•							•
Manzate Pro-Stick	349				•	•						
Parasol WG	334				•	•						
Penncozeb 75 DF	349				•	•						
Phostrol	350					•	•	2,3				
Quadris	324	•			•	•				2	2	
Quadris Top	361	4		4	•							
Quash	362				•							
Ranman 400SC	337					•	•					
Reason 500SC	366				1	1						
Revus	368					•						
Revus Top	369				•	•						
Ridomil Gold/Bravo, Ridomil Gold SL/Bravo	369		•		•	•	•	3	3			
Ridomil Gold 480EC, Ridomil Gold 480SL	369								2			
Scala SC	371				1							
Serenade Max/CIPB	373				3							3
Tanos 50 DF	375				•	•						
Tattoo C	376					•						
Torrent 400 SC	337					•	•					
Vertisan	381		•		3					3		
Zampro	383					•	•					

Note: Before using any pesticide on potatoes, consult the list of Agricultural Pesticides Approved for Use, available from Simplot Canada and McCain Foods (Canada).

• Fungicide effective against the disease

1. Must not be used alone, only as a tank mix (consult individual labels)

2. In-furrow treatment (suppression only)

3. Suppression only (foliar application)

4. Suppression only

Table 3. Foliar Fungicides for Disease Control in Cereals

FUNGICIDES	Page	CROPS						DISEASES																	
		Wheat (W)	Barley (B)	Oat (O)	Rye (R)	Triticale (T)	Corn (C)	Fusarium Head Blight (wheat, barley, oat, rye)*	Fusarium and Gibberella Ear Rot** (corn)	Gray Leaf Spot (corn)	Northern Leaf Blight (corn)	Southern Leaf Blight (corn)	Eye Spot (corn)	Net Blotch (barley)	Powdery Mildew (wheat, barley, oat, rye, triticale)	Rusts					Scald (barley)	Septoria Glume Blotch (wheat)	Septoria Leaf Spot (wheat, barley, oat, rye, triticale)	Spot Blotch (wheat, barley)	Tan Spot (wheat, barley, rye, triticale)
																Common Rust (corn)	Crown Rust (oat)	Leaf Rust (wheat, barley, rye, triticale)	Stem Rust (wheat, barley, oat, triticale)	Stripe Rust (wheat, barley)					
Acapela	319	X	X	X	X	X				•				•				W,R,T					W,B,R,T		W
Blanket	326	X	X		X								•					W,B		•	•		W,B,R		W,B,R
Bravo 500	330	X					W															•	W		W
Bumper 418EC	355	X	X	X		X				•			•	W,B	•	•	W,B	W,B	W	•	•	W,B,O	B	W	
Caramba	329	X	X	X	X	X	•						•			•	•			•		W,O,T	•*	•	
Dithane Rainshield	349	X															W					W		W	
Echo 720	330	X					W														•	W		W	
Folicur 432F/ Folicur 250EW	377	X	X	X			W						•	W,B		•	W,B	W,B,O	•	•	•	W,B	B	W	
Fuse	377	X	X	X			W						•	W,B		•	W,B	W,B,O	•	•	•	W,B	B	W	
Headline EC	341	X	X	X	X	X			•				•	W,R	•	•	W,R		•	•		W	•	W	
Manzate Pro-Stick	349	X															W					W		W	
Palliser	377	X	X	X			W						•	W,B		•	W,B	W,B,O	•	•	•	W,B	B	W	
Penncozeb 75DF	349	X															W					W		W	
Pivot 418EC	355	X	X	X		X				•			•	W,B	•	•	W,B	W,B	W	•	•	W,B,O	B	W	
Proline 480SC	353	X	X	X		X	W,B	•	•	•		•	•		•	•	W			•	•	W	B	W	
Propel	355	X	X	X		X				•			•	W,B	•	•	W,B	W,B	W	•	•	W,B,O	B	W	
Nufarm Propiconazole Fungicide	355	X	X	X		X				•			•	W,B	•	•	W,B	W,B	W	•	•	W,B,O	B	W	
Prosaro 250EC	359	X	X				W,B						•	W,B			W,B	W,B	•	•	•	W,B	B	W	
Quadris	324					X										•									
Quilt	363	X	X	X		X			•	•	•	•	•		•	•	W,B		•	•		W,B,O		W	
Regalia Maxx	367	X																				W*			
Tilt 250E	355	X	X	X		X				•			•	W,B	•	•	W,B	W,B	W	•	•	W,B,O	B	W	
Twinline	380	X	X	X	X	X							•	W,R,T		•	W,R,T		•			W,T	•	W,T	
Vertisan	381	X	X	X	X	X	X		•*						•		W,R,T	•				(W,B,T)*			

* Suppression only

X Product registered for the crop

• Fungicide registered for disease control on all crops listed unless otherwise noted by the abbreviations listed: wheat (W), barley (B), oats (O), rye (R), triticale (T)

Table 4. Foliar Fungicides for Disease Control in Specialty and Other Crops

FUNGICIDES	Page	CROPS					DISEASES							
		Alfalfa (A)	Canary-seed (CS)	Coriander (CR)	Grasses grown for seed (G)	Timothy (TY)	Blossom Blight (A & CR)	Common Leaf Spot (A)	Spring Black Stem & Leaf Spot (A)	Sclerotinia (A)	Septoria Leaf Mottle** (CS)	Leaf Rust & Stem Rust (G)	Powdery Mildew (G)	Purple Eye Spot (TY)
Bumper 418EC	355		X							•				
Dithane Rainshield	349	X*						•	•					
Fontelis	338	X								•				
Headline EC	341	X*			X			•			•	G**		
Lance	346	X*					A	•	•	•				
Manzate Pro-Stick	349	X*						•	•					
Overall 240SC	345	X*								•				
Penncozeb 75DF	349	X*						•	•					
Pivot 418EC	355		X			X			•		•		•	
Propel	355		X			X					•		•	
Nufarm Propiconazole Fungicide	355		X								•			
Quadris	324			X			CR							
Rovral Flo / Rovral RX	345	X*								•				
Tilt 250E	355		X			X					•		•	

* Alfalfa grown for seed

** Suppression only

X Product registered for the crop

• Fungicide registered for disease control on all crops listed unless otherwise noted by the abbreviations listed: alfalfa (A), canaryseed (CS), coriander (CR), grasses (G), timothy (TY)

1. In-furrow treatment

Table 5. Foliar Fungicides for Disease Control in Oilseed Crops

FUNGICIDES	Page	CROPS					DISEASES									
		Canola (CA)	Flax (F) (and Crambe for Proline only)	Rapeseed, Oriental Mustard (M)	Soybean (S)	Sunflower (SF)	Alternaria Spot (canola, sunflower)	Anthrachnose (soybean)	Blackleg (canola)	Brown Spot (soybean)	Frogeye / Cercospora Leaf Spot (soybean)	Pasmo (flax)	Powdery Mildew (soybean)	Rust: Asian Soybean Rust (pulses, soybean)	Rust: <i>Puccinia helminthii</i> (sunflower)	Sclerotinia White Mould / Head Rot (oilseeds)
Acapela	319				X					•	•			•		S*
Allegro 500F	322				X											S
Astound	323	X														CA
Bumper 418EC	355	X			X				•		•					
Caramba	329				X									•		
Contans	332	X			X	X										(CA,S,SF)*
Folicur 432F/ Folicur 250W	377				X						•		•†	•		
Headline EC	341	X	X	X	X	X	CA		•		•	•		•	•	
Lance	346	X				X	•									CA,SF*
Overall 240SC	345	X					CA*									CA
Pivot 418EC	355	X			X				•		•					
Proline 480SC	353	X	X	X	X						•			•		CA,EM
Propel	355	X			X				•		•		•	•		
Nufarm Propiconazole Fungicide	355	X			X				•		•					
Quadris	324	X			X		CA		•		•		•	•		CA
Quash	362	X														CA
Quilt	363				X			•			•		•	•		
Rovral Flo/Rx	345	X					CA*									CA
Serenade Max/CPB	373	X			X					•*	•*					(CA,S)*
Stratego 250EC	374				X						•			•		
Tilt 250E	355	X			X				•		•		•	•		
Vertisan	381	X			X	X				•*	•*			•*	•*	CA, SF*

* Suppression only

X Product registered for the crop

† Folicur 432F is not registered for control of powdery mildew in soybeans

• Fungicide registered for disease control on all crops listed unless otherwise noted by the abbreviations listed: canola (CA), flax (F), rapeseed/oriental mustard (M), soybean (S), sunflower (SF)

Table 6. Foliar Fungicides for Disease Control in Pulse Crops

FUNGICIDES	Page	CROPS						DISEASES								
		Bean (B)	Chickpea (CP)	Fababean (FB)	Lentil (L)	Field Pea (P)	Anthracnose (bean, lentil)	Ascochyta Blight (bean, chickpea, lentil, pea)	Bacterial Blight (bean)	Botrytis Gray Mould (bean, chickpea, lentil, pea)	Downy Mildew (bean, pea)	Mycosphaerella Blight (pea, fababean)	Powdery Mildew (pulses)	Rust: <i>Uromyces</i> (bean)	Rust: Asian Soybean Rust (pulses)	Sclerotinia White Mould (pulses)
Acapela	319	X	X		X	X						P			B,CP,L,P	(B,CP,L,P)*
Allegro 500F	322	X														B
Bravo 500	330		X		X	X	L	CP, L				P				
Bumper 418EC	355	X												•		
Contans	332	X														B*
Copper 53W	334	X					B		•		B					
Dithane Rainshield	349				X		L	L								
Echo 720	330		X		X	X	L	CP, L				P				
Headline Duo	340		X		X	X	L	•		L, P		P	P			L
Headline EC	341	X	X	X	X	X	•	•			P	•	B FB, P	•	B FB, P	
Kocide 2000	334	X							•							
Lance	346	X	X	X	X	X		•		CP, L, P		P				B,CP,FB,L
Manzate Pro-Stick	349				X		L	L								
Parasol WG	334	X							•							
Penncozeb 75DF	349				X		L	L								
Pivot 418EC	355	X												•		
Priaxor DS	352		X		X	X	L	CP, L				P	P		P	(CP,L,P)*
Proline 480SC	353		X		X			CP, L								
Propel	355	X	X		X	X		CP, L				B,CP,L,P		•	B,CP,L,P	
Nufarm Propiconazole Fungicide	355	X												•		
Propulse	358	X	X		X			CP, L								B,CP,L
Quadris	324	X	X		X	X	•	•				P	B, P		B,CP,L,P	B,CP,L
Quash	362	X	X		X	X		(CP,L)*					P			(B,CP,L,P)*
Quilt	363	X	X		X	X	L					P	B,CP,L,P		B,CP,L,P	
Senator 70WP	372	X														B
Serenade Max/CPB	373	X	X		X	X			•*							(B,CP,L,P)*
Tilt 250E	355	X	X		X	X						B,CP,L,P		•	B,CP,L,P	
Vertisan	381	X	X		X	X		•		•					(B,CP,L,P)*	

* Suppression only

X Product registered for the crop

• Fungicide registered for disease control on all crops listed unless otherwise noted by the abbreviations listed: bean (B), chickpea (CP), fababean (FB), lentil (L), field pea (P)

Foliar Fungicide Product Pages

Acapela

Fungicide Group – 11
(Refer to page 312)

Company:

E.I. duPont Canada Company – PCP# 30470

Formulation:

250 g per L picoxystrobin formulated as a suspension concentrate.

Container size 2 x 9.6 L

Crops, Diseases, Rate and Timing:

Crop:	Diseases:	Application Rate (mL per acre):	Application Timing:
Dry legumes (chickpea, field pea, lentil, field, kidney, navy, lima, pinto, mung beans)	Control of <i>Mycosphaerella</i> blight (<i>Mycosphaerella pinodes</i>) on field pea; control of Asian soybean rust (<i>Phakospora pachyrhizi</i>)	240 to 350	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Suppression of <i>Sclerotinia</i> rot / white mould (<i>Sclerotinia sclerotiorum</i>)	350	For white mould make initial preventative application at early bloom and follow with second application 7 to 10 days later at full bloom.
Wheat	Leaf rust (<i>Puccinia recondita</i>), Septoria leaf blotch (<i>Septoria tritici</i>), powdery mildew (<i>Erysiphe graminis</i>), tan spot (<i>Pyrenophora tritici-repentis</i>)	175 to 350	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high. Apply at Feekes 9, "flag leaf out". Do not apply after flowering (Feekes 10.5)
Barley	Septoria leaf blotch (<i>Septoria tritici</i>), powdery mildew (<i>Erysiphe graminis</i>)	175 to 350	
Oats	Powdery mildew (<i>Erysiphe graminis</i>)	175 to 350	
Rye & triticale	Leaf rust (<i>Puccinia recondita</i>), Septoria leaf blotch (<i>Septoria tritici</i>), powdery mildew (<i>Erysiphe graminis</i>)	175 to 350	
Corn (field corn, sweet corn, seed popcorn)	Northern corn leaf blight (<i>Setosphaeria turcica</i> , <i>Exserohilum turcicum</i>)	215 to 325	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
Soybean	Asian soybean rust (<i>Phakospora pachyrhizi</i>), Brown spot (<i>Septoria glycines</i>), Frog-eye leafspot (<i>Cercospora soja</i>)	175 to 350	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Suppression of <i>Sclerotinia</i> stem rot / white mould (<i>Sclerotinia sclerotiorum</i>)	350	Initial preventative application at 100% bloom (1 flower blooming on all plants) and follow with 2 nd application 7 to 10 days later at full bloom.

Application Information:

Water Volume: Use sufficient water to obtain thorough coverage of plants.

Ground: minimum 45 L per acre.

Aerial: minimum 20 L per acre.

How it Works:

The active ingredient picoxystrobin is a broad spectrum strobilurin fungicide and is to be used as a preventative application when environmental conditions are favorable for disease development. Picoxystrobin has curative and locally systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Dry legumes – DO NOT exceed 1 application before switching to a fungicide with a different mode of action; maximum season use rate 700 ml per acre. Cereal grains, soybeans – DO NOT exceed 2 applications before switching to a fungicide with a different mode of action; maximum season use rate 1100

ml per acre. Corn – DO NOT exceed 2 applications before switching to a fungicide with a different mode of action; maximum season use rate for field, seed or popcorn is 1100 ml per acre and sweet corn is 1400 ml per acre.

Grazing: No restrictions listed.

Pre-harvest interval: Dry legumes and soybeans – 14 days, Cereal grains – 45 days, (7 days for forage, 14 days for hay), Corn – 7 days

Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: Crops that are on the product label may be replanted immediately after harvest. All other crops -10 months following last application of picoxystrobin.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closer. To prevent contamination, store this product away from food or feed.

Environment: Observe prescribed buffer zones. Minimize off-target drift to reduce the effects on beneficial insects at the field boundary. DO NOT apply to areas prone to run-off and delay spraying if heavy rainfall is forecast.

Hazard Rating:

None listed.

For an explanation of the symbols used here see page 11.

Acrobat 50 WP

Fungicide Group – 40
(Refer to page 312)

Company:

BASF Canada – PCP# 27700

Formulation:

50% dimethomorph formulated as a wettable powder.
Container size - 8 x 1.82 kg.

Crops, Diseases and Timing:

Control of late blight (*Phytophthora infestans*) and reduction of late blight tuber rot on potato.

Make the first application when the disease threatens or when the first visible signs of disease occur in the field or nearby. Apply every 5 to 7 days under high disease pressure or every 7 to 10 days under low disease pressure. A minimum of 5 days between applications is required. It is recommended to apply this product alternately with a fungicide having a different mode of action. Apply after top kill to control tuber rot under high levels of late blight infection.

Rate:

Apply at 182 g per acre. **DO NOT** apply *Acrobat 50 WP* alone. This fungicide must be tank mixed with one of the following fungicides: *Dithane Rainshield*, or *Bravo 500*, at the recommended product label rate.

Application Information:

Water Volume: Use sufficient water to obtain adequate spray coverage.

Ground: 20 to 40 L per acre for concentrate sprays to 91 to 648 L per acre for dilute sprays.

Aerial: optimum 20 L per acre.

How it Works:

The active ingredient dimethomorph is a Carboxylic Acid Amide (CAA) fungicide with contact, systemic and antisporeulant activity. To be used as a preventative and inhibitive (antisporeulant) fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides: Must be tank mixed with one of the following fungicides: *Dithane Rainshield*, or *Bravo 500*.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 3 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 4 days.

Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: DO NOT replant in treated area within 120 days of last application.

Storage: Store under cool, dry conditions in secure, well ventilated buildings away from food or feed.

Environment: DO NOT apply to terrain where there is a potential for surface runoff to enter aquatic systems. This product is highly toxic to aquatic organisms. DO NOT apply within 100 m of streams, ponds, rivers and lakes when applying by air and within 50 m when applying by ground. When using *Acrobat 50 WP* consult the labels of the tank mix partner and observe the largest buffer zone of the product used in the tank mix.

Hazard Rating:



Caution – Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Allegro 500F

Fungicide Group – 29
(Refer to page 312)

Company:

Syngenta Canada – PCP# 27517

Formulation:

40% fluazinam. Container size – 2 x 10 L.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Potato	Late blight (<i>Phytophthora infestans</i>)	162 mL	Begin applications when plants are 15 to 20 cm tall or when conditions favour disease development. Repeat application at 7 to 10 day intervals. DO NOT make more than 3 consecutive applications.
	White mould (<i>Sclerotinia sclerotiorum</i>)	162 to 242 mL	Begin applications at full bloom. Repeat application intervals of 7 to 10 days. When white mould pressure is low to moderate use 162 mL. When conditions favour moderate to high white mould pressure, increase the rate to 242 mL. DO NOT make more than 3 consecutive applications.
Dry shelled beans	White mould (<i>Sclerotinia sclerotiorum</i>)	240 to 405 mL	Begin applications when plants are at early to mid bloom (10 to 50% bloom). Repeat application 7 to 10 days later.
Soybeans	White mould (<i>Sclerotinia sclerotiorum</i>)	356 to 474 mL	Timing: Begin application at the R1 (early bloom) to R2 (full bloom) stage of development and if needed, again 10 to 14 days later at early pod formation (R3).

Application Information:

Water Volume: 80 to 240 L per acre. Spray volumes vary with amount of plant growth; apply in sufficient water to obtain adequate coverage of foliage.

How it Works:

The active ingredient fluazinam is a pyridinamine fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Note: Syngenta supports the following mixes that are not on the Allegro 500F label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Reglone (potatoes)

Fungicides: Quadris

Insecticides: Matador (dry beans)

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Bean and soybean – DO NOT exceed 2 applications of this product per season. Potato – DO NOT exceed 10 application of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 14 days (potatoes); 30 days (soybeans and dry shelled beans).

Re-entry: DO NOT re-enter treated areas within 24 hours of application.

Re-cropping: Can be replanted with potatoes as soon as practical after the last application, 30 days for other root crops and leafy vegetables, and 70 days for all other crops. Fluazinam will carry over, DO NOT use in areas treated with this product during the previous season.

Storage: Store product in a dry place separate from other pesticides, fertilizer, food, and feed.

Environment: DO NOT contaminate aquatic habitats when cleaning and rinsing spray equipment or containers. DO NOT overspray non-target terrestrial or aquatic habitats.

Hazard Rating:



Caution – Poison.



Warning – skin irritant. Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Astound

Fungicide Group – 9, 12
(Refer to page 312)

Company:

Syngenta Canada – PCP #29648

Formulation:

37.5% Cyprodinil and 25.0% Fludioxonil formulated as wettable granules. Container size - 2 x 6.28 kg.

Crops, Diseases and Timing:

Control of sclerotinia stem rot caused by *Sclerotinia sclerotiorum* in canola. One application at 20 to 50% flowering, optimum at 20 to 30%.

Rates:

Apply at 314 to 395 g per acre.

Application Information:

Ground and aerial application. Apply higher rate under conditions of high disease pressure.

Apply in sufficient water volume to obtain thorough coverage; a minimum spray volume of 81 L per acre is recommended for ground application or 18 L per acre for aerial application.

How it Works:

The active ingredient cyprodinil is a anilinopyrimidine fungicide and the active ingredient fludioxonil is a phenylpyrrole fungicide. Together they provide contact and systemic activity and inhibit spore germination and penetration. To be used as a preventative and curative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Astound fungicide can be tank mixed with *Matador 120EC* insecticide at a rate of 314 to 395 g per acre *Astound*

and 34 mL per acre *Matador 120EC* for control of insects and diseases on canola. Refer to labels for diseases and insects controlled, specific application instructions, and precautions. Pests and crops must be at the correct stage as specified on both labels. DO NOT apply more than 1 application per season of this tank mix.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: No restrictions listed.

Grazing: No restrictions listed.

Preharvest interval: DO NOT apply within 35 days of harvest.

Re-entry: DO NOT re-enter treated area within 12 hours of application.

Re-cropping: DO NOT plant any other crop for a period of 30 days after harvest or crop failure unless *Astound* is registered for that use.

Storage: Store in a dry place.

Environment: This product is toxic to aquatic organisms; DO NOT apply directly to aquatic habitats. For ground application, buffer zones must be 1m for protection of aquatic habitats. For aerial application (fixed and rotary wing), buffer zones must be 10 m for aquatic habitats less than 1 m deep and 2 m for aquatic habitats deeper than 1 m. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. DO NOT apply this product through any type of irrigation system.

Hazard Rating:



Caution – Eye irritant.

Keep out of reach of children.

For an explanation of the symbols used here see page 11.

Azoxystrobin

Fungicide Group – 11

(Refer to page 312)

Quadris / MANA Azoxystrobin

MANA Azoxystrobin 250 is currently only available as a co-pack with Bumper 418EC (see Blanket, page 326).

Company:

Syngenta Canada – (Quadris – PCP#26153)

MANA Canada – (MANA Azoxystrobin – PCP#30489)

Formulation:

250 g per L azoxystrobin formulated as a flowable suspension concentrate.

Container size – Quadris 4 x 3.78 L jugs.

MANA Azoxystrobin 4.8 L jug part of Blanket co-pack

Quadris ONLY – Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rates (per acre):	Application Timing:
Bean	Anthrachnose (<i>Colletotrichum lindemuthianum</i>); Powdery mildew (<i>Erysiphe polygoni</i>); Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	200 ml	Apply before disease is established and no later than onset of flowering; make 2nd application 10 to 14 days later.
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200 ml	Apply at the 2 to 6 leaf stage.
	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	280 to 400 ml	Apply at early bloom (prior to 30% bloom). This timing will also suppress alternaria black spot. Use the higher rate if there is a history of sclerotinia infection in the area and when conditions favour development.
	Alternaria black spot (<i>Alternaria brassicae</i> , <i>A. raphani</i>)	200 ml	Apply at pod stage (90% petal fall).
Chickpea	Ascochyta blight (<i>Ascochyta rabiei</i>), Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	200 ml	Apply before disease is established and no later than onset of flowering; make 2nd application 10 to 14 days later.
Coriander*	Blossom blight (<i>Aureobasidium</i> spp.)	180 to 450 ml	Apply once prior to disease establishment. Use high rate if high disease pressure.
Corn*	Rust (<i>Puccinia sorghi</i>)	180 ml	Apply before disease is established and make 2nd application 7 to 14 days later.
Lentil	Anthrachnose (<i>Collectotrichum truncatum</i>), Ascochyta blight (<i>Ascochyta lentis</i>), Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	200 ml	Apply before disease is established and no later than onset of flowering; make 2nd application 10 to 14 days later.
Pea	Mycosphaerella blight (<i>Mycosphaerella</i> , <i>Ascochyta</i> spp.), Powdery mildew (<i>Erysiphe pisi</i>), Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	200 ml	Apply before disease is established and no later than onset of flowering; make 2nd application 10 to 14 days later.

Continued

Quadris ONLY – Crops, Diseases, Rates and Timing continued:

Crop:	Diseases Controlled:	Application Rates (per acre):	Application Timing:
Potato	Early blight (<i>Alternaria solani</i>)	200 to 320 ml	Apply prior to disease development and repeat on a 7 to 14 day interval. Use the higher rate if extending treatment interval to 14 days. Apply in alternation with fungicides with a different mode of action. If late blight becomes established, discontinue use of <i>Quadris</i> and use alternative fungicides.
	Late blight (<i>Phytophthora infestans</i>)	320 ml	
	Rhizoctonia stem rot, Stolon canker, Black scurf (<i>Rhizoctonia solani</i>)*; Silver scurf (<i>Helminthosporium solani</i>)*	4 to 6 ml per 100 m of row	Apply once as an in-furrow spray in 20 to 56 L per acre water at planting. Mount the spray nozzle so that spray is directed into the furrow as a 15 to 20 cm band just before the seed is covered.
	Black dot (<i>Colletotrichum coccodes</i>)	200 to 320 ml	Apply on a 7 to 14 day interval prior to disease development. Use the high rate and short application interval under high disease pressures.
Soybean	Powdery mildew (<i>Microsphaera diffusa</i>), Cercospora leaf spot (<i>Cercospora kikuchi</i>), Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	200 ml	Apply at the R1 to R3 stage, or when 5% disease in the field; make 2nd application 14 days later.

* DO NOT apply by air.

MANA Azoxystrobin ONLY – Crops, Diseases, Rates and Timing:

MANA Azoxystrobin is only registered as a tank mix with Bumper 418EC fungicide on cereals. Refer to fungicide co-pack product Blanket on page 325 for details.

Application Information:**Water Volume:**

Ground: Use sufficient water volume to obtain adequate coverage. Use minimum 40 L per acre. *Quadris* only: in-furrow treatment in 20 to 56 L per acre.

Aerial: Use minimum of 18 L per acre. Ensure uniform application.

How it Works:

The active ingredient azoxystrobin is a methoxyacrylate compound (strobilurin) with broad spectrum contact and systemic activity. To be used as a preventative and curative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes

Insecticides: For cereals, legumes and field corn, *Quadris* may be tank-mixed with Matador 120EC insecticide. For control of potato diseases and insects, *Quadris* can be tank-

mixed with Actara 240 insecticide. Consult each label for pests controlled, appropriate timing, precautions, and specific application instructions.

Fungicides: MANA Azoxystrobin is only registered as a tank mix with Bumper 418EC fungicide on cereals. Refer to fungicide co-pack product Blanket on page 325 for details.

For control of Asian soybean rust in legumes, *Quadris* may be tank-mixed with Tilt 250E. For the control of early blight of potato, *Quadris* may be tank-mixed with Bravo 500. For control of Rhizoctonia stem, stolon canker and black scurf in potato, *Quadris* can be tank-mixed with Ridomil Gold 480EC. For control of ascochyta blight in chickpea, *Quadris* must be tank-mixed with Bravo 500 (minor use registration). *Quadris* may be tank-mixed with Tilt 250E in wheat and barley.

Note: Syngenta supports the following mixes that are not on the *Quadris* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: Touchdown Total, Traxion

Fungicides: Allegro 500E

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Coriander – DO NOT exceed 1 application of this product per season. Bean, canola, chickpea, corn, lentil, pea, soybean – DO NOT exceed 2 applications of this product per season. Potato – DO NOT exceed 3 applications of this product per season.

Grazing: DO NOT feed dried pea vines to livestock.

Preharvest interval: Canola - 30 days. Coriander - 21 days. Corn - 7 days. Legumes - 15 days. Potatoes - 1 day; 90 days for in-furrow application.

Re-entry: DO NOT re-enter treated areas until residues have dried.

Re-cropping: DO NOT plant broadleaf or root crops within 30 days of application. DO NOT plant cereals within 45 days of application.

Storage: Store in a cool, dry, well-ventilated area. DO NOT store below 0°C.

Environment: This product is toxic to fish and aquatic organisms. Observe buffer zones outlined in the label.

Hazard Rating:

None.

Other precautions: May irritate eyes.

Blanket

Fungicide Group – 3, 11

(Refer to page 312)

Blanket is a co-pack of **MANA Azoxystrobin** (PCP #30489), and **Bumper** (PCP #28017; see propiconazole on page 355). Information listed is restricted to crops, diseases, and rates. For other detailed information on the component products, refer to the product labels and the product page listed above.

Company:

MANA Canada

Formulation:

Blanket has two components:

MANA Azoxystrobin 250 (PCP #30489): 250 g per L azoxystrobin formulated as a suspension. (4.8 L jug)

Bumper 418 EC (PCP #28017): 418 g per L propiconazole formulated as an emulsifiable concentrate (4.8 L jug)

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rates (per acre):		Application Timing:
		MANA Azoxystrobin	Bumper	
Barley	Net blotch (<i>Pyrenophora teres</i>), scald (<i>Rhynchosporium secalis</i>), and barley leaf rust (<i>Puccinia hordei</i>).	90 to 120 mL	120 mL	Apply once between stem elongation and half-head emergence (BBCH Growth Stage 29-55). DO NOT make more than one application per season of this tank mixture. An additional application of <i>Bumper 418 EC</i> Fungicide can be made, if required. Refer to the <i>Bumper 418 EC</i> Fungicide label for details of the rate and timing. A total of two applications of <i>Bumper</i> may be applied per season either in a tank mix with MANA <i>Azoxystrobin</i> or alone.
Wheat, barley, rye	Septoria leaf spot (<i>Septoria</i> sp.) and tan spot (<i>Pyrenophora tritici-repentis</i>)			
Winter wheat, spring wheat, and barley	Stripe rust (<i>Puccinia striiformis</i>) and wheat leaf rust (wheat only) (<i>Puccinia triticina</i>)	80 to 120 mL	100 to 120 mL	

Restrictions:

See product labels and pages listed above for restrictions and recommendations for resistance management. Follow the most stringent restrictions for either product.

Hazard Rating:



Warning – Poison



Warning – Eye and Skin Irritant. Potential Skin Sensitizer.

For an explanation of the symbols used here see page 11.

Cabrio Plus

Fungicide Group – 11, M3
(Refer to page 312)

Company:

BASF Canada – PCP#30395

Formulation:

5% pyraclostrobin and 55% metiram formulated as a water dispersible granule.

Container size – 20 kg.

Crops, Diseases and Timing:

Control of early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*) in potato. Application should begin prior to row closure or when conditions become favourable for the development of disease. For early blight apply on a 7 to 14 day interval and for late blight apply on a 7 to 10 day interval.

Rates:

Apply at 0.91 to 1.35 kg per acre.

Application Information:

Water Volume:

Ground: Minimum of 80 L per acre.

Aerial: Minimum of 20 L per acre.

How it Works:

The active ingredient metiram is a dithiocarbamate fungicide with contact activity. The active ingredient pyraclostrobin is a member of the strobilurin class of chemistry

used as a broad spectrum fungicide. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 3 applications of this product per season.

Grazing: Crop can be grazed or fed to livestock.

Preharvest Interval: 3 days.

Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: A plant back interval of 14 days is required for all crops not listed on the label.

Storage: Store in original tightly closed container.

Environmental Hazards: Avoid run-off from treated areas into aquatic areas. Toxic to aquatic organisms, non-target terrestrial plants and small wild animals.

Hazard Rating:



Warning – Poison



Warning – Eye and Skin Irritant.

For an explanation of the symbols used here see page 11.

Cantus WDG Fungicide

Fungicide Group – 7
(Refer to page 312)

Company:

BASF Canada – PCP#30141

Formulation:

70% boscalid formulated as a water dispersible granules.
Container size – 4 x 2.83 kg.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre)*:	Application Timing:
Potato	Early blight (<i>Alternaria solani</i>)	71 to 127 g	Apply prior to disease development and at 14 day intervals if conditions continue to favour disease development.

Application Information:

Water Volume:

Ground: Use a minimum water volume of 40 L per acre and ensure thorough coverage of foliage.

Aerial: Use a minimum water volume of 16 L per acre and ensure thorough coverage of foliage.

Pivot and Sprinkler Irrigation: DO NOT exceed 0.64 cm (1/4 inch) or 25,700 L per acre. Apply only through sprinkler systems including centre pivot, lateral move, end two, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. The system must contain functional valves to prevent water source contamination from back-flow.

How it Works:

The active ingredient boscalid is a carboxamide (SDHI) fungicide with systemic activity. It inhibits spore germination, mycelia growth and sporulation of the fungus on the leaf surface. To be used as a preventative application when environmental conditions are favourable for disease development. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: 4 applications per season on potatoes.

Grazing: No restriction listed.

Preharvest interval: Potato - 30 days

Re-entry: DO NOT re-enter treated area for 12 hours after application or until dry.

Re-cropping: A plant back restriction of 14 days is required for all crops not on the label.

Storage: Store in a cool, dry, locked, well-ventilated area without a floor drain.

Environment: Toxic to aquatic organisms and non-target terrestrial plants. DO NOT apply to areas where runoff is likely to occur, or near any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Hazard Rating:



Caution Poison – Potential Skin Sensitizer.



Warning – Eye irritant.

For an explanation of the symbols used here see page 11.

Caramba

Fungicide Group – 3

(Refer to page 312)

Company:

BASF Canada – FCI# 29767

Formulation:

90 g per L metconazole formulated as an emulsifiable concentrate.

Container size – Case (2 x 8.1L); 128 L drum; or 400 L tote.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre)*:	Application Timing:
Wheat (all types) Triticale	Leaf rust (<i>Puccinia triticina</i>), tan spot (<i>Pyrenophora tritici-repentis</i>), septoria leaf spot (<i>Septoria</i> sp.), suppression of spot blotch (<i>Cochliobolus sativus</i>)	202 to 283 mL	Prior to disease development or at onset of disease.
Barley	Net blotch (<i>Pyrenophora teres</i>), scald (<i>Rhynchosporium secalis</i>), leaf rust (<i>Puccinia hordei</i>) and suppression of spot blotch (<i>Cochliobolus sativus</i>)		
Oats	Crown rust (<i>Puccinia coronata</i>), septoria leaf blotch (<i>Septoria avenae</i>)		
Rye	Leaf rust (<i>Puccinia recondita</i>)		
Wheat Oats Rye Barley	Suppression of fusarium head blight (FHB) (<i>Fusarium</i> spp.)	405 mL	To be used as a preventative fungicide application when environmental conditions are favourable for disease development. Fusarium head blight outbreaks are favoured by warm and wet weather when the cereal crop is at the flowering to soft dough stage. For optimum suppression of FHB, apply within the time period when at least 75% of the heads for wheat, oats and rye on the main stem are fully emerged to when 50% of the heads on the main stem are in flower. For barley apply between full head emergence and up to 3 days after full emergence of main stem heads.
Soybeans	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	283 mL	Initial application between early flowering and pod set (R1 and R3 growth) or prior to rust development.

* In cereals a case can treat 40 acres after heading (suppression of FHB) or 60 to 80 acres before heading (leaf disease). In cereals a drum can treat 320 acres after heading (suppression of FHB) or 460 to 640 acres before heading (leaf disease).

Application Information:

Water Volume:

Ground: Minimum of 40 L per acre.

Aerial: Minimum of 20 L per acre.

Consult nozzle manufacturers for specific nozzle and pressure recommendations.

How it Works:

The active ingredient metconazole is a broad spectrum triazole demethylation inhibitor (DMI) fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None listed.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Wheat, oats, rye, barley – DO NOT exceed 1 application of this product per season. Soybean – DO NOT exceed 2 applications of this product or other DMI (Group 3) fungicides per season.

Grazing: All crops can be grazed or fed to livestock.

Preharvest interval: Wheat, barley, oats, rye and soybeans 30 days and sugar beets 14 days.

Re-entry: Soybean – DO NOT re-enter treated areas within 4 days of application. Wheat, barley, oats, rye – DO NOT re-enter treated areas within 5 days of application.

Re-cropping: A plant back interval of 35 days is required for all crops not listed on the label.

Storage: Store in original tightly closed container. Protect from freezing.

Environment: Avoid run-off from treated areas into aquatic areas. Toxic to aquatic organisms, non-target terrestrial plants and small wild animals.

Hazard Rating:



Warning – Eye irritant.

Check label for first-aid information.

For an explanation of the symbols used here see page 11.

Chlorothalonil

Fungicide Group – M5
(Refer to page 312)

Bravo 500/Bravo Zn/Echo 720/Echo 90DF

Company:

Syngenta Canada (Bravo 500 – PCP#15723, Bravo Zn – PCP#28900)

United Agri Products (Echo 720 – PCP#29355, Echo 90DF – PCP#29356)

Formulation:

Bravo 500 - 500 g per L chlorothalonil formulated as a suspension. Container sizes - 2 x 10 L case and 450 L.

Bravo Zn - 500 g per L chlorothalonil formulated as a suspension. Container sizes - 10 L and 450 L.

Echo 720 - 720 g per L chlorothalonil formulated as a suspension. Container size - 2 x 9.46 L case, 450 L and 984.1 L.

Echo 90DF - 90% chlorothalonil formulated as a dry flowable. Container size - 10 kg.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate* (per acre):		Application Timing:
		Bravo 500	Echo 720	
Wheat	Tan spot (<i>Pyrenophora tritici-repentis</i>), Septoria glume blotch, Septoria leaf spot (<i>Septoria</i> sp.)	600 to 1000 mL	405 to 690 mL	Begin application at flag leaf emergence; repeat 10 to 14 days later when heads are visible. A third application when heads are fully emerged may be necessary.
	Suppression of Fusarium head blight (scab) (<i>Fusarium</i> spp.)	800 to 1000 mL	570 to 690 mL	For suppression of fusarium head blight apply at early flowering (before flowering has started in the majority of tillers) and before the beginning of weather favouring disease.
Pea	Mycosphaerella blight (<i>Mycosphaerella pinodes</i>)	800 to 1200 mL	570 to 850 mL	Begin application at early flowering and repeat 10 days later at early pod set or mid-flowering if necessary. Make a third application 10 to 14 days after the second application at pod fill or later flowering should conditions remain favourable for disease.

Continued

Crops, Diseases, Rates and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate* (per acre):		Application Timing:
		<i>Bravo 500</i>	<i>Echo 720</i>	
Lentil	Ascochyta blight (<i>Ascochyta lentis</i>), Anthracnose (<i>Colletotrichum truncatum</i>)	800 to 1600 mL	570 to 1130 mL	For one application only, apply at early flowering. For two applications: apply first before flowering when bud formation is evident; apply second at early to mid-flowering 10 to 14 days after the first application but before rows close in.
Chickpea	Ascochyta blight (<i>Ascochyta rabiei</i>)	1200 to 1600 mL for first application; 800 to 1200 mL for subsequent applications.	850 to 1130 mL for first application; 570 to 850 mL for subsequent applications.	Make first application at very early flowering and remaining applications at 10 day intervals.

Crop	Diseases Controlled:	Application Rate* (per acre):			Application Timing:
		<i>Bravo 500</i>	<i>Bravo Zn</i>	<i>Echo 720</i>	
Potato	Late blight (<i>Phytophthora infestans</i>)	480 to 1000 mL	480 to 1000 mL	320 to 690 mL	Begin application when plants are 6 to 8 inches (15 to 20 cm) high or when disease threatens. Repeat applications at 7 to 10 day intervals or as necessary to maintain disease control. * Under high disease pressure, use higher rate and shorter spray intervals.
	Early blight (<i>Alternaria solani</i>)	640 to 1000 mL	640 to 1000 mL	445 mL	
	Botrytis vine rot (<i>Botrytis cinerea</i>)	640 to 1000 mL	-	480 to 1000 mL	

Crop	Diseases Controlled:	Application Rate* (per acre): <i>Echo 90DF</i>	Application Timing:
Potato	Late blight (<i>Phytophthora infestans</i>)	280 to 530 grams	Begin application when plants are 6 to 8 inches (15 to 20 cm) high or when disease threatens. Repeat applications at 7 to 10 day intervals or as necessary to maintain disease control. * Under high disease pressure, use higher rate and shorter spray intervals.
	Early blight (<i>Alternaria solani</i>), Botrytis vine rot (<i>Botrytis cinerea</i>)	370 to 530 grams	

Application Information:

Water Volume: Volume will vary with crop and amount of plant growth. Use sufficient water to obtain adequate coverage of foliage.

Ground: Spray volume will usually range from 90 to 640 L per acre for dilute sprays and 20 to 40 L per acre for concentrate sprays.

Chickpea - 90 L per acre. Ground application only.

Aerial: Use minimum of 12 L per acre.

How it Works:

The active ingredient chlorothalonil is a chloronitrile fungicide with multi-site contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

DO NOT combine with pesticides, surfactants or fertilizers unless prior use has shown the combination is physically compatible and non-injurious under your conditions of use.

Fungicides: For control of early blight in potato, *Bravo 500* may be tank-mixed with 200 mL per acre *Quadris*. DO NOT apply sequential applications of this tank-mix and DO NOT exceed 3 tank-mix applications per season. DO NOT apply to potatoes later than 2 days before harvest. For control of early blight, late blight, and botrytis vine rot in potato and for suppression of storage rots, pythium leak and pink rot, in potato *Bravo Zn* may be tank mixed with 80 mL per acre *Ridomil Gold 480 EC* or *Ridomil Gold 480 SL*.

Herbicides: On lentils, DO NOT apply in combination with *Poast* herbicide and *Merge* surfactant or within 48 hours of the application of *Poast* and *Merge*.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Lentil – DO NOT exceed 2 applications of this product per season. Wheat, pea, chickpea – DO NOT exceed 3 applications of this product per season. Potato (*Echo 901D1*) – DO NOT exceed 12 applications of this product per season.

Grazing: DO NOT graze treated areas. DO NOT feed straw from treated crop to livestock.

Preharvest interval: Potato – 1 day; Lentil – 48 days;

Chickpea – *Bravo 500* – 14 days, *Echo 720* – 48 days; Wheat – 30 days; Pea – 32 days.

Re-cropping: None.

Re-entry: DO NOT re-enter treated area within 48 hours of application. If required, and at least 4 hours have passed since application, individuals may re-enter treated area for short-term tasks not involving hand labour. Long pants, long-sleeved shirt, and chemical resistant gloves must be worn.

Storage: DO NOT store near feed or food stuffs. Store in a cool, dry, ventilated place. Protect from excessive heat.

Environment: DO NOT apply if weather conditions favour drift from area being treated. DO NOT contaminate lakes, streams or ponds. Observe a buffer zone of 100 m for aerial applications and 15 m for ground applications to protect aquatic systems.

Hazard Rating:



Caution – Poison.



Warning – causes severe eye damage.

For an explanation of the symbols used here see page 11.

Contans WG

Fungicide Group –
Not classified, bio-fungicide.

Company:

Prophyta Biologischer Pflanzenschutz, distributed by United Agri Products - PCP#29066.

Formulation:

Wettable Granules - 5.30% *Coniothyrium minitans* strain CON/M/91-08. Contains minimum of 1×10^9 cfu per gram.

Container size - 20kg.

Crops, Diseases Controlled, Rates and Timing

Crop:	Diseases Suppressed:	Application Rate (per acre):	Application Timing:
Pre-plant - Soils where Canola, Sunflower, Dry Edible Bean or Soybean will be planted	White mould or stem rot (<i>Sclerotinia sclerotiorum</i>) and rots caused by <i>S. minor</i>	0.4 to 0.8 kg	Prior to planting of spring crop; three months before the typical onset of sclerotinia white mould or stem rot. In fall, prior to spring planting of susceptible crop. After application to the soil, the product should be incorporated to within 5 cm of the top-soil. Incorporation should take place as soon as possible after application (within 1 week maximum).
		0.8 to 1.6 kg	If soil incorporation is to a depth greater than 5 cm, higher rate should be applied.
Post-harvest - On harvest residue of susceptible crops		0.2 to 0.4 kg	Prior to the next soil treatment, the residues of the susceptible crops in rotation can be also treated to help reduce inoculum loads of sclerotia in the field.

Application Information:

Use sufficient water volumes to give thoroughly coverage of the soil surface and/or the crop residue (10 gallons per acre of water volume).

DO NOT allow spray mixture to stand overnight or for prolonged periods; should be used within 24 hours of being prepared.

After incorporation treated soils should not be disturbed to avoid bringing untreated sclerotia from lower soil depths to the top soil layer.

As part of an overall long term pest management strategy it is recommended to use other management practices along with *Contans* such as in season foliar fungicide applications and proper crop rotations.

DO NOT apply by air.

How it Works:

The active ingredient, *Coniothyrium minitans* is a fungus that infects the sclerotia of *Sclerotinia sclerotiorum* and *S. minor*. Infection of sclerotial bodies prevents production of ascospores and mycelial structures that infect plants. Regular use of *Contans* in successive years within a long-term management strategy will improve disease control.

Tank Mixes:

DO NOT tank-mix with fungicides or fertilizers. Also, DO NOT tank mix with acids, alkalis or any product that

attacks organic materials. Contact UAP Canada for more information on what products are compatible with *Contans*.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: No restrictions listed.

Grazing: No restrictions listed.

Preharvest interval: Can be applied up to and including the day of harvest.

Re-entry: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Maximum storage period of one year at 4°C or below. Up to 6 weeks at temperatures between 4°C and 23°C. Store in a dry area inaccessible to children. Store in original container away from food or feed.

Environment: DO NOT apply this product directly to freshwater habitats, estuarine/marine habitats. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:

 Caution – Potential sensitizer.

For an explanation of the symbols used here see page 11.

Copper

Copper 53W/Copper Spray/Kocide 2000/Parasol WG

Fungicide Group – M1
(Refer to page 312)

Company:

United Agri Products, (Copper 53W – PCP#09934, Copper Spray – PCP#19146)

E.I. du Pont Canada Company, (Kocide 2000 – PCP#27348)

Nufarm Agriculture Inc. (Parasol WG – PCP#29063)

Formulation:

Copper 53 W - 53% tribasic copper sulphate formulated as a wettable powder. Container size - 10 kg.

Copper Spray - 50% copper oxychloride formulated as a wettable powder. Container size - 10 x 2 kg case.

Kocide 2000 - 35% copper as copper hydroxide formulated as a dry flowable. Container size - 10 to 25 kg.

Parasol WG - 50% elemental copper as copper hydroxide formulated as a wettable granule. Container size - 10 kg.

Crops, Diseases and Timing:

Potatoes - Control of early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*).

Copper 53W: Apply when plants are 5 to 7 inches (12 to 18 cm) tall. Repeat at 7 to 10 day intervals.

Copper Spray: Apply when plants are 4 to 8 inches (10 to 20 cm) tall. Repeat at 7 to 10 day intervals.

Kocide 2000: Apply when plants are 15 cm tall at 7 to 10 day intervals, until harvest.

Parasol WG: Apply when plants are 6 inches (15 cm) tall. Apply combined with 0.7 to 0.9 kg of a mancozeb (80%) product per acre, at 7 to 10 day intervals. *Kocide 2000* and *Parasol* may be applied with a desiccant at vine kill or alone after vine kill, prior to harvest, to reduce the risk of late blight tuber infection. See table for rates.

Beans - Control of anthracnose (*Copper 53W*), downy mildew (*Copper 53W*), and bacterial blight (common and halo) (*Copper 53W*, *Kocide 2000*, *Parasol WG*).

First application when plants are 6 inches (15 cm) tall, as a protectant. Repeat every 7 to 14 days depending on local conditions.

See label for more information on user requested minor use label expansion (URMULE) for use as a seed treatment on beans:

For suppression of the following seed borne diseases; common blight (*Xanthomonas axonopodis* pv. *phaseoli*), halo blight (*Pseudomonas syringae* pv. *phaseolicola*) and bacterial brown spot (*Pseudomonas syringae* pv. *syringae*), in dry edible beans and succulent beans.

For every 100 kg seed to be treated, add 113 g *Kocide 2000* to 200 ml of warm water and stir until dissolved. Ensure even coverage and allow treated seed to dry before planting. Treated seed must be labelled as follows: "This seed has been treated with *Kocide 2000*; DO NOT use treated seed for food, feed or oil processing".

Kocide 2000 used as a seed treatment may cause some delay in seed germination. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment.

All seed treated with this product must be conspicuously coloured at the time of treatment. When handling contaminated equipment or treated seed, wear long sleeved coveralls over normal work clothing, chemical resistant gloves and goggles; use an approved respirator with a canister approved for pesticides to minimize exposure to dust when handling, mixing leading product, bagging seed, sewing bags of treated seed or when transferring treated seed to a storage bin.

Refer to the label for additional application instructions and/or use precautions regarding this URMULE.

Rates:

Product	Potato (kg per acre)	Acres treated by 10 kg container	Beans (kg per acre)	Acres treated by 10 kg container
<i>Copper 53W</i>	2.2	4.5	2.2	4.5
<i>Copper Spray</i>	1.6	6.25	—	—
<i>Kocide 2000</i>	0.32 to 0.64 0.967 (vine kill)	31.3 to 15.4 10.34 (vine kill)	0.65 to 0.93	15.4 to 10.7
<i>Parasol WG</i>	0.44 to 1.0 1.36 (vine kill)	23 to 10 7.3 (vine kill)	0.9 to 1.3	11 to 7.7

Application Information:

Water Volume:

Ground:

Kocide 2000 - Enough to ensure thorough coverage.

Copper 53W, *Copper Spray* - 88 gallons per acre (400 L per acre).

Parasol(s) - Enough to ensure thorough coverage.

Aerial: DO NOT apply *Copper Spray*, *Copper 53W*, *Kocide 2000*, and *Parasol WG* by air.

How it Works:

The active ingredients tribasic copper sulphate, copper oxychloride and copper hydroxide are inorganic fungicides with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides: *Kocide 2000* - Can be combined with 0.7 to 0.9 kg per acre of *Manzate Pro-Stick*; and with a desiccant for vine kill.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Bean (*Kocide 2000*, *Parasol WG*) - DO NOT exceed 6 applications of this product per season. Potato (*Kocide 2000*, *Parasol WG*) - DO NOT exceed 10 applications of this product per season.

Labelling (*Kocide 2000*): Treated seed must be labeled as follows: "This seed has been treated with *Kocide 2000*; DO NOT use treated seed for food, feed or oil processing."

Grazing: No restrictions listed.

Preharvest interval: 1 day.

Re-entry: DO NOT re-enter treated areas within 48 hours of application.

Re-cropping: No restrictions listed.

Storage: Store in cool, dry, ventilated area, away from feed or food. Keep away from heat, fire and sparks.

Environment: DO NOT apply or allow to drift onto streams or any body of water.

Hazard Rating:



Warning - Poison - (*Copper 53W*, *Copper Spray*)



Caution - Poison - (*Kocide 2000*, *Parasol WG*)



Warning - Eye irritant (*Kocide 2000*)

For an explanation of the symbols used here see page 11.

Curzate 60 DF

Fungicide Group – 27
(Refer to page 312)

Company

E.I du Pont Canada Company - PCP#26284

Formulation:

60% cymoxanil formulated as a dry flowable.

Container size - 1.8 kg

Crops, Diseases Timing:

Control of late blight (*Phytophthora infestans*) in potato. Initial applications should start when local conditions indicate that late blight is imminent. Make additional applications at 5 to 7 day intervals; however at least 20 days must pass between the 2nd and 3rd application.

Rate:

Apply Curzate 60 DF at 90 g per acre

Plus

Manzate DF or Manzate Pro-Stick at 540 g to 650 g per acre

Application Information:

Water Volume: Utilize sufficient water to obtain thorough coverage - 80 to 400 L per acre.

Aerial Application: DO NOT apply by air.

How it Works:

The active ingredient cymoxanil is a cyanoacetamide-oxime fungicide with locally systemic activity. To be used as a preventative, curative and inhibitive (antisporeulant) fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

DO NOT use Curzate 60 DF alone. Use only in a tank mix with Manzate DF or Manzate Pro-Stick.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 7 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 8 days

Re-entry: DO NOT re-enter treated area within 24 hours of application.

Re-cropping: No restrictions listed.

Storage: Store product in original container in a secure, dry area away from food or feed. Protect against humid air and water. Not for use or storage in or around the home. Keep container tightly closed.

Environment: A buffer zone of 50 m is required between the down-wind edge of the boom and sensitive aquatic habitats such as ponds, lakes, rivers, streams, and wetlands. DO NOT contaminate these habitats when cleaning and rinsing equipment or containers. DO NOT clean sprayer near well or water source or near desirable vegetation.

Hazard Rating:



Danger – Poison.



Caution – Eye irritant.

For an explanation of the symbols used here see page 11.

Cyazofamid

Ranman 400SC/Torrent 400SC

Fungicide Group – 21
(Refer to page 312)

Company:

ISK Biosciences Corporation; distributed by FMC of Canada
(Ranman 400SC– PCP# 27984)

ISK Biosciences Corporation; distributed by Engage Agro
Corporation (Torrent 400SC– PCP# 30392)

Formulation:

400 g per L cyazofamid formulated as a suspension concentrate.

Container size: Ranman 400SC - 500ml and 200L.
Torrent 400SC - 6 x 1 L and 4 x 4L.

Crops, Diseases and Timing:

Control of late blight (*Phytophthora infestans*) on potato. Begin applications on a 7 day schedule when warning systems forecast disease infection periods or at row closure. Use the low rate under low disease pressure and increase the rate as disease pressure and/or crop development increases, up to the maximum rate. For late blight tuber rot control, ensure that the last 2 to 3 applications prior to desiccation are made at the maximum rate following resistance management practices.

Rates:

0.04 to 0.08 L per acre. Ranman 400SC and Torrent 400SC should be tank mixed with a non-ionic or organo-silicone surfactant (such as Sylgard 309 at 0.06 L per acre).

Application Information:

DO NOT apply by air. DO NOT make sequential applications. After one application alternate with at least one application of fungicide with a different mode of action.

Water Volume: Use sufficient volume to obtain coverage of the foliage, 80 to 240 L per acre.

How it Works:

The active ingredient cyazofamid is a cyanoimidazole fungicide with contact activity. To be used as a preventative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 6 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 7 days.

Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: A plant back interval of 30 days is required.

Storage: Store product in original container in a secured dry place separate from other pesticides, fertilizer, food and feed.

Environment: Toxic to aquatic organisms. The specified buffer zones (Field sprayer, 5 m; Field sprayer with use of shrouds, 2 m; Field sprayer with use of cones, 4 m) are required between the point of direct application and the closest downwind edge of sensitive freshwater habitats.

Hazard Rating:

None listed.

Folicur 432 F/Folicur 250 EW (2013)

See Tebuconazole on page 377.

Fontelis

Fungicide Group - 7
(Refer to page 312)

Company:

E.I. duPont - PCP#30331

Formulation:

200 g per L penthiopyrad formulated as a suspension.
Container size - 4 x 3.79 L jug.

Crops, Diseases, Rates and Timing:

Crop:	Disease Controlled:	Application Rate (per acre)*:	Application Timing:
Alfalfa	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	500 to 700 mL	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.

Application Information:

Water Volume:

Ground: 45 L per acre.

Aerial: 16 L per acre.

Use sufficient water to obtain thorough coverage of plants. DO NOT apply under periods of dead calm. Avoid application of this product when winds are gusty.

How it Works:

The active ingredient penthiopyrad is a carboxamide fungicide with broad spectrum, locally systemic and curative properties recommended for foliar and soil borne plant diseases. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed two sequential applications of this product before switching to a fungicide with a different mode of action. DO NOT exceed 1.4 L per acre in one season.

Grazing: No restrictions listed.

Preharvest interval: 14 days.

Re-entry: DO NOT re-enter treated areas until 12 hours after application.

Re-cropping: Crops and crop groups on the *Fontelis* label as well as the following crops may be planted immediately after harvest: canola, cereal grains crop group, corn, cotton, legume vegetables crop subgroup, soybean, sugarbeet, tuberous and corm vegetables and leaves crop subgroup. All other crops cannot be planted until 12 months after the last application.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

Environment: This product is toxic to aquatic organisms. When using *Fontelis* consult the product label for buffer zones.

Hazard Rating:

Potential Skin Sensitizer.

Gavel 75 DF

Fungicide Group – 22, M3

(Refer to page 312)

Company:

Gowan Canada - PCP# 26842

Formulation:

66.7% mancozeb and 8.43% zoxamide formulated as a dry flowable. Container sizes - 3.5 kg, 13.6 kg, 20 kg, 25 kg.

Crops, Diseases and Timing:

Control of early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*) in potato. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program. Begin applications at the first sign of disease or when blight is reported in the area. Apply at 0.90 kg per acre every 7 days under high disease pressure when either disease is present or environmental conditions favour continued disease development.

Apply at 0.70 kg per acre every 7 days under low disease pressure and environmental conditions unfavorable for disease development.

Rate:

Apply at 0.70 to 0.90 kg per acre.

Application Information:

Thorough, uniform coverage is essential for good disease control.

Water Volume:

Ground: 90 L per acre.

Aerial: 18 to 36 L per acre. Use 36 liters of water under high disease pressure to provide better crop coverage.

How it Works:

To be used as a preventative fungicide application. The active ingredient zoxamide is a benzamide fungicide with contact activity. The mancozeb component is a dithio-carbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 6 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 3 days.

Re-entry: DO NOT re-enter treated areas within 48 hours of application.

Re-cropping: A 30 day plant back interval (PBI) is required for leafy vegetables and root and tuber vegetables. For all other crops not included on the label, the PBI should be 140 days.

Storage: DO NOT allow product to freeze. Keep away from fire and sparks. Store in a cool, dry, well ventilated place away from feed or food.

Environment:

Ground application: a buffer zone of 25 m for application by ground sprayer should be established between the last spray swath and the edge of aquatic systems. A buffer zone of 5 m for application by ground sprayer should be established between the last spray swath and the edge of terrestrial habitats such as hedgerows, windbreaks, woodlots, vegetative strips and other vegetation. This pesticide is toxic to fish.

Aerial application: a buffer zone of 20 m is required between the downwind edge of the boom and the closest edge of sensitive aquatic habitats.

Hazard Rating:



Caution – causes moderate eye irritation.

For an explanation of the symbols used here see page 11.

Headline DUO

Fungicide Group – 7, 11
(Refer to page 312)

Company:

BASF Canada – PCP#28862 (Headline Duo One) and PCP#28863 (Headline Duo Two)

Formulation:

Headline Duo One contains 25.2% boscalid and 12.8% pyraclostrobin formulated as a water dispersible granule. Headline Duo Two contains 70% boscalid formulated as a water dispersible granule.

Container size - Headline Duo One 6.4 kg + Headline Duo Two 1.16 kg. Jugs are split internally and contain both Headline Duo One and Headline Duo Two. Each split jug treats 20 acres (2 per case).

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Rates:
Chickpea	Ascochyta blight (<i>Ascochyta rubiei</i>) Ascochyta blight can develop quickly once established so early detection is essential.	Headline Duo One must be tank-mixed at a rate of 0.320 kg per acre with Headline Duo Two at a rate of 0.058 kg per acre at the beginning of flowering OR at the onset of symptoms. Each split jug of Headline DUO treats 20 acres (2 per case).
Field pea	Ascochyta Blight (<i>Ascochyta</i> spp.); gray mould (<i>Botrytis cinerea</i>); mycosphaerella blight (<i>Mycosphaerella</i> spp.); powdery mildew (<i>Erysiphe</i> spp.)	
Lentil	Ascochyta blight (<i>Ascochyta</i> spp.); white mould (<i>Sclerotinia sclerotiorum</i>); gray mould (<i>Botrytis cinerea</i>); anthracnose (<i>Colletotrichum</i> spp.)	

Application Information:

DO NOT apply sequential applications of this tank mixture combination; alternate to a fungicide with a mode of action other than Group 7 or 11 for at least one application.

Water Volume:

Ground: Use a minimum water volume of 40 L per acre on chickpeas. Ensure thorough coverage of foliage.

Aerial: Use a minimum water volume of 20 L per acre. Ensure thorough coverage of foliage

How it Works:

The active ingredient pyraclostrobin is a strobilurin fungicide with broad spectrum contact and systemic activity. The active ingredient boscalid is a carboxamide (SDHI) fungicide with systemic activity. To be used as a preventative application when environmental conditions are favourable for disease development. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 2 applications of this tank mixture per season.

Grazing: Chickpeas can be grazed or fed to livestock.

Preharvest interval: 30 days.

Re-entry: DO NOT re-enter treated areas within 48 hours of application.

Re-cropping: Wait 14 days before planting crops not on the label.

Storage: Store in a cool, dry, locked, well-ventilated area without a floor drain. DO NOT freeze.

Environment: Avoid overspray or drift to sensitive habitats. Maintain specified buffer zones.

Hazard Rating:



Danger Poison – Skin and eye irritant.

For an explanation of the symbols used here see page 11.

Headline EC

Fungicide Group – 11
(Refer to page 312)

Company:

BASF Canada – PCP#27322

Formulation:

250 g per L of pyraclostrobin formulated as an emulsifiable concentrate.

Container size - Case (2 x 6.5 L); 120 L shuttle; 400 L tote.

Crops, Diseases, Rates, and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Wheat	Tan spot (<i>Pyrenophora tritici-repentis</i>); Septoria leaf spot (<i>Septoria tritici</i> ; <i>S. nodorum</i>); Leaf rust (<i>Puccinia recondita</i>)	120 to 240 mL	<p>Wheat, barley, rye and oats: For control of leaf diseases apply single application immediately after flag leaf emergence; use higher rate to obtain extended protection; if disease persists or weather conditions are favourable for disease development, apply a second application 10 to 14 days later with a fungicide that contains a different mode of action.</p> <p>To maximize yields in cereals, it is important to protect the flag leaf from disease.</p>
	Powdery mildew (<i>Erysiphe graminis</i> f. sp. <i>tritici</i>); Spot blotch (<i>Cochliobolus sativus</i>); Stripe rust (<i>Puccinia striiformis</i>)	160 to 240 mL	
Barley	Net blotch (<i>Pyrenophora teres</i>)	120 to 240 mL	
	Scald (<i>Rhynchosporium secalis</i>); Spot blotch (<i>Cochliobolus sativus</i>); Stripe rust (<i>Puccinia striiformis</i>)	160 to 240 mL	
Rye	Leaf rust (<i>Puccinia recondita</i>)	120 to 240 mL	
	Powdery mildew (<i>Erysiphe graminis</i>)	160 to 240 mL	
Oat	Crown rust (<i>Puccinia coronata</i>)	120 to 160 mL	<p>Apply in tank mix with supported canola herbicides to control blackleg at the 2 to 6-leaf (rosette) stage. Apply to control alternaria black spot at 20 to 50% bloom (suppression) to early pod stage (90% bloom) for control.</p> <p>Can be tank-mixed with <i>Lance</i> WDG Fungicide at 20 to 50% flower to control sclerotinia stem rot and suppress black spot.</p>
Canola, rape-seed, canola quality <i>Brassica juncea</i> , mustard (oilseed and condiment)	Black spot (<i>Alternaria brassicae</i> and <i>A. raphani</i>) Blackleg (<i>Leptosphaeria maculans</i>)	120 to 160 mL	
Corn	Common rust (<i>Puccinia sorghi</i>); Gray leaf spot (<i>Cercospora zeae-maydis</i>)	160 to 240 mL	<p>Begin all applications prior to disease development. If disease persists or weather conditions are favourable for disease development, apply a second time 10 to 14 days later with a fungicide that contains a different mode of action. Use higher rate and shorter interval when disease pressure is high.</p>

Continued

Crops, Diseases, Rates, and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Chickpea	Ascochyta blight (<i>Ascochyta rabiei</i>)	160 to 240 mL Must be tank-mixed with 0.14 to 0.17 kg per acre <i>Lance</i> WDG	Apply a tank-mix of <i>Headline</i> EC with <i>Lance</i> at the beginning of flowering or the onset of symptoms. Ascochyta blight can develop quickly once established so early detection is essential. DO NOT apply sequential applications of this tank-mix; alternate to a fungicide with a mode of action other than Group 7 or 11 for at least one application.
Lentil	Anthracnose (<i>Colletotrichum truncatum</i>) Ascochyta blight (<i>Ascochyta lentis</i>)	160 mL	Lentil, field pea, dry bean, faba bean and soybean: Apply at the beginning of flowering or at the onset of symptoms for more aggressive diseases* (whichever comes first). If disease persists or weather conditions are favourable for disease development, apply a second application 10 to 14 days later with a fungicide that contains a different mode of action.
Field pea	Mycosphaerella blight (<i>Mycosphaerella</i> , <i>Ascochyta</i> spp.) Powdery mildew (<i>Erysiphe</i> spp.)	160 mL	*Anthracnose in lentils, downy mildew in peas, and Asian soybean rust can develop quickly once can develop quickly once established so early detection is critical.
	Asian soybean rust (<i>Phakopsora pachyrhizi</i>) Downy mildew (<i>Peronospora viciae</i> f.sp. <i>pisi</i>)	160 to 240 mL	
Dry bean (<i>Phaseolus</i> spp.)	Anthracnose (<i>Colletotrichum lindemuthianum</i>) Powdery mildew (<i>Erysiphe</i> spp.) Rust (<i>Uromyces</i> spp.)	160 mL	
	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	160 to 240 mL	
Faba bean	Mycosphaerella blight (<i>Mycosphaerella</i> spp.) (<i>Ascochyta fabae</i>); Powdery mildew (<i>Erysiphe</i>)	160 mL	
	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	160 to 240 mL	
Soybean	Frogeye leaf spot (<i>Cercospora sojina</i>); Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	160 to 240 mL	
Sunflowers	Rust (<i>Puccinia helianthi</i>)	160 mL	Apply for optimum disease suppression prior to disease development. If disease persists or weather conditions are favourable for disease development, apply a second application 10 to 14 days later with a fungicide that contains a different mode of action.
Flax (including low-linolenic acid varieties)	Pasmo (<i>Septoria linicola</i>)	120 to 160 mL	Apply at the mid flower stage (7 to 10 days after the initiation of flowering). If disease persists or weather conditions are favourable for disease development, apply a second application 10 to 14 days later with a fungicide that contains a different mode of action.
Alfalfa (for seed production)	Common leaf spot (<i>Pseudopeziza medicaginis</i>)	160 mL	Apply at the beginning of flowering (10 to 30% bloom) or at the onset of disease.
Bluegrasses; fescues; rye-grasses (for seed production)	Leaf and stem rust (<i>Puccinia recondita</i> ; <i>P. graminis</i>); Powdery mildew – suppression (<i>Erysiphe graminis</i>)	160 to 270 mL	Apply prior to disease development; if disease conditions exist, apply again 12 to 14 days later with a fungicide that contains a different mode of action. Use higher rate and shorter interval when high disease pressure.

Continued

Crops, Diseases, Rates, and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Potato	Early blight (<i>Alternaria solani</i>) (Spray interval: 7 to 14 days) Late blight (<i>Phytophthora infestans</i>) (Spray interval: 7 to 10 days)	180 to 270 mL In tank-mix with <i>Bravo 500</i>	Apply prior to row closure or when conditions become favourable for disease development (whichever comes first). Use higher rates, shorter spray intervals and/or tank-mix with <i>Bravo</i> 500 under heavy disease pressure. DO NOT make more than 1 application in potatoes for late blight and early blight control before before alternating to a fungicide with a different mode of action.

Application Information:

Water Volume:

Ground: Use a minimum water volume of 40 L per acre on oilseeds, cereals, pulses, alfalfa and grasses; use 80 L per acre on potatoes. Ensure thorough coverage of foliage.

Aerial: Use a minimum water volume of 20 L per acre. Ensure thorough coverage of foliage. DO NOT apply more than 160 mL per acre by aerial application.

Pivot or Sprinkler irrigation: DO NOT exceed 0.64 cm (1/4 inch) (63,500 L) per hectare. DO NOT apply registered tank mixes in potato, chickpea, and canola by pivot or sprinkler irrigation. Apply only through overhead sprinkler systems including centre pivot and lateral move containing low pressure drop nozzles.

How it Works:

The active ingredient pyraclostrobin is a strobilurin fungicide with broad spectrum contact and systemic activity. To be used as a preventative application when environmental conditions are favourable for disease development. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Herbicides: *Headline EC* at a rate of 120 to 160 mL per acre can be tank mixed with *Odyssey* on canola quality *Brassica juncea* with Clearfield trait, *Ares*, *Odyssey*, *Odyssey DLX*, and *Tensile* on Clearfield canola, *Liberty Herbicide* (150SN or 200SN) in glufosinate ammonium tolerant canola (eg: LibertyLink canola), registered glyphosate herbicides in glyphosate tolerant canola (eg: Roundup Ready), *Poast Ultra* in canola, and *Equinox EC* in canola and canola quality *Brassica juncea*.

Fungicides: On chickpea, *Headline EC* at a rate of 160 to 240 mL per acre must be applied in tank-mix with 140 to 170 grams per acre *Lance* for control of ascochyta blight. On potatoes, *Headline EC* at rates of 180 to 270 mL per acre may be applied in tank-mix with *Bravo 500* at label rates, additional use recommendations, restrictions, and precautions for the control of late blight. On canola, *Headline EC* can be tank mixed with *Lance Fungicide* at 142 grams per acre at 20 to 50% flowering to control sclerotinia stem rot and suppress black spot.

Restrictions:

Resistance management: Refer to page 310.

Note: BASF does not recommend use of *Headline* alone on potato due to potential for resistance.

Maximum Number of Applications: DO NOT exceed 1 sequential application of this product per season. Any subsequent applications of this product must be in combination with a fungicide that contains a different mode of action. Alfalfa – DO NOT exceed 1 application of this product per season. Canola, rapeseed, canola quality *Brassica juncea*, mustard, flax, dry bean, faba bean, lentil, field pea, chickpea, bluegrass, fescue grass, ryegrass, corn, soybean, sunflower – DO NOT exceed 2 applications of this product per season. Potato – DO NOT exceed 3 applications of this product per season.

Grazing: DO NOT graze treated corn crops within 6 days of last application. DO NOT feed alfalfa hay or forage to livestock. All other crops listed can be grazed or fed to livestock.

Preharvest interval: Barley, rye, wheat, oat – apply no later than the end of flowering, Corn – 7 days, Pulses – 30 days, Forage grasses – 14 days, Alfalfa – not applicable, Oilseeds – 21 days, Potatoes – 3 days.


Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: Crops listed on label may be planted immediately following last application. Wait 14 days before planting all other crops.

Storage: Store in a cool, dry, locked, well-ventilated area without a floor drain. DO NOT freeze.

Environment: Avoid overspray or drift to sensitive habitats. Maintain specified buffer zones. DO NOT spray non-target terrestrial or aquatic habitats.

Hazard Rating:

 Danger Poison – Skin and eye irritant.

For an explanation of the symbols used here see page 11.

Inspire

Fungicide Group – 3

(Refer to page 312)

Company:

Syngenta Canada – PCP #30004

Formulation:

250 g per L difenoconazole formulated as an emulsifiable concentrate.

Container size – 4x1.12 L.

Crops, Diseases, Rates, and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Potato	Early blight (<i>Alternaria solani</i>)	118 to 207 mL	Begin application at first sign of disease or when conditions are conducive for disease development. Repeat application 7 to 14 days after the first application. Maximum 2 consecutive applications before rotating to fungicide with another mode of action. Under severe disease pressure, use higher rate at 7 day interval.

* DO NOT apply more than 826 mL per acre per season. DO NOT apply *Inspire* or other products containing difenoconazole more than one year out of two.

Application Information:

Water Volume:

Ground: Minimum of 60 L per acre.

Aerial: Minimum of 18 L per acre.

Consult nozzle manufacturers for specific nozzle and pressure recommendations. DO NOT apply this product through any type of irrigation system.

How it Works:

The active ingredient difenoconazole is a triazole fungicide with broad spectrum, systemic activity. To be used as a preventative and curative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides: *Inspire* may be mixed with *Revus* (also available as co-pack *Revus Top*). DO NOT make more than 4 applications of this tank mix per season.

Inspire may be mixed with *Bravo 500*.

Restrictions:

Resistance management: Refer to page 310.

Preharvest interval: DO NOT apply within 14 days of harvest.

Re-entry: DO NOT re-enter treated area within 12 hours after application.

Re-cropping: DO NOT plant any other crop for a period of 60 days after application unless *Inspire* is registered for that crop.

Storage: Store in cool, dry place. DO NOT store near food, beverages, or tobacco products.

Environment: This product is toxic to aquatic organisms and certain beneficial insects. Buffer zones of 11 m must be left from the downwind edge of the boom to the sensitive habitats. Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy. Avoid run-off from treated areas into aquatic habitats by including a vegetative strip between the edge of the treated area and the water body.

Hazard Rating:



Warning – causes severe eye damage.

Check label for first-aid information.

For an explanation of the symbols used here see page 11.

Iprodione

Rovral Flo / Rovral RX / Overall 240SC

Fungicide Group – 2

(Refer to page 312)

Company:

FMC of Canada (*Rovral Flo* - PCP#29315)

FMC of Canada (*Rovral RX* - PCP#24378)

MANA Canada (*Overall 240SC* - PCP#30275)

Formulation:

240 g per L iprodione formulated as a liquid flowable.
Container size - 8.4 L.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rates (per acre):	Application Timing
Canola	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	<p>Single application*: 0.85 to 1.25 L</p> <p>Split application* (Rovral Flo/RX only): 0.42 to 0.63 L at 20% bloom; followed by 0.42 L at 50% bloom.</p> <p>Single application for low disease pressure and light crop stands (Rovral Flo/RX only): 0.63 L</p>	Apply at 20 to 50% bloom stage (approximately 4 to 8 days after crop begins to flower). Best protection achieved when applied at the 20 to 30% bloom stage (prior to petal fall). Can be applied until 50% bloom stage (when crop is at its maximum yellow color and prior to significant petal fall).
	Alternaria black spot (<i>Alternaria brassicae</i> and <i>A. raphani</i>) (suppression only)	<p>Single application: 0.85 L</p> <p>Split applications: 0.42 L each application.</p>	<p>Single application: Apply at early green pod stage.</p> <p>Split application: Apply first application at 20 to 50% bloom, followed by second application at early green pod stage.</p>
Alfalfa (grown for seed)	Sclerotinia (<i>Sclerotinia sclerotiorum</i>)	0.85 to 1.25 L *	Make a single application at the 20 to 50% bloom stage.

* Lower rate recommended for most crops; use higher rate for fields with history of severe disease pressure and dense crop stands.

Application Information:

Water Volume: Good coverage of the plants is essential.

Ground: 40 L per acre

Air (canola only): Minimum of 18 L per acre

How it Works:

The active ingredient iprodione is a dicarboximide fungicide with contact activity. To be used as a preventative and eradicator fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 2 applications of this product per season.

Grazing: DO NOT use treated alfalfa for animal feed.

Preharvest interval: 38 days.

Re-entry: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Protect from frost.

Environment: DO NOT apply directly to water. DO NOT contaminate sensitive areas through spray drift, direct application, disposal of waste or cleaning equipment. Observe specified buffer zones.

Hazard Rating:

Caution – Poison.



Warning – Skin and Eye Irritant.

For an explanation of the symbols used here see page 11.

Lance WDG Fungicide

Fungicide Group – 7
(Refer to page 312)

Company:

BASF Canada – PCP# 27495

Formulation:

70% boscalid formulated as a water dispersible granular.
Container size - 2 x 2.83 kg per case.

Crops, Diseases, Rates and Timing:

(Ground, Aerial, and Pivot or Sprinkler Irrigation Applications)

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Alfalfa (seed production only)	Blossom blight (<i>Sclerotinia sclerotiorum</i>)/ <i>Botrytis cinerea</i> ; common leaf spot (<i>Pseudopeziza medicaginis</i>); Spring black stem (<i>Phoma medicaginis</i>); leaf spot (<i>Leptosphaerulina briosiani</i>)	170 g	Apply at 20 to 50% flowering. Apply every 7 to 14 days if disease persists, or weather conditions are favourable for disease development.
Canola and Mustard (oilseed and condiment)	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	142 g	Apply at 20 to 50% flowering. Apply a second time 7 to 14 days later up to full bloom if disease persists, or weather conditions are favourable for disease development.
	Black spot (<i>Alternaria brassicae</i> and <i>A. raphani</i>)	142 g	Apply at late flowering to early green pod.
Dry beans (including field bean and faba bean)	White mould (<i>Sclerotinia sclerotiorum</i>)	227 to 312 g	Apply at 20 to 50% flowering. Apply a second time 7 to 14 days later if disease persists, or weather conditions are favourable for disease development. Use the higher rate to obtain extended protection and maximum yield benefit.

Crops, Diseases Controlled, Rates and Timing *continued*:

(Ground and Aerial Applications)

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Chickpea** Lentil	Ascochyta blight (<i>Ascochyta</i> spp.), white mould (<i>Sclerotinia sclerotiorum</i>), gray mould (<i>Botrytis cinerea</i>)	170 g	Apply at the beginning of flowering. Apply a second time 7 to 14 days later if disease persists, or weather conditions are favourable for disease development.
Sunflower	Suppression of sclerotinia head rot (<i>Sclerotinia sclerotiorum</i>), leaf spot (<i>Alternaria helianthi</i>)	142 to 259 g	Apply at early flower for optimal disease suppression. Use the higher rate when disease pressure is high or there is a history of high disease in the field.

** For the control of ascochyta blight in chickpea, *Lance* should be mixed with 160 to 240 mL per acre *Headline EC*.

(Ground Application Only)

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Pea***	Ascochyta blight (<i>Ascochyta</i> spp.); mycosphaerella blight (<i>Mycosphaerella</i> spp.); gray mould (<i>Botrytis cinerea</i>)	170 g	Apply at the beginning of flowering. Apply a second time 7 to 14 days later if disease persists, or weather conditions are favourable for disease development.

*** DO NOT apply by air

Application Information:

Water Volume:

Aerial (registered for all crops but peas): Use a minimum water volume of 16 L per acre and ensure thorough coverage of foliage.

Ground: Use a minimum water volume of 40 L per acre and ensure thorough coverage of foliage.

Pivot and Sprinkler Irrigation: DO NOT exceed 0.64 cm (1/4 inch) or 25,700 L per acre. Apply only through sprinkler systems including centre pivot, lateral move, end two, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems.

How it Works:

The active ingredient boscalid is a carboxamide (SDHI) fungicide with systemic activity. To be used as a preventative application when environmental conditions are favourable for disease development. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Insecticides: For control of corn borer in succulent beans, *Lance* can be tank-mixed with *Matador 120 EC* at 37 mL per acre.

Fungicides: For the control of ascochyta blight on chickpea, *Lance* at rate of 140 to 170 g per acre should be applied with 160 to 240 mL per acre *Headline EC*.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Sunflower – DO NOT exceed 1 application of this product per season. Canola, mustard, dry bean, chickpea, lentil, pea – DO NOT exceed 2 applications of this product per season. Alfalfa – DO NOT exceed 3 applications of this product per season.

Grazing: All crops except alfalfa (grown for seed) can be grazed or fed to livestock.

Preharvest interval: Beans, canola, chickpea, lentil, pea - 21 days; alfalfa - not applicable.

Re-entry: DO NOT re-enter treated area for 12 hours after application or until dry.

Re-cropping: A plant back restriction of 14 days is required for all crops not on the label.

Storage: Store in a cool, dry, locked, well-ventilated area without a floor drain.

Environment: DO NOT apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.

Hazard Rating:



Caution Poison – Potential Skin Sensitizer.



Warning – Eye Irritant.

For an explanation of the symbols used here see page 11.

Luna Tranquility

Fungicide Group – 7, 9
(Refer to page 312)

Company:

Bayer CropScience – PCP#30510

Formulation:

125 g per L fluopyram + 375 g per L pyrimethanil formulated as a suspension concentrate.

Container size – 2 x 4.86 L.

Crops, Diseases, Rates, and Timing:

Crop:	Diseases:	Application Rate (per acre):	Application Timing:
Potato	Early blight (<i>Alternaria solani</i>)	243 mL	Begin fungicide applications preventatively. Continue as needed on a 7 to 14 day interval. When disease pressure is severe, use the shorter intervals.
	Brown leaf spot (<i>Alternaria alternata</i>)		
	White mold (<i>Sclerotinia sclerotiorum</i>)	324 mL	
	Black dot – suppression only (<i>Colletotrichum coccodes</i>)		

Application Information:

Ground: Use a minimum water volume of 80 L per acre and ensure thorough coverage of foliage.

Aerial: Use a minimum water volume of 20 L per acre and ensure thorough coverage of foliage.

How it Works:

The active ingredient fluopyram is a carboxamide fungicide with systemic activity. The active ingredient pyrimethanil is a anilinopyrimidine fungicide with contact and systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT apply more than 1.3 L per acre of this product per season.

Grazing: No restriction listed.

Pre-harvest interval: 7 days.

Re-entry: DO NOT re-enter treated areas until 12 hours after application.

Re-cropping: A plant back restriction of 30 days is required for canola, cereal grains, corn, soybean, dry bean, chickpea, lentil, and alfalfa.

Storage: Do not store below freezing. If stored for one year or longer, shake well before using. Store the tightly closed container away from feeds, seeds, fertilizer, plants and foodstuffs. Keep the product in the original container during storage.

Environment: Toxic to aquatic organisms and birds. DO NOT apply directly to water, to areas where surface water is present, or to intertidal areas below the high water mark. non-target areas. Observe buffer zones outlined in the label.

Hazard Rating:

None listed.

Mancozeb

Fungicide Group – M3

(Refer to page 312)

Dithane Rainshield/Manzate Pro-Stick/Penncozeb 75 DF

Company:

Dow AgroSciences (*Dithane Rainshield* – PCP# 20553)

United Phosphorus Inc; (*Manzate Pro-Stick* – PCP# 28217)

United Phosphorus Inc. distributed by United Agri Products (*Penncozeb 75 DF* – PCP# 25397)

Formulations:

Dithane Rainshield - 75% mancozeb formulated as a water dispersible granule. Container size - 3.5 to 544 kg.

Manzate Pro-Stick - 75% mancozeb formulated as a dry flowable. Container sizes - 10 to 20 kg.

Penncozeb 75 DF - 75% mancozeb formulated as a wettable granule. Container sizes - 2.5 to 250 kg.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Potato	Early blight (<i>Alternaria solani</i>), Late blight (<i>Phytophthora infestans</i>)	0.45 to 0.90 kg	Apply at 0.45 kg per acre rate when plants are 10 to 15 cm high; increase to 0.7 kg per acre as plants increase in size, and 0.9 kg per acre at row closure. Apply every 7 to 10 days. Spray interval may be reduced to 5 to 6 days during periods of wet weather favouring late blight and / or vigorous crop growth.
Wheat	Tanspot (<i>Pyrenophora tritici-repentis</i>), Leaf rust (<i>Puccinia recondita</i>), Septoria leaf blotch (<i>Septoria</i> sp)	0.45 to 0.9 kg	Apply 0.45 kg per acre early (when crop is in the 3-leaf to tillering stage); apply 0.9 kg per acre later (when head is fully emerged, but prior to flowering).
Lentil	Anthrachnose (<i>Colletotrichum truncatum</i>) Ascochyta blight (<i>Ascochyta lentis</i>)	0.9 kg	Apply first application before flower when bud formation is evident; apply second application 10 to 14 days later at early to mid-bloom but prior to row closure. A third application may be applied 10 to 14 days later.
Alfalfa (for seed)	Leaf and stem spot diseases (<i>Pseudopeziza medicaginis</i>)	0.6 kg	Apply first application prior to 50% bloom; apply second application 7 to 10 days later; apply third application 10 days after second.

Application Information:

Water Volume:

Thorough uniform coverage is essential for good disease control.

Ground: 40 L per acre (wheat); 40-80 L per acre (lentil).

Air: 16 L per acre (wheat, lentil, potato).

How it Works:

The active ingredient mancozeb is a dithiocarbamate fungicide with multi-site contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides (potato only): For early and late blight control, *Manzate Pro-stick* can be tank-mixed with *Kocide 2000*; and for control of late blight, with *Curzate 60 DF*.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Wheat – DO NOT exceed 2 applications of this product per season. Lentil, alfalfa – DO NOT exceed 3 applications of this product per season.

Grazing: DO NOT graze or feed treated crop or straw to livestock. DO NOT graze or cut treated alfalfa for hay.

Preharvest interval: Potato - 1 day; Lentil - 35 days; Wheat - 40 days

Re-entry: DO NOT re-enter treated areas within 24 hours of application.

Re-cropping: No restrictions listed.

Storage: Store in cool, dry, well-ventilated place. Keep away from fire and sparks.

Environment: Toxic to aquatic organisms. DO NOT con-

taminate any body of water by direct application, drift or by cleaning equipment.

Hazard Rating:



Warning - Poison.



Danger - Eye irritant.

For an explanation of the symbols used here see page 11.

Phostrol

Fungicide Group - 33
(Refer to page 312)

Company:

Engage Agro Corporation - PCP#30449

Formulation:

53.6% mono- and dibasic sodium, potassium, and ammonium phosphites formulated as a liquid flowable.

Container size - 2 x 10 L and 1000 L

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (L per acre)	Application Timing:
Potato	Suppression of pink rot (<i>Phytophthora erythroseptica</i>)	2.3 to 4.7	In furrow: Apply in a band at planting directly over the seed pieces prior to row closure.
	Late blight (<i>Phytophthora infestans</i>)	1.2 to 4.7	Foliar applications: For preventative control of late blight and preventative suppression of pink rot begin applications when conditions favouring disease development exist and continue on a 7 to 14 day interval.
	Suppression of pink rot (<i>Phytophthora erythroseptica</i>)	2.3 to 4.7	
	Late blight (<i>Phytophthora infestans</i>) Pink rot (<i>Phytophthora erythroseptica</i>)	0.17 to 0.81	Post harvest control: Apply directly to the tubers and ensure complete and even coverage.

Application Information:

Water Volume:

Ground: Minimum of 12 L per acre for in-furrow treatment and minimum of 81 L per acre for foliar applications.

Aerial: Minimum of 20 L per acre.

How it Works:

The active ingredient mono- and dibasic sodium, potassium, and ammonium phosphite is a phosphonates fungicide with systemic activity to suppress pathogen inoculum. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Pink rot: *Ridomil Gold 480SL* (in furrow), *Ridomil Gold MZ* and *Ridomil Gold Bravo Twin Pack* (foliar)

Late Blight: May be tank mixed with one of the following fungicides: *Bravo 500*, *Bravo ZN*, *Echo 720*, *Echo 90DF*, *Ridomil Gold Bravo Twin Pack*, *Dithane Rainshield*, *Manzate Pro-Stick*, *Gavel 75DF*, *Penncozeb 75DF*

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Do NOT exceed 7 applications of this product per season

Grazing: Do NOT graze treated fields or feed treated forage to livestock.

Preharvest interval: May be applied up to the day of harvest and post harvest.

Re-entry: Re-entry interval after application is 12 hours.

Re-cropping: No restriction listed.

Storage: Store in original tightly closed container.

Environment: Avoid run-off from treated areas into aquatic areas.

Toxicity: Toxic to aquatic organisms, non-target terrestrial plants and small wild animals.

Hazard Rating:



Caution – Skin irritant.

For an explanation of the symbols used here see page 11.

Priaxor DS

Fungicide Group – 7, 11
(Refer to page 312)

Company:

BASF Canada – PCP# 30646

Formulation:

250 g per L fluxapyroxad and 250 g per L pyraclostrobin formulated as a suspension concentrate.

Container size – Case (2 x 9.6 L); 192 L drum.

Crops, Diseases, Rates and Timing:

Crop:	Diseases:	Application Rate (per acre)	Application Timing:
Chickpea	Ascochyta blight (<i>Ascochyta rubiei</i>)	160 mL	Apply at beginning of flowering or at the onset of disease symptoms. If disease persists or weather conditions are favourable for disease development, make a second application 10 to 14 days later with a fungicide that contains a different mode of action.
	White mould suppression (<i>Sclerotinia sclerotiorum</i>)	120 to 160 mL	
Field Pea	Mycosphaerella blight (<i>Mycosphaerella pinodes</i>)	120 to 160 mL	
	Powdery mildew (<i>Erysiphe pisi</i>), Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	160 mL	
	White mould suppression (<i>Sclerotinia sclerotiorum</i>)	120 to 160 mL	
Lentil	Ascochyta blight (<i>Ascochyta lentis</i>) and suppression of white mould (<i>Sclerotinia sclerotiorum</i>)	120 to 160 mL	
	Anthraxnose (<i>Colletotrichum truncatum</i>)	120 to 160 mL	

*Use of a non-ionic surfactant (0.125% v/v) is recommended but not required.

Application Information:

Water Volume:

Ground: Minimum of 40 L per acre.

Aerial: Minimum of 20 L per acre.

Consult nozzle manufacturers for specific nozzle and pressure recommendations.

How it Works:

The active ingredient fluxapyroxad is a carboximide (SDHI) fungicide with systemic activity. The active ingredient pyraclostrobin is a member of the strobilurin class of chemistry used as a broad spectrum fungicide. To be used as a preventative application when environmental conditions are favourable for disease development. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT make sequential applications of this product before alternating to at least

one application of a fungicide with a different mode of action. DO NOT exceed 2 applications of this product per season.

Grazing: All crops can be grazed or fed to livestock.

Preharvest interval: DO NOT apply within 30 days of harvest.

Re-entry: DO NOT re-enter treated areas within 12 hours after application.

Re-cropping: Tuberous and corm vegetables, sugar beets, legume vegetables, pome fruits, stone fruits, cereals and oilseeds may be planted immediately following the last application. A plant-back interval of one year is required for all other crops.

Storage: Store in original tightly closed container. Protect from freezing. Store away from food or feed.

Environment: Toxic to aquatic organisms, small mammals and non-target terrestrial plants. Maintain specified buffer zones. The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable and/or depth of the water table is shallow.

Hazard Rating:



Danger – Eye irritant.

Check label for first-aid information

For an explanation of the symbols used here see page 11.

Proline 480 SC

Fungicide Group – 3
(Refer to page 312)

Company:

Bayer CropScience - PCP#28359

Formulations:

480 g per L prothioconazole formulated as a suspension concentrate. Container size - 5.1 litre.

Crops, Diseases, Rates, and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Barley; Wheat	Suppression of fusarium head blight (<i>Fusarium</i> spp.)	126 to 168 ¹ mL	<p>Timing of application is critical:</p> <p>Barley: Apply within the time period when 70 to 100% of barley heads on the main stem are fully emerged to 3 days after full head emergence.</p> <p>Wheat: Apply within the time period from when at least 75% of wheat heads on the main stem are fully emerged to when 50% of heads on the main stem are in flower.</p> <p>Use higher rate when disease pressure is expected to be high or to provide the highest level of myco-toxin reduction.</p> <p>Spray coverage is essential for optimum efficacy: Spray equipment must be set to provide good coverage to heads. Use forward and backward mounted nozzles, or nozzles that have a two-directional spray.</p>
Wheat	Glume blotch (<i>Stagonospora nodorum</i>)	168 ¹ mL	
	Speckled leaf blotch (<i>Septoria tritici</i>); tan spot (<i>Pyrenophora tritici-repentis</i>); leaf rust (<i>Puccinia recondita</i>)	126 ¹ mL	
Oat	Crown rust (<i>Puccinia coronata</i>)	126 ¹ mL	
Barley	Net blotch (<i>Pyrenophora teres</i>); scald (<i>Rhynchosporium secalis</i>); spot blotch (<i>Cochliobolus sativus</i>)	84 to 126 mL	
Corn	Suppression of <i>Fusarium</i> and <i>Gibberella</i> ear rots (<i>Fusarium</i> spp. and <i>Gibberella</i> spp.); rusts (<i>Puccinia sorghi</i> , <i>Puccinia polysora</i>), eyespot (<i>Kabatiella zeae</i> or <i>Aureobasidium zeae</i>), grey leaf spot (<i>Cercospora zeae-maydis</i>), and northern blight (<i>Setosphaeria turcica</i>)	168 ¹ mL	Apply from silking (tip of stigmata visible) to silk browning (stigmata drying).
Canola; Rapeseed; Oriental mustard	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	126 to 147 ² mL	Apply at 20 to 50% bloom stage (prior to petal fall). Use high rate if history of heavy disease or if dense crop stand.

Continued

Crops, Diseases, Rates, and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Chickpea	Ascochyta blight (<i>Ascochyta rubiei</i>)	126 to 168 ² mL	Apply at first sign of disease. Repeat applications every 10 to 14 days. Use high rate when conditions favour disease or when growing susceptible varieties.
Lentil	Ascochyta blight (<i>Ascochyta lentis</i>)	126 to 168 ² mL	Apply at the first sign of disease. Repeat application 10 to 14 days later. Use high rate when conditions favour disease or when growing susceptible varieties.
Flax (linseed) and Crambe	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	128 to 149 ² mL	Apply when the crop is in the 20 to 50% bloom stage. Best protection will be achieved when the fungicide is applied prior to petals beginning to fall, allowing the maximum number of petals to be protected. Use high rate in fields with a history of heavy disease pressure or for dense crop stands.
Soybean	Asian soybean rust (<i>Phakopsora pachyrhizi</i>) and frogeye leaf spot (<i>Cercospora sojina</i>).	84 ² mL	Apply when first symptoms of disease can be found or that the risk of infection is imminent.

¹ Apply with non-ionic surfactant, i.e. AgSurf or Agral 90 at 0.125% v/v.

² May be applied with the lowest rate of non-ionic surfactant, i.e. AgSurf or Agral 90.

Application Information:

DO NOT apply during periods of dead calm or when winds are gusty. Ensure uniform coverage.

Water volumes:

Ground (all crops): Minimum of 40 L per acre.

Aerial (canola only): Minimum of 20 L per acre. Follow detailed label recommendations for aerial application.

How it Works:

The active ingredient prothioconazole is a triazole fungicide with broad-spectrum systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Bayer CropScience supports the following mixes that are not on the Proline 480 SC label. Apply mixes according to the most restrictive use limitations for either product:

Insecticides: Decis, Lorsban, Matador, Sevin XLR

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Corn, flax, soybean – DO NOT exceed 1 application of this product per season. Wheat, barley, oat, canola, lentil – DO NOT exceed 2 applications of this product per season. Chickpea – DO NOT exceed 3 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 30 days (barley, wheat, oat); 36 days (canola, flax); 7 days (chickpea, lentil); 14 days (corn); 20 days (soybean).

Re-entry: DO NOT re-enter treated fields within 24 hours of application.

Re-cropping: May be re-planted with any crop specified on the label as soon as practical. For crops not listed, wait 30 days.

Storage: DO NOT store at temperatures below freezing. Keep in original tightly closed container and store away from feeds, seeds, fertilizer, plants and food stuffs. Keep away from sources of heat. Shake well before using if stored for more than 1 year.

Environment: Toxic to aquatic organisms. DO NOT apply directly to freshwater, estuaries or marine habitats. DO NOT contaminate bodies of water by cleaning of equipment or disposal of wastes. Observe the specified buffer zones.

Hazard Rating:



Caution - Poison.

For an explanation of the symbols used here see page 11.

Propiconazole

Fungicide Group - 3

(Refer to page 312)

Tilt 250E/Bumper 418 EC/Pivot 418 EC/Propel EC/Propel/Nufarm Propiconazole Fungicide

Company:

Syngenta Canada (*Tilt 250E* - PCP#19346)

MANA Canada (*Bumper 418 EC* - PCP#28017)

Interprovincial Cooperative Ltd. (*Pivot 418 EC* - PCP#28219)

Loveland Products (formerly Viterra) (*Propel* - PCP#29548)

Nufarm Agriculture Inc (*Nufarm Propiconazole Fungicide* - PCP#30367)

Formulations:

Tilt 250E - 250 g per L propiconazole formulated as an emulsifiable concentrate. Container sizes - 2 x 8L and 4 x 5 L.

Bumper 418 EC - 418 g per L propiconazole formulated as an emulsifiable concentrate. Container size - 4.8 L.

Pivot 418 EC - 418 g per L propiconazole formulated as an emulsifiable concentrate. Container size - 2 x 4.8L.

Propel - 250 g per L propiconazole formulated as an emulsifiable concentrate. Container size - 8L.

Nufarm Propiconazole Fungicide - 418 g per L propiconazole formulated as an emulsifiable concentrate. Container size - 4.8L.

Foliar Fungicides

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rates (per acre):		Application Timing:
		<i>Tilt 250E</i> <i>Propel</i>	<i>Bumper 418 EC</i> <i>Pivot 418 EC</i> <i>Nufarm Propiconazole Fungicide</i>	
Wheat (durum, winter wheat, spring wheat)	Septoria leaf spot (<i>Septoria</i> sp.) and tan spot (<i>Pyrenophora tritici-repentis</i>) **	100 mL	60 mL	An early application of the low product rate can be timed with herbicide application. At growth stage (G.S.) 12 to 23, 100 ml per acre (<i>Tilt</i> or <i>Propel</i>) or 60 ml per acre (<i>Bumper</i>) may be applied for early season disease suppression under normal field conditions. At this stage, if there is a history of high disease pressure in the field and/or if field conditions favour disease development use the full-rate fungicide application (200 ml per acre for <i>Tilt</i> or <i>Propel</i> ; 120 ml per acre for <i>Bumper</i>). This rate can also be used to suppress disease at later growth stages (G.S. 29-55).
Barley	Net blotch (<i>Pyrenophora teres</i>) **			

Continued

Crops, Diseases, Rates and Timing *continued*:

Crop:	Diseases Controlled:	Application Rates (per acre):		Application Timing:
		Tilt 250E Propel	Bumper 418 EC Pivot 418 EC Nufarm Propicon- azole Fungicide	
Wheat (durum, winter wheat, spring wheat)	Septoria leaf spot (<i>Septoria</i> sp.), tan spot (<i>Pyrenophora tritici-repentis</i>), septoria glume blotch, stripe rust (<i>Puccinia striiformis</i>), leaf rust (<i>Puccinia tritica</i>), stem rust (<i>Puccinia graminis</i>) and powdery mildew (<i>Erysiphe graminis</i> f. sp. <i>tritici</i>)	200 mL	120 mL	Application may be made at the very early stage of disease (during tillering or stem elongation). The product lasts about 3 weeks in the plant, so another application will be necessary if disease pressure continues. Make the 2nd application up to when the head is half-emerged.
Barley	Spot blotch (<i>Cochliobolus sativus</i>), net blotch (<i>Pyrenophora teres</i>), scald (<i>Rhynchosporium secalis</i>), leaf rust (<i>Puccinia hordei</i>), stem rust (<i>Puccinia graminis</i>), septoria leaf spot (<i>Septoria</i> sp.) and powdery mildew (<i>Blumeria graminis</i>)			
Oat	Septoria leaf blotch (<i>Septoria avenae</i>) and crown rust (<i>Puccinia coronata</i>)			
Canaryseed*	Septoria leaf mottle (<i>Septoria triseti</i>)**	200 mL	120 mL	Apply only once at flag leaf emergence.
Corn	Rust (<i>Puccinia sorghi</i>)	200 mL	120 mL	Apply when rust pustules first appear; make 2nd application 14 days later.
	Northern leaf blight (<i>Setosphaeria turcicum</i>)	100 to 200 mL	60 to 120 mL	Apply when disease first appears. Use low rate if disease pressure is low.
Canola	Blackleg (<i>Leptosphaeria maculans</i>)	200 mL	120 mL	Apply during the rosette stage (after second true leaf and prior to bolting).
Bean	Rust (<i>Uromyces</i> spp.)	200 mL	120 mL	Apply when disease first appears; make 2nd application 14 to 21 days later.
Soybean* (grown for seed)	Frogeye leaf spot (<i>Cercospora soja</i>)	200 to 300 mL	120 to 182 mL	Apply when disease first appears; make 2nd application 14 days later if severe disease pressure exists.

Continued

Crops, Diseases, Rates and Timing *continued*:

Crop:	Diseases Controlled:	Application Rates (per acre):		Application Timing:
		<i>Tilt 250E</i> <i>Propel</i>	<i>Bumper</i> 418 EC <i>Pivot 418</i> EC <i>Nufarm</i> <i>Propiconazole</i> <i>Fungicide</i>	
Legume*** (soybean, bean, lentil, pea, chickpea)	Asian soybean rust (<i>Phakopsora pachyrhizi</i>)	200 to 300 mL	-	Apply at the first sign of rust; make 2nd application 14 days later if disease con- tinues.
	Powdery mildew (<i>Microsphaera diffusa</i> , <i>Erysiphe pisi</i> , <i>E. polygoni</i>)	200 mL	-	
Timothy *†	Purple eyespot (<i>Cladosporium phlei</i>)	200 mL	120 mL	Apply at the first sign of disease (usu- ally at the beginning of flowering). Can be applied up to full flowering. Spray interval of 14 days.

* Ground application only.

** Suppression only.

*** Only *TILT 250E* and *Propel* are registered for use on these crops.

† Only *TILT 250E*, *Pivot 418 EC* and *Propel* are registered for use on this crop.

Application Information:

Water Volume:

Ground: Minimum 80 L per acre

Aerial: 16 to 20 L per acre

How it Works:

The active ingredient propiconazole is a triazole fungicide with broad spectrum systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Herbicides: In wheat and barley only, propiconazole may be tank-mixed with one of the following: 2,4-D amine, Estemine 2,4-D, MCPA amine, Estemine MCPA, Buctril-M or Pardner and in wheat only, may be applied with *Horizon 240 EC*. In wheat and barley only, *Pivot 418 EC* may be tank-mixed with *Logic M* or *Brotex 240*; *Pivot 418 EC*, *Bumper 418 EC*, or *Nufarm Propiconazole Fungicide* may be tank-mixed with *Badge* or *Bromotril 240 EC*. In spring wheat and barley only, *Tilt 250E* or *Propel* may be tank-mixed with *Axial 100 EC*. Refer to labels for tank-mix precautions.

Fertilizers: Propiconazole may be applied with up to 4 kg per acre (9 lb per acre) of actual nitrogen. The appropriate amount of urea can be dissolved in water and added to the spray tank before adding the fungicide. Excessive nitrogen or application during hot weather may result in crop injury. DO NOT add nitrogen when tank-mixing propiconazole with a herbicide.

Insecticides: In field corn, propiconazole can be tank-mixed with one of the following: *Matazor 120 EC/Silencer 120 EC* or *Ripcord*. In legumes, *Tilt 250E* or *Propel* can be tank-mixed with *Matazor 120 EC*.

Fungicides: For control of Asian soybean rust in legumes, *Tilt 250E* or *Propel* can be tank-mixed with *Quadris*. *Tilt 250E* or *Propel* may be tank-mixed with *Quadris* in wheat and barley. Consult each label for diseases controlled and appropriate timing.

Note: Syngenta supports the following mixes that are not on the respective labels. Apply mixes according to the most restrictive use limitations for either product:

Propel Tank Mixes:

Herbicides: *Axial*, *Broadband*, *Horizon NG*, *Touchdown Total*, *Truxion*

Tilt Tank Mixes:

Herbicides: *Liberty*, *Sierra 2.0*, *Touchdown Total*, *Truxion*, a ½ rate of *Tilt* may also be tank mixed with *Broadband*.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Wheat, barley, corn, bean, legume, timothy – DO NOT exceed 2 applications of this product per season.

Grazing: DO NOT graze animals on treated green crops within 3 days of application. DO NOT feed straw treated with herbicide tank mixes to livestock. DO NOT use treated soybean seed for animal feed.

Preharvest interval: Wheat, oats, barley - 45 days. Canola - 60 days. Corn - 14 days if tank-mixed with an insecticide. Soybeans - 50 days. Beans - 28 days. Timothy - 14 days.

Re-entry: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: DO NOT freeze. Store products away from food or feed.

Environment: Toxic to aquatic organisms. DO NOT contaminate any body of water by direct application, drift or by cleaning equipment.

Hazard Rating:



Warning - Poison (*Bumper 418 EC, Pivot 418 EC, Nufarm Propiconazole Fungicide*).



Caution - Poison (*Tilt 250 EC, Propel*).



Warning - Eye and Skin Irritant.
Potential Skin Sensitizer.

For an explanation of the symbols used here see page 11.

Propulse

Fungicide Group - 3, 7
(Refer to page 312)

Company:

Bayer CropScience - PCP#30511

Formulation:

200 g per L prothioconazole + 200 g per L fluopyram formulated as a suspension concentrate.

Container size - 6.1 litres.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre)*:	Application Timing:
Chickpea	White mould (<i>Sclerotinia sclerotiorum</i>)	304 mL	Begin fungicide applications preventatively. When disease pressure is high or when agronomic or weather conditions are conducive to disease development, continue applications as needed, on a 10 to 14 day interval. Use the higher rate when conditions for heavy infestation exist and/or growing less resistant cultivars.
	Ascochyta blight (<i>Ascochyta rabiei</i>)	202 to 304 mL	
Dry bean	White mould (<i>Sclerotinia sclerotiorum</i>)	304 mL	Begin fungicide applications preventatively. When disease pressure is high or when agronomic or weather conditions are conducive to disease development, continue applications as needed on a 7 to 14 day interval. Use shorter intervals for best protection.
Lentil	White mould (<i>Sclerotinia sclerotiorum</i>)	304 mL	Begin fungicide applications preventatively. When disease pressure is high or when agronomic or weather conditions are conducive to disease development, continue applications as needed, on a 10 to 14 day interval. Use the higher rate when conditions for heavy infestation exist.
	Ascochyta blight (<i>Ascochyta lentis</i>)	202 to 304 mL	

Application Information:

Water Volume:

Ground: Minimum of 40 L per acre.

Use sufficient water and spray pressure to provide thorough and uniform coverage of plants. DO NOT apply

under periods of dead calm. Avoid application of this product when winds are gusty.

How it Works:

The active ingredient prothioconazole is a triazole fungicide with broad-spectrum systemic activity. The active ingredient fluopyram is a carboxamide fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310. Make no more than 2 sequential applications before switching to a fungicide with a different mode of action.

Maximum number of applications: DO NOT apply more than 607 mL per acre per season.

Grazing: DO NOT graze treated area, and do not harvest for forage or hay.

Preharvest interval: DO NOT apply within 14 days of harvest.

Re-entry: DO NOT re-enter treated areas until 24 hours after application.

Re-cropping: Dry beans, chickpeas, lentils and peanuts may be replanted immediately following the last application. Following dry beans, the following crops may be planted 30 days after the last application: Alfalfa, canola, cereal grains, corn, potato and soybean. Do not rotate to any other crops.

Aerial Application: DO NOT apply by air.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Do not store below freezing. If stored for one year or longer, shake well before using.

Environment: This product is toxic to birds and aquatic organisms. When using *Propulse* consult the product label for buffer zones.

Hazard Rating:

None listed.

Prosaro 250 EC

Fungicide Group – 3
(Refer to page 312)

Foliar Fungicides

Company:

Bayer CropScience - PCP#29821

Formulation:

125 g per L prothioconazole + 125 g per L tebuconazole, formulated as an emulsifiable concentrate.

Container size – 6.5 litres; 104 L tote.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Barley; Wheat	Suppression of fusarium head blight (<i>Fusarium</i> spp.)	324 mL	<p>Barley: Apply as a preventative spray within the time period when 70 to 100% of barley heads on the main stem are fully emerged to 3 days after full head emergence.</p> <p>Wheat: Apply as a preventative spray within the time period from when at least 75% of wheat heads on the main stem are fully emerged to when 50% of heads on the main stem are in flower.</p>
Barley	Net blotch (<i>Pyrenophora teres</i>); scald (<i>Rhynchosporium secalis</i>); spot blotch (<i>Cochliobolus sativus</i>), leaf blotch (<i>Septoria passerinii</i>), leaf, stem, and stripe rusts (<i>Puccinia hordei</i> , <i>P. graminis</i> , <i>P. striiformis</i>), powdery mildew (<i>Erysiphe graminis</i>)		
Wheat	Leaf and glume blotch (<i>Septoria tritici</i> , <i>Stagonospora nodorum</i>), tan spot (<i>Pyrenophora tritici-repentis</i>); leaf, stem, and stripe rusts (<i>Puccinia recondita</i> , <i>P. graminis</i> , <i>P. striiformis</i>), powdery mildew (<i>Erysiphe graminis</i>)		

Application Information:

DO NOT apply during periods of dead calm or when winds are gusty. Ensure uniform coverage.

Water Volume:

Ground: Minimum of 40 L per acre.

Aerial: Minimum of 20 L per acre. Follow detailed label recommendations for aerial application.

How it Works:

The active ingredients prothioconazole and tebuconazole are demethylation inhibitors with broad-spectrum systemic activity. To be used as a preventative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes

None registered.

Bayer CropScience supports the following mixes that are not on the *Prosaro 250 EC* label. Apply mixes according to the most restrictive use limitations for either product:

Insecticides: *Lorsban*

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 1 application of this product per season.

Grazing: DO NOT allow livestock to graze or feed green forage to livestock prior to 6 days after treatment with *Prosaro 250 EC* fungicide. Straw cut after harvest may be fed or used for bedding.

Preharvest interval: 36 days.

Re-entry: DO NOT re-enter treated fields until 12 hours post-application.

Re-cropping: Treated areas may be replanted with any crop specified on the label as soon as practical after application. For oat and soybean, DO NOT plant back within 30 days of application. For all other crops, DO NOT plant back until 120 days after application. Tebuconazole is persistent and will carryover. It is recommended that any products containing tebuconazole not be used in areas treated with this product during the previous season.

Storage: DO NOT store in or around the home. DO NOT store at temperatures below freezing. Keep in original tightly closed container and store away from feeds, seeds, fertilizer, plants and food stuffs. Keep away from sources of heat. Shake well before using if stored for more than 1 year. DO NOT contaminate water, food, or feed by storage or disposal.

Environmental: Toxic to birds, small wild animals, aquatic organisms, and non-target plants. As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT apply to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff is hazardous to aquatic organisms in neighbouring areas. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body. Follow buffer zones as per the product label.

Hazard Rating:



Danger Eye irritant.



Caution – Skin irritant

For an explanation of the symbols used here see page 11.

Quadris (2013)

See Azoxystrobin on page 324.

Quadris Top

Fungicide Group – 3, 11
(Refer to page 312)

Company:

Syngenta Canada – PCP#30518

Formulation:

200 g per L azoxystrobin and 125 g per L difenconazole formulated as a flowable suspension concentrate.

Container size – 2 x 10.125 L.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre)*:	Application Timing:
Potato	Early Blight (<i>Alternaria solani</i>)	229 to 405	Apply as a broadcast foliar spray in sufficient water (60 L/acre) for thorough coverage. Apply on a 7 to 14 day interval, starting prior to disease establishment.
	Suppression of brown spot (<i>Alternaria alternata</i>) and black dot (<i>Colletotrichum coccodes</i>)	229 to 405	Apply as a broadcast foliar spray in sufficient water (60 L/acre) for thorough coverage. Apply prior to disease. Apply no more than 1 application to target these diseases. If disease pressure is high, use the highest rate.

Application Information:

Water Volume:

Ground: Use sufficient water volume to obtain adequate coverage. Use minimum 60 L per acre.

Aerial: Use sufficient water volume to obtain adequate coverage. Use minimum 60 L per acre.

How it Works:

The active ingredient azoxystrobin belongs to a strobilurin group of fungicides and difenconazole is a triazole fungicide. Together they provide broad spectrum preventative and systematic. To be used as a preventative and curative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed three applications per season.

Grazing: No restrictions listed.

Preharvest interval: 14 days.


Re-entry: DO NOT re-enter treated areas until 12 hours after application.

Re-cropping: Do not plant any other crop for a period of 60 days following application to the preceding crop unless *Quadris Top* or *Inspire* are registered for that crop.

Storage: Store in cool, dry place. Do not store food, beverages or tobacco products in storage area.

Environment: This product is toxic to aquatic organisms (or invertebrates), fish and mammals. Observe buffer zones outlined in the label.

Hazard Rating:

 Caution – Eye irritant.

For an explanation of the symbols used here see page 11.

Quash

Fungicide Group – 3

(Refer to page 312)

Company:

Valent Canada – PCP#30402

Formulation:

50.0% metconazole formulated as water dispersible granules. Container sizes – Pouch (2 x 280g)

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Canola	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	113 g	Make a single, preventative application between 20% and 50% bloom.
Dry bean	Suppression of white mould (<i>Sclerotinia sclerotiorum</i>)	113 g	Apply prior to disease development. Make first application at 20 to 50% bloom stage, before disease symptoms are visible. Make second application at full bloom. Do not make the second application until 9 days have passed since the first application.
Field pea	Powdery mildew (<i>Erysiphe pisi</i>), suppression of white mould (<i>Sclerotinia sclerotiorum</i>)		
Chickpea Lentil	Suppression of Ascochyta blight (<i>Ascochyta rabiei</i> and <i>A. lentis</i>), suppression of white mould (<i>Sclerotinia sclerotiorum</i>)		
Potato	Early blight (<i>Alternaria solani</i>)	71 to 113 g	Apply prior to infection for preventative control. If conditions are favourable for disease development, make additional applications at 7 to 10 day intervals.

Application Information:

DO NOT apply during periods of dead calm or when winds are gusty. Ensure uniform coverage.

Water Volume:

Ground: Minimum of 80 L per acre.

Aerial: Minimum of 20 L per acre.

Consult nozzle manufacturers for specific nozzle and pressure recommendations. Ensure thorough coverage for optimal disease control/suppression.

How it Works:

The active ingredient metconazole is a broad spectrum triazole demethylation inhibitor (DMI) fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Canola – DO NOT exceed 1 application of this product per season. Dry bean, field pea, chickpea, lentil – DO NOT exceed 2 applications of this product per season. Potato – DO NOT exceed 3 applications of this product per season at the high rate, DO NOT exceed 4 applications of this product per season at the low rate.

Grazing: No restrictions listed.

Preharvest interval: Potato – 1 day; Dry bean, field pea, chickpea, lentil – 21 days; Canola – 45 days.

Re-entry: Potato – DO NOT re-enter treated areas within 12 hours of application. Canola – DO NOT re-enter treated areas within 6 days of application. Dry bean, field pea, chickpea, lentil – DO NOT re-enter treated areas within 9 days of application.

Re-cropping: A plant back interval of 30 days is required for all crops not listed on the *Quash* label.

Storage: Store in original tightly closed container. Protect from freezing. Store in cool, dry place. Store this product away from food or feed.

Environment: Avoid run-off from treated areas into aquatic areas. Toxic to aquatic organisms, non-target terrestrial plants and small wild animals.

Hazard Rating:



Caution – Poison

Check label for first-aid information.

For an explanation of the symbols used here see page 11.

Quilt

Fungicide Group – 3, 11
(Refer to page 312)

Company:

Syngenta Canada – PCP# 28328

Formulations:

75 g per L azoxystrobin and 125 g per L propiconazole formulated as a suspension.

Container size – 2 x 10.125 L case and 101.25 tote.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Soybeans; Legume vegetables, including: Bean Chickpea Field pea Lentil	Asian (soybean) rust (<i>Phakopsora pachyrhizi</i>)	405 to 607 mL	Make the first application at the first sign of disease. Apply the high rate only under conditions of high disease pressure. A second application at a 14 day interval may be needed if conditions persist. It is important to protect the developing pod of soybean and podded legume vegetables.
	Powdery mildew (<i>Microsphaera diffusa</i> , <i>Erysiphe pisi</i> , <i>E. polygoni</i>) Mycosphaerella blight (<i>Mycosphaerella pinodes</i>) on peas	405 mL	
Lentil Soybean	Anthracnose (<i>Colletotrichum truncatum</i>)	405 to 607 mL	Make the first application at the first sign of disease. A second application at 14 days interval may be needed if conditions persist. Good spray coverage and canopy penetration are important for best results.
Soybean	Frogeye leaf spot (<i>Cercospora sojae</i>)	405 to 607 mL	Make the first application at growth stage R3 (early pod set) and make a second application 14 days later at approximately growth stage R5.

Continued

Crops, Disease, Rates and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Barley (B) Wheat (W) Spring Wheat (SW) Winter Wheat (WW)	Net blotch (<i>Pyrenophora teres</i>) (B); Scald (<i>Rhynchosporium secalis</i>) (B); Septoria leaf spot (<i>Septoria</i> sp.) (B & W); Tan spot (<i>Pyrenophora tritici- repentis</i>) (W) Stripe rust (<i>Puccinia striiformis</i>) (B, SW & WW); Wheat leaf rust (<i>Puccinia triticina</i>) (SW & WW).	304 mL	Apply once between stem elongation and half-head emergence. When disease pressure from stripe rust or wheat leaf rust is expected to be high, the 405 ml per acre applica- tion rate should be used.
	Stripe rust (<i>Puccinia striiformis</i>) (B, SW & WW); Wheat leaf rust (<i>Puccinia triticina</i>) (SW & WW).	304 to 405 mL	Please note which crops (abbrevia- tions in brackets) correspond to each disease controlled by product.
	Barley leaf rust (<i>Puccinia hordei</i>) (B)	405 mL	
Field, Sweet, and Popping Corn (includ- ing seed production)	Rust (<i>Puccinia sorghi</i>); Northern corn leaf blight (<i>Setosphaeria tur- cicum</i>); Southern corn leaf blight (<i>Cochliobolus heterostrophus</i>); Eye spot (<i>Aureobasidium zeae</i>); Grey leafspot (<i>Cercospora zeae-maydis</i>)	304 to 405 mL	Make first application at the first sign of disease, followed by a second application 14 days after the first, if environmental conditions are favou- rable for disease development.
Oat	Septoria leaf spot (<i>Septoria</i> sp); net blotch (<i>pyrenophora teres</i>)	304 mL	Apply between stem elongation and half-head emergence.
	Crown rust (<i>puccinia coronata</i>)	304 to 405 mL	

Application Information:

Water Volume:

Ground: Apply in a minimum of 18 L of water per acre for legume vegetables and soybean. Apply in a minimum of 40 L of water per acre for other crops.

Aerial: Apply in a minimum of 18 L of water per acre.

How it Works:

The active ingredient azoxystrobin is a methoxyacrylate compound (strobilurin) with broad spectrum contact and systemic activity. The active ingredient propiconazole is a triazole fungicide with broad-spectrum systemic activity. To be used as a preventative and curative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Insecticides: *Quilt* can be tank-mixed with insecticide *Matador 120EC* for foliar disease and insect control in cereals. Consult each label for pests controlled, precautions, and specific application instructions

Note: Syngenta supports the following mixes that are not on the *Quilt* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides: *Axial*, *Broadband* + registered tank mixes, *Horizon NG*, *Sierra*, *Touchdown Total*, *Traxion*

Fungicides: *Quadris*

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Barley, wheat, oat – DO NOT exceed 1 application of this product per season. Soybean, bean, chickpea, field pea, lentil, corn – DO NOT exceed 2 applications of this product per season.

Preharvest interval: 30 days (soybeans and dry legume vegetables); 15 days (succulent podded and shelled legume vegetables); 14 days (soybean hay and dry pea hay); 45 days (wheat, barley and oat); 14 days (field corn, sweet corn, and popcorn).

Re-entry: DO NOT re-enter treated fields until 12 hours post-application.

Re-cropping: Oats and rye may be planted 45 days after *Quilt* application. DO NOT plant any other crop intended for food, grazing, or any component of animal feed or bedding within 105 days of *Quilt* application to the preceding crop unless the second crop appears on the *Quilt* label.

Storage: Store in a cool, dry, well ventilated area away from feed and foodstuffs, and out of reach of children and animals. DO NOT store at temperatures below freezing. Keep

in original container, tightly closed, during storage.

Environment: Azoxystrobin is persistent and will carry over. *Quilt* is toxic to aquatic organisms and is extremely phytotoxic to certain apple varieties. Avoid spraying when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt), or aquatic habitat. DO NOT contaminate irrigation or drinking water supplies by cleaning of equipment or disposal of wastes. Avoiding spray drift is the responsibility of the applicator.

Hazard Rating:



Caution – Poison and skin irritant

For an explanation of the symbols used here see page 11.

Ranman 400SC (2013)

See Cyazofamid on page 337.

Reason 500SC

Fungicide Group – 11
(Refer to page 312)

Company:

Bayer CropScience – PCP# 27462

Formulation:

500 g per L fenamidone formulated as a suspension concentrate. Container sizes - 2, 4 or 10 L.

Crops, Diseases and Timing:

Control of early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*) on potato. Begin application when plants are 15 to 20 cm high or when disease threatens. Apply a fungicide with a different mode of action within 7 to 10 days after each application using the shorter interval when conditions favor disease development. Ensure even application.

Rates:

Apply at 80 mL per acre as a tank mix with either Dithane DG* at 500 g per acre or Bravo 500 at 500 mL per acre.

*When using other formulations of mancozeb, adjust application rates to apply 375 g active ingredient per acre.

Application Information:

Water Volume:

Aerial: Use minimum of 14 L per acre at a pressure no less than 300kPa.

How it Works:

The active ingredient fenamidone is a strobilurin fungicide with contact activity. To be used as a preventative and

inhibitive (spore germination and antisporeulant) fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides: To be applied ONLY as a tank-mix with mancozeb fungicides or Bravo 500. Follow mixing instructions provided on the label.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 6 applications or 0.48 L per acre of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 14 days.

Re-entry: DO NOT re-enter treated areas until residues have dried.

Re-cropping: A 30 day plant-back interval is required for potato and all other crops.

Storage: DO NOT allow product to freeze. If stored more than 1 year, shake well before using. Keep away from fire, open flame or other sources of heat. Store in tightly closed container away from fertilizer, seeds, feed or food.

Environment: For ground application maintain an 8 m buffer zone between areas sprayed and aquatic systems. For aerial application allow a 10m buffer. Toxic to fish and other aquatic organisms; DO NOT apply where runoff is likely to occur.

Hazard Rating:



Caution Poison – Eye Irritant.

For an explanation of the symbols used here see page 11.

Regalia Maxx

Company:

Marrone Bio Innovations – PCP#30199
Distributed by Engage Agro Corporation

Formulation:

20% extract of *Reynoutria sachalinensis* formulated as a liquid concentrate.
Container size – 4 x 5 L.

Crops, Diseases and Timing:

Partial suppression of septoria leaf spot (*Septoria tritici*) in wheat. Apply preventatively or when disease systems first appear after initial jointing. Repeat applications in 7 to 14 day intervals depending upon crop growth and disease pressure.

Rates:

0.10% v/v in 160 to 240 L of water per acre.

Application Information:

DO NOT apply by air. When environmental conditions and plant stage are conducive to rapid disease development use *Regalia Maxx* in a rotational program with other registered fungicides.

Water Volume:

Ground: Minimum of 160 to 240 L per acre

Fungicide Group –
Not classified, bio-fungicide
(Refer to page 312)

How it Works:

Reynoutria sachalinensis is a plant extract to induce the plants' natural defense mechanisms against certain fungal and bacterial disease. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Do NOT exceed 3 applications of this product per season.

Grazing: No restrictions listed.

Preharvest Interval: May be applied up to the day of harvest.

Re-entry: DO NOT re-enter into treated areas until the spray is dried.

Re-cropping: No restrictions listed.

Storage: Store in original tightly closed container.

Environmental Hazards: DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:

None listed.

Revus

Fungicide Group – 40
(Refer to page 312)

Company:

Syngenta Canada – PCP# 29074

Formulation:

250 g per L mandipropamid. Container size – 4 x 3.78 L.

Crops, Diseases and Timing

Control of late blight (*Phytophthora infestans*) on potato. Begin applications prior to disease development. Continue applications on 7 to 10 day intervals, following resistance management guidelines.

Rates:

0.16 to 0.24 L per acre. The use of a non-ionic adjuvant (0.25% v/v) is recommended.

Application Information:

Water Volume:

Ground: Use a minimum water volume of 40 L per acre. In situations where dense canopy or pest pressure is high, use greater water volumes.

Aerial: Use a minimum water volume of 18 L per acre.

Nozzles: DO NOT apply using any type of ultra low volume (ULV) spray system.

How it Works:

The active ingredient mandipropamid is a carboxylic acid amide (CAA) fungicide with contact and systemic activity. To be used as a preventative and inhibitive (prevents spore germination) fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides: Bravo 500.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 4 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 14 days.

Re-entry: DO NOT re-enter treated area within 12 hours of application.

Re-cropping: DO NOT plant any crop which is not registered for use with *Revus* for a period of 30 days after the last application.

Storage: Store in a cool dry place away from food, beverages, and tobacco products.

Environment: To reduce runoff into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Runoff into aquatic habitats may be reduced by including a vegetative strip between the treated area and the edge of the water body.

Hazard Rating:



Caution – Poison.



Warning – Skin Irritant.
Potential Skin Sensitizer.

For an explanation of the symbols used here see page 11.

Revus Top

Fungicide Group – 3, 40

(Refer to page 312)

Revus Top is a co-pack of *Revus Fungicide* (mandipropamid, page 368) and *Inspire Fungicide* (difenoconazole, page 344). Information listed is restricted to crop, diseases, rates. For other detailed information on the component products see the product pages listed above.

Company:

Syngenta Canada

Formulation:

Revus Top has two components:

Revus (PCP #29074): 250 g per L mandipropamid formulated as a suspension

Inspire (PCP #30004): 250 g per L difenoconazole as an emulsifiable concentrate

See the component products for other information. Use the most stringent restrictions for either product.

Container size: 2x3.78 L of *Revus* + 2x3.78 L of *Inspire*

Crops, Diseases and Timing:

Control of early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*) on potato. Begin applications prior to late blight development, at early stages of early blight, or

when conditions are conducive for development of either disease.

Rates:

0.16 to 0.24 L per acre of *Revus*.

0.12 to 0.21 L per acre of *Inspire*.

See pages listed above for recommendations of resistance management and the maximum number of applications per season. Follow the most stringent restrictions for either product.

Hazard Rating:



Caution – Eye and skin irritant.



Caution – Poison.

Check label for first aid information.

For an explanation of the symbols used here see page 11.

Ridomil Gold/Bravo, Ridomil Gold SL/Bravo, Ridomil Gold 480EC, Ridomil Gold 480SL

Fungicide Group

Ridomil Gold/Bravo,
Ridomil Gold SL/Bravo - 4, M5
Ridomil Gold 480EC,
Ridomil Gold 480SL - 4

(Refer to page 312)

Company:

Syngenta Canada

(*Ridomil Gold/Bravo* – PCP#26443;

Ridomil Gold SL/Bravo – PCP#29239;

Ridomil Gold 480EC – PCP#25384;

Ridomil Gold 480SL – PCP#28474)

Formulation:

Ridomil Gold/Bravo - 500 g per L chlorothalonil and 480 g per L metalaxyl-M. Container size 8.83 L jug twin-pak.

Ridomil Gold SL/Bravo - 500 g per L chlorothalonil and 480 g per L metalaxyl-M formulated as a soluble concentrate. Container size - 8.83 L jug twin-pak.

Ridomil Gold 480EC - 480 g per L metalaxyl-M formulated as an emulsifiable concentrate. Container size - 4 x 3.78 L jugs.

Ridomil Gold 480SL - 480 g per L metalaxyl-M formulated as a soluble concentrate. Container size - 10 x 0.5 L or 4 x 3.78 L jugs.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate:	Application Timing:
Potato	Ridomil Gold/Bravo, Ridomil Gold SL/Bravo: Early blight (<i>Alternaria solani</i>), late blight (<i>Phytophthora infestans</i>), late blight tuber rot, Botrytis vine rot (<i>Botrytis cinerea</i>). Suppression of Pythium leak (<i>Pythium</i> spp.) and pink rot (<i>Phytophthora erythroseptica</i>)	Ridomil Gold/Bravo, Ridomil Gold SL/Bravo: One 8.83 L jug treats 10 acres. The entire contents of the jug must be added to the spray tank or an improper mixture will result.	Begin preventive applications early in the season when conditions are favorable for disease (before infection), no later than when the plant foliage meets within the row uniformly across the field. Apply a second and third application at 14 day intervals. Other registered contact fungicides should be applied 7 days after each application.
	Ridomil Gold 480EC, Ridomil Gold 480SL: Suppression of pink rot (<i>Phytophthora erythroseptica</i>) as in-furrow treatment.	Ridomil Gold 480EC, Ridomil Gold 480SL: 4 mL per 100 m row, applied in-furrow at planting.	

Application Information:

Water Volume:

Ground: use sufficient water to ensure thorough coverage of foliage. Use a water volume of 90 to 640 L per acre.

In-furrow treatment: use a minimum of 12 L per acre. For tank mixes with *Quadris* water volume should be 20 to 56 L per acre.

Aerial: use a minimum water volume of 20 L per acre.

How it Works:

The active ingredient metalaxyl is an acylalanine fungicide with systemic activity. The active ingredient chlorothalonil is a chloronitrile fungicide with contact activity. To be used as a preventative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Ridomil Gold/Bravo, Ridomil Gold SL/Bravo - None registered.

Ridomil Gold 480EC, Ridomil Gold 480SL - May be tank mixed with *Quadris* for in-furrow treatment to control *Rhizoctonia* stem rot, stolon canker, black scurf and suppression of pink rot.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: Ground / aerial (**Ridomil Gold/Bravo, Ridomil Gold/SL/Bravo**) - DO NOT exceed 3 applications of this product per season.
In-furrow (**Ridomil Gold 480EC, Ridomil Gold 480SL**) - DO NOT exceed 1 application of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 90 days for in-furrow application.

Re-entry: **Ridomil Gold 480EC, Ridomil Gold 480SL** - DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: A plant back interval of 30 days for root crops is required after the in-furrow application.

Storage: Protect from excessive heat.

Environment: DO NOT apply where runoff is likely to occur. DO NOT use on coarse textured gravelly soils, soils with less than 2% organic matter or in areas where the water table may be high. Avoid application by ground or air near or around bodies of water. DO NOT contaminate streams or ponds by spray drift, by cleaning equipment, or disposal of wastes. A buffer zone of 100 m for aerial application and 15 m for ground application should be observed to protect water bodies.

Hazard Rating:

Ridomil Gold/Bravo, Ridomil Gold SL/Bravo



Warning Poison - Eye irritant.

Ridomil Gold 480EC



Caution Poison. Warning - Eye irritant.

Ridomil Gold 480SL



Caution Poison. Warning - Eye irritant, skin irritant.

For an explanation of the symbols used here see page 11.

Scala SC

Fungicide Group – 9
(Refer to page 312)

Company:

Bayer CropScience – PCP#28011

Formulation:

400 g per L pyrimethanil formulated as a flowable concentrate. Container size - 2 L.

Crops, Diseases and Timing:

Control of early blight (*Alternaria solani*) on potato. Apply when plants are 15 to 20 cm high or when disease threatens. Repeat applications at 7 to 14 day intervals or as necessary to maintain disease control. If severe disease conditions exist, use the 7 day interval. Minimum spray interval is 7 days. Ensure complete coverage.

Rates:

Apply at 300 mL per acre as a tank mix with *Bravo 500*.

Application Information:

Water Volume:

Ground: Minimum of 120 L per acre.

Aerial: Minimum of 14 L per acre.

How it Works:

The active ingredient pyrimethanil is a anilinopyrimidine fungicide with contact and systemic activity. To be used as a preventative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicides: To be applied ONLY as a tank mix with *Bravo 500*. Follow mixing instructions provided on the label.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 6 applications or 2.4 L per acre of this product per season.

Grazing: No restrictions listed.

Preharvest interval: 7 days.

Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: A 30 day plant-back interval is required for potatoes and wheat and 130 days for all other crops.

Storage: DO NOT allow product to freeze. If stored more than 1 year, shake well before using. Store in tightly closed container away from fertilizer, seeds, feed or food.

Environment: Maintain a 1 m buffer zone between areas sprayed and aquatic systems. Toxic to aquatic organisms; DO NOT apply where runoff is likely to occur.

Hazard Rating:



Caution Poison – Skin Irritant.

For an explanation of the symbols used here see page 11.

Senator 70WP

Fungicide Group – 1
(Refer to page 312)

Company:

Nippon Soda Company Ltd. – PCP# 25343
Distributed by Engage Agro Corporation

Formulation:

70% thiophanate-methyl formulated as wettable powder.
Container size - 2 kg.

Crops, Diseases and Timing:

Control of sclerotinia (white mould) (*Sclerotinia sclerotium*) on white beans. Apply when conditions favour disease (i.e. warm, humid weather and heavy, dense foliage), usually during early bloom stage and prior to rows closing in. If conditions favouring disease persist, repeat applications may be warranted.

Rates:

0.7 to 0.9 kg per acre (one container treats 1.67 to 2.14 acres)

Application Information:

Water Volume:

Ground: 400 L per acre.

Aerial: 20 to 24 L per acre.

How it Works:

The active ingredient thiophanate-methyl is a methylbenzimidazole carbamate fungicide with systemic activity. To be used as a preventative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered. *Senator 70WP* is compatible with most pesticides; DO NOT mix with lime or other alkaline materials.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: No restrictions listed.

Grazing: DO NOT feed or allow livestock to graze on treated crops.

Preharvest interval: No restrictions listed.

Re-entry: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store in a dry place.

Environment: DO NOT apply to any body of water. Avoid drifting of spray onto any body of water or non-target areas. Specified buffer zones should be observed.

Hazard Rating:

None listed.

Serenade Max/Serenade CPB

Fungicide Group – 44
(Refer to page 312)

Company:

Bayer CropScience

Formulation:

Serenade Max: PCP# 28549 - 14.6% *Bacillus subtilis* (QST 713 strain) formulated as a wettable powder.

Container size - 5.44 kg.

Serenade CPB: PCP# 30143 - 1.34% *Bacillus subtilis* (QST 713 strain) formulated as an aqueous suspension.

Container size - 2 x 9.46 L per case, 511 L and 1000L tote.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Suppressed:	Application Rate (per Acre)		Application Timing:
		Serenade Max	Serenade CPB	
Bean, chickpea, lentil, pea (all types)	White mould or stem rot (<i>Sclerotinia sclerotiorum</i>); Botrytis blight or pod rot (<i>Botrytis cinerea</i>)	1.2 to 2.4 kg	1.6 to 6.1 L	Product should be applied prior to or in the early stages of disease development; repeat applications on 7 to 10 day intervals if conditions for disease persist. Use maximum label rates and shortened spray intervals for conditions conducive to rapid disease development. When conditions are conducive to heavy disease pressure, use in a rotational program with other registered fungicides.
Soybean	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)			Begin application soon after emergence and when conditions are conducive to disease development. Repeat as necessary on a 7 to 10 day interval.
	Brown spot (<i>Septoria glycines</i>) Frogeye (<i>Cercospora sojae</i>)	0.1 to 0.4 kg	0.4 to 1.6 L	
Potato	White mould (<i>Sclerotinia sclerotiorum</i>) Potato early blight (<i>Alternaria solani</i>)	1.2 to 2.4 kg	—	Begin application soon after emergence and when conditions are conducive to disease development. Repeat as necessary on a 7 to 10 day interval.
Canola	Sclerotinia stem rot (<i>Sclerotinia sclerotiorum</i>)	0.1 to 0.4 kg	0.4 to 1.6 L	Begin application at 20% to 30% bloom. A second application may be made 7 to 10 days later, at approximately 50% bloom and prior to significant petal fall, if conditions for disease development remain favourable. Use higher rates in fields with a history of heavy disease pressure.

Application Information:

Water Volume:

Use water volumes to give good canopy penetration and coverage of plant parts to be protected.

Ground application only for all crops, except canola (ground or air).

How it Works:

Bacillus subtilis is a bacterium that works as a bio-fungicide to prevent infection of labeled diseases by multi-site biochemical activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: No restrictions listed.

Grazing: No restrictions listed.

Preharvest interval: Can be applied up to and including the day of harvest.

Re-entry: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Maximum storage period of two years at room temperatures up to 25°C. Store in a dry area inaccessible to children. Store in original container.

Environment: DO NOT contaminate water, food, or feed by storage and disposal.

Hazard Rating:

None listed.

Potential skin sensitizer.

Stratego 250EC

Fungicide Group – 3, 11
(Refer to page 312)

Company:

Bayer CropScience - PCP# 27528

Formulation:

125 g per L propiconazole and 125 g per L trifloxystrobin formulated as an emulsifiable concentrate. Container size - 8.1 L and 129.6 L.

Crops, Disease, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Soybean	Asian soybean rust (<i>Phakopsora pachyrhizi</i>) and frogeye leaf spot (<i>Cercospora soja</i>)	202 mL	Begin applications preventatively from early flowering (R1) to complete pod fill (R5) when risk of rust infection is high. Spray between R3 and R4 (early pod fill) or when first symptoms occur for frogeye leaf spot.

Application Information:

Water Volume:

Ground: 40 to 80 L per acre.

Air: Minimum of 20 L per acre.

How it Works:

The active ingredient propiconazole is a triazole fungicide with broad spectrum systemic activity.

The active ingredient trifloxystrobin is a strobilurin fungicide with broad spectrum preventative activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 2 applications of this product per season.

Grazing: DO NOT graze or harvest as forage if 2 applications made in one season. If a single application is made, DO NOT allow livestock to graze or harvest as forage within 30 days after application.

Preharvest interval: 20 days.

Re-entry Interval: 12 hours.

Re-cropping: DO NOT plant back within 30 days for last application for crops not listed on label.

Storage: Store in cool dry place away from food, drink and animal feeding stuff. Keep leftover product in original container and tightly closed. Protect from freezing.

Environment: Toxic to fish and other aquatic organisms. DO NOT apply directly to any body of water and avoid drifting onto any body of water or other non-target areas. DO NOT contaminate water by cleaning of equipment or disposal of wastes. Specified buffer zones should be observed.

Hazard Rating:

Danger – Eye irritant, potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Tanos 50 DF

Fungicide Group – 11, 27
(Refer to page 312)

Company:

E.I. duPont Canada Company – PCP# 27435

Formulation:

25% famoxadone and 25% cymoxanil formulated as a dry flowable.

Container size - 2.5 to 5 kg.

Crops, Diseases and Timing:

Potato and Field Tomato - Early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*). Make the first application following one or two applications of a preventative broad spectrum fungicide such as chlorothalonil or mancozeb. A minimum 12 day application interval must pass between the 1st and 2nd application of *Tanos 50 DF*. A minimum 24 day application interval must pass between the 2nd and 3rd application of *Tanos 50 DF*. Fungicides other than *Tanos 50 DF* may be used as necessary to protect the crop during these intervals.

Rates:

225 to 340 g per acre.

Application Information:

Water volume: Use sufficient water to obtain thorough coverage. With a conventional sprayer use no less than 100 to 120 L per acre. With an air-assisted sprayer use no less than 44 L per acre. Ground application only.

How it Works:

The active ingredient cymoxanil is a cyanoacetamide-oxime fungicide with locally systemic activity. The active ingredient famoxadone is a strobilurin fungicide with broad spectrum activity. To be used as a preventative, curative and inhibitive (against sporulation) fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered. Tank mix solutions containing boron may affect product solubility. When using boron containing solutions, add the correct amount of *Tanos 50 DI* first and boron containing solution last.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 6 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: Potato 14 days and field tomato 7 days.

Re-entry: DO NOT re-enter treated areas within 24 hours of application.

Re-cropping: Crops that are on the product label (potato and field tomato) may be planted back at any time. A 30-day plantback interval is required for cereal grains. All other crops may be planted following a 1 year interval.

Storage: Store product closed in original container only. Protect against humid air and water. Avoid contact with food, drink and livestock feed material.

Environment: Toxic to fish and aquatic organisms. Observe prescribed buffer zones. Toxic to birds, mammals and harmful to beneficial arthropods. Minimize off-target drift to reduce the effects on wildlife at the field boundary. DO NOT apply to areas prone to run-off.

Hazard Rating:

Warning Poison – Eye Irritant.

For an explanation of the symbols used here see page 11.

Tattoo C

Fungicide Group – 28, M5
(Refer to page 312)

Company:

Bayer CropScience – PCP# 24544

Formulation:

375 g per L propamocarb HCl and 375 g per L chlorothalonil formulated as a suspension. Container size - 10 L.

Crops, Diseases and Timing:

Control of late blight (*Phytophthora infestans*) on potato. Begin applications when conditions are favorable for disease, but before infection, and continue on 7 to 14 day intervals until threat of disease is over. Use the 7 day

interval when the risk and conditions for disease are high. To avoid resistance, rotating and alternating applications with fungicides having different modes of action is recommended if multiple fungicide applications are required.

Rates:

1.09 L per acre.

Application Information:

Water Volume (ground only): 80 to 120 L per acre.

How it Works:

The active ingredient propamocarb HCl is a carbamate fungicide with systemic activity. Chlorothalonil is a chloronitrile fungicide with contact activity. To be used as a preventative fungicide application. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 3 applications of this product per season.

Grazing: DO NOT feed treated crops to livestock.

Preharvest interval: 7 days.

Re-entry: DO NOT re-enter treated areas within 48 hours after treatment. If required, individuals may re-enter treat-

ed areas within 48 hours for short tasks not involving hand labour, provided that 4 hours have passed since application and that long pants and a long-sleeve shirt are worn.

Re-cropping: DO NOT plant a new crop in the treated area within 120 days of the last application.

Storage: Keep away from fire, open flame or other sources of heat. DO NOT store below freezing. Store the tightly closed container away from seeds, fertilizers, plants and food-stuffs.

Environment: DO NOT apply directly to water or areas where surface water is present. DO NOT apply where runoff is likely to occur. DO NOT contaminate water when disposing of equipment wash waters. Allow a buffer zone of 15 m around bodies of water when applying.

Hazard Rating:

Ⓐ Danger – Corrosive to eyes.

Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Tebuconazole

Folicur 432F/Folicur 250EW/Palliser/Fuse

Fungicide Group – 3
(Refer to page 312)

Company:

Bayer CropScience (Folicur 432F – PCP#25940, Folicur 250EW – PCP#29820)

Loveland Products (formerly Viterra) (Palliser – PCP#30491)

Syngenta Canada (Fuse – PCP#30492)

Formulation:

Folicur 432F/Fuse – 432 g per L tebuconazole formulated as a flowable liquid. Container size – 4.73 L.

Folicur 250EW – 250 g per L tebuconazole formulated as an emulsion in water. Container size – 8.1 L.

Palliser – 432 g per L tebuconazole formulated as a flowable liquid. Container size – 9.46 L.

Crop, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):		Timing:
		<i>Folicur 432F</i> */ <i>Palliser/Fuse</i> *	<i>Folicur</i> <i>250EW</i>	
Wheat (includes spring, winter and durum)	Fusarium head blight (scab) – suppression only (<i>Fusarium</i> spp.)	118 mL	202 mL	<p>Timing of application is critical: Apply within the time period from when at least 75% of the wheat heads on the main stem are fully emerged to when 50% of the heads on the main stem are in flower.</p> <p>Spray coverage is essential for optimum efficacy: Spray equipment must be set to provide good coverage to heads (e.g. forward and backward mounted nozzles, or nozzles that have a two-directional spray).</p>
	Septoria glume blotch (<i>Stagonospora</i> <i>nodorum</i>)			
	Tan spot (<i>Pyrenophora tritici- repentis</i>), septoria leaf blotch (<i>Septoria tritici</i>), leaf rust (<i>Puccinia triticina</i>) stem rust (<i>Puccinia graminis</i>) stripe rust (<i>Puccinia striiformis</i>)	89 to 118 mL	152 to 202 mL	
	Powdery mildew (<i>Erysiphe graminis</i>)	118 mL	202 mL	<p>Apply at the first sign or very early stage of disease, up to the end of the flowering stage.</p> <p>Use the higher rate when weather conditions are conducive for disease.</p>
Barley	Net blotch (<i>Pyrenophora teres</i>), spot blotch (<i>Cochliobolus sativus</i>), scald (<i>Rhynchosporium secalis</i>), leaf rust (<i>Puccinia hordei</i>), stem rust (<i>Puccinia graminis</i>), stripe rust (<i>Puccinia striiformis</i>), septoria leaf blotch (<i>Septoria passerinii</i>), powdery mildew (<i>Erysiphe graminis</i>)	89 to 118 mL	152 to 202 mL	
Oat	Crown rust (<i>Puccinia coro- nata</i>), stem rust (<i>Puccinia graminis</i>)	89 mL	152 mL	

* *Folicur 432F*, *Palliser* and *Fuse* are recommended to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 1.25 L per 1000 L of spray solution.

Continued

Crop, Diseases, Rates and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate (per acre):		Timing:
		<i>Folicur</i> 432F*	<i>Folicur</i> 250EW	
Soybean	Asian soybean rust (<i>Phakopsora pachyrhizi</i>), frog-eye leaf spot (<i>Cercospora sojina</i>), powdery mildew (<i>Microspheeta diffusa</i>) **	89 to 118 mL	152 to 202 mL	Apply when first symptoms of disease can be found or risk of infection is imminent. Use the higher rate when disease pressure is severe.

* *Folicur* 432F is to be used with a registered non-ionic surfactant, such as Agral 90 or AgSurf, at 1.25 L per 1000 L of spray solution.

** *Folicur* 432F is not registered for control of powdery mildew in soybeans.

Application Information:

Water Volume:

Ground: Minimum 40 L per acre. Ensure thorough coverage of all wheat heads. Avoid excessive water volumes (maximum 80 L per acre) at flowering time because this can increase the risk of infection.

Aerial: Minimum 19 L per acre.

How it Works:

The active ingredient tebuconazole is a triazole demethylation inhibitor (DMI) fungicide with systemic broad-spectrum activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Herbicides: In spring wheat and barley, *Folicur* 432F *Palliser* or *Fuse* may be tank mixed with *Refine Extra* for leaf diseases and respective weeds controlled (consult labels). In spring wheat only, *Folicur* 432F *Palliser* or *Fuse* may be tank-mixed with *Buctril M* for leaf diseases and respective weeds controlled (consult labels).

Insecticides: For control of orange wheat blossom midge (*Sitodiplosis mosellana*) in wheat, *Folicur* 432F, *Palliser* or *Fuse* may be tank mixed with *Lorsban 4E* labeled rates. See respective labels for directions and use precautions.

Fungicides: None registered.

Bayer CropScience also supports the following mixes that are not on the *Folicur* EW label. Apply mixes according to the most restrictive use limitations for either product:

Insecticides: *Decis*, *Lorsban*, *Serim NLR*

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed one application of this product per season.

Grazing: DO NOT allow livestock to graze or feed green forage to livestock prior to 6 days after treatment. Straw cut after harvest may be fed or used for bedding.

Preharvest interval: 36 days.

Re-entry: DO NOT re-enter treated areas within 12 hours of application.

Re-cropping: Treated areas may be replanted immediately following harvest with any crop listed on the *Folicur* 432F/*Folicur* 250EW labels. For crops not listed on these labels, DO NOT replant treated areas for 120 days after last application.

Storage: Store in a cool, dry place and prevent cross contamination with other pesticides, fertilizers, food and feed.

Environment: Any products containing tebuconazole should not be used in areas treated with this product during the previous season (use only in alternate years). This product is toxic to birds, small wild animals, aquatic organisms, and non-target plants.

DO NOT apply directly to water, or to areas where surface water is present. Maintain a buffer zone of 30 m near aquatic areas. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:



Danger – Skin irritant.



Caution – Eye irritant.

For an explanation of the symbols used here see page 11.

Twinline

Fungicide Group – 3, 11
(Refer to page 312)

Company:

BASF Canada – PCP#30337

Formulation:

130 g per L pyraclostrobin and 80 g per L metconazole formulated as a liquid.

Container size – Case (2 x 8.1L); 64 L drum; 128 L Shuttle; or 400 L tote.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Wheat (all types) Triticale	Tan spot (<i>Pyrenophora tritici-repentis</i>), septoria leaf spot (<i>Septoria tritici</i> or <i>Stagonospora nodorum</i>), leaf rust (<i>Puccinia recondita</i>), spot blotch (<i>Cochliobolus sativus</i>), stripe rust (<i>Puccinia striiformis</i>), powdery mildew (<i>Erysiphe graminis</i> f. sp. <i>tritici</i>)	154 to 202 mL	Prior to disease development or at onset of disease. Optimal application timing is at the flag leaf stage. Use the 202 mL per acre rate to obtain extended protection with maximum yield benefits.
Barley	Net blotch (<i>Pyrenophora teres</i>), spot blotch (<i>Cochliobolus sativus</i>), scald (<i>Rhynchosporium secalis</i>), stripe rust (<i>Puccinia striiformis</i>)		
Oats	Crown rust (<i>Puccinia coronata</i>)		
Rye	Leaf rust (<i>Puccinia recondita</i>), powdery mildew (<i>Erysiphe graminis</i>)		

Application Information:

Water Volume:

Ground: Minimum of 40 L per acre.

Aerial: Minimum of 20 L per acre.

How it Works:

The active ingredient metconazole is a broad spectrum triazole demethylation inhibitor (DMI) fungicide with systemic activity. The active ingredient pyraclostrobin is a strobilurin fungicide with broad spectrum contact and systemic activity. Best utilized as a preventative application when environmental conditions are favourable for disease development. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 2 applications of this product per season.

Grazing: No restrictions listed.

Preharvest interval: Apply no later than the end of flowering.

Re-entry: DO NOT re-enter treated areas within 6 days of application.

Re-cropping: A plant back interval of 35 days is required for all crops not listed on the label.

Storage: Store in original tightly closed container. Protect from freezing.

Environment: Avoid run-off from treated areas into aquatic areas. Toxic to aquatic organisms, non-target terrestrial plants and small wild animals. For ground application, buffer zones must be 1 m for protection of terrestrial habitats and aquatic habitats greater than 1 m deep and buffer zones

must be 5 m from aquatic habitats less than 1 m deep. For aerial application, buffer zones must be 10 m for protection of terrestrial habitats and aquatic habitats greater than 1 m deep and buffer zones must be 250 m from aquatic habitats less than 1 m deep.

Hazard Rating:



Danger – Poison

Eye and skin irritant.

For an explanation of the symbols used here see page 11.

Vertisan

Fungicide Group – 7
(Refer to page 312)

Company:

E.I. duPont – PCP #30332

Formulation:

200 g/L penthiopyrad formulated as an emulsifiable concentrate.

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Canola	Sclerotinia stem rot / white mould (<i>Sclerotinia sclerotiorum</i>)	500 to 600 ml	Apply at 20 to 50% bloom stage prior to disease development. Under high disease pressure, make a second application 7 to 14 days later. Use higher rate and shorter interval when disease pressure is high.
Dry legumes (chickpea, lentil, field, navy, pinto bean, field pea):	Ascochyta blight (<i>Ascochyta</i> spp.)	400 to 600 ml	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Gray mould (<i>Botrytis cinerea</i>)	500 to 600 ml	
	Suppression of Asian soybean rust (<i>Phakospora pachyrhizi</i>)	400 to 700 ml	
Cereals	Suppression of septoria leaf spot (<i>Septoria tritici</i>) in wheat, barley and triticale. Black stem rust (<i>Puccinia graminis</i> f.sp. <i>tritici</i>) in wheat, oats, triticale, barley. Brown leaf rust, orange leaf rust (<i>Puccinia recondita</i> f. sp. <i>tritici</i>) in wheat, rye, triticale.	500 to 700 ml	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high. For optimizing yield and flag leaf disease control apply at Feeke's 9, "flag leaf out". Do not apply after flowering (Feeke's 10.5.1)
Corn	Suppression of grey leaf spot (<i>Cercospora zeae-maydis</i>) and control of common rust (<i>Puccinia sorghi</i>)	400 to 700 ml	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high. For suppression of grey leaf spot, add a non-ionic surfactant.
Soybean	Suppression of Asian soybean rust (<i>Phakospora pachyrhizi</i>), brown spot (<i>Septoria glycines</i>), frog-eye leafspot (<i>Cercospora sojina</i>)	400 to 700 ml	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.

Continued

Crops, Diseases, Rates and Timing *continued*:

Crop:	Diseases Controlled:	Application Rate (per acre):	Application Timing:
Sunflower	Suppression of rust (<i>Puccinia helianthi</i>) and sclerotinia head rot (<i>Sclerotinia sclerotiorum</i>)	700 ml	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
Potato	Suppression of early blight (<i>Alternaria solani</i>)	400 to 700 ml	Begin applications prior to disease development and continue on a 7 to 14 day interval. Use higher rate and shorter interval when disease pressure is high.
	Gray mould (<i>Botrytis cinerea</i>).	500 to 600 ml	
	Stem rot (<i>Rhizoctonia solani</i>).	15.5 to 31 ml per 100 m row	In furrow at planting, using 1.4 to 1.75 L water per 100 row meters. Do not exceed 0.7 L per acre.

Application Information:

Water Volume:

Ground: 45 L per acre.

Aerial: 16 L per acre. Use sufficient water to obtain thorough coverage of plants. DO NOT apply under periods of dead calm. Avoid application of this product when winds are gusty.

How it Works:

The active ingredient penthiopyrad is a carboxamide fungicide with broad spectrum, locally systemic and curative properties recommended for foliar and soil borne plant diseases. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310. Make no more than 2 sequential applications before switching to a fungicide with a different mode of action.

Maximum seasonal use rate: Canola, dry legumes, soybean – 1.2 L per acre, cereal grains, corn – 1.4 L per acre and sunflower – 1.8 L per acre, potato – 2 L per acre.

Grazing: Cereals, corn and soybeans may be used for grazing or forage 0 days after the last application.

Pre-harvest interval: DO NOT apply within the following number of days prior to harvest canola and dry legumes (21), soybean and sunflower (14), corn and potato (7), cereal grains – no restriction

Re-Entry: DO NOT re-enter treated areas until 12 hours after application. For corn detasselling do not enter treated areas for 3 days.

Re-cropping: Crops and crop groups on the *Vertisan* label as well as the following crops may be planted immediately after harvest: alfalfa, low growing berries (strawberries), *Brassica* (cole) leafy vegetable, bulb vegetable (onion), cucurbit vegetables (cucumber, melons, squash), fruiting vegetables (tomato, pepper), leafy vegetables (lettuce, celery, spinach), legume vegetables (succulent), root vegetables (carrot, radish, turnip). All other crops cannot be planted until 12 months after last application.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

Environment: This product is toxic to aquatic organisms. When using *Vertisan* consult the product label for buffer zones.

Hazard Rating:



Danger – Eye irritant and skin irritant

Potential skin sensitizer.

For an explanation of the symbols used here see page 11.

Zampro

Fungicide Group – 40, 45
(Refer to page 312)

Company:

BASF Canada – PCP#30321
Distributed by Engage Agro Corporation

Formulation:

225 g per L dimethomorph and 300 g per L ametoctradin formulated as a suspension.
Container size – 4 x 4.14 L

Crops, Diseases, Rates and Timing:

Crop:	Diseases Controlled:	Application Rate (L per acre):	Application Timing:
Potato	Late blight (<i>Phytophthora infestans</i>)	0.32 to 0.4	Begin applications prior to disease development and continue on a 5 to 10 day interval. Use the higher rate and shorter interval when disease pressure is high. The addition of a spreading/penetrating adjuvant is recommended to improve disease control performance.
	Tuber blight (<i>Phytophthora infestans</i>)	0.4	When used in accordance to label recommendations, <i>Zampro</i> also reduces tuber blight when applied immediately prior to or after vine kill.

Application Information:

Water Volume:

Ground: Minimum of 80 L per acre.

Aerial: Minimum of 20 L per acre.

How it Works:

The active ingredient dimethomorph is a carboxylic acid amide fungicide with contact, systemic and antispore activity. The active ingredient ametoctradin is a quinone x inhibitor fungicide with contact and antispore activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Maximum number of applications: DO NOT exceed 3 applications of this product per season

Grazing: No restrictions listed.

Preharvest Interval: 4 days.

Re-entry: Re-entry interval after application is 12 hours.

Re-cropping: A plant back interval of 30 days is required for all crops not listed on the label

Storage: Store in original tightly closed container. Protect from freezing.

Environmental Hazards: Avoid run-off from treated areas into aquatic areas. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Toxicity: Toxic to aquatic organisms.

Hazard Rating:



Warning Poison – Eye Irritant.

For an explanation of the symbols used here see page 11.

Seed Treatment Tables

Table 7. Seed Treatment Products for Cereal Crops:

PRODUCTS	Page	CROPS							DISEASES										INSECTS			
		Wheat	Barley	Oat	Rye	Corn	Triticale	Pearl Millet and Proso Millet	Seed and Seedling Rots / Blights	Pythium Seed Rot and Damping off	Common root rot / Crown Rot **	Loose smut (wheat, barley, oat)	Common bunt (wheat)	Covered smut (barley, oat)	False Loose smut (barley)	Seed-borne Septoria (wheat, barley, rye)	Early season leaf diseases	Seed and Soil-Borne <i>Rhizoctonia</i> spp. (barley, wheat, oats, corn)	Fusarium (seed and soil borne)	Wireworms	Seed Corn Maggot	Corn flea beetle
Agrox B-2/ Agrox CD	390					X			•												•	
Agrox FL	391					X			•													
Allegiance FL	424	X*	X*	X*	X*	X*	X*		•													
Armour/ Armour RTU	444	X	X	X					(1)		•	•	•	•	•							
Belmont 2.7 FS	424	X*	X*	X*	X*	X*	X*		•													
Charler RTU	444	X	X	X					(1)		•	•	•	•	•							
Cruiser 5FS†	396	X	X		X	X	X													•	•	
Cruiser Maxx Cereals/ Cruiser Maxx Cereals Commercial†	399	X	X	X	X		X		(1)	•	•	(4)	•	•	•					•		
Cruiser Maxx Vibrance Cereals	399	X	X	X	X		X		(1)	•	•	•	•	•	•	•	(9)	•		•		
DB-Red L	403	X	X	X	X				(1)		•		(3)	•	•							
Dividend XL RTA	405	X	X	X	X	X	X		(1)	•	•	(4)	(3)	•	•	•	(9/10)					
EverGol Energy	407	X	X	X	X	X	X	X	(1)	•	•	•	•	•	•		(6)					
Gemini	410	X	X	X					(1)	(2)	(2)	•	•	•	•							
Insure Cereal	416	X	X	X	X		X		•	•	•	•	•	•	•							
Maxim Quattro†	422					X			•	•								•				
Poncho 600FS†	428					X														(5)	(5)	
Proseed†	427	X	X	X	X	X	X		(1)									•				
Rancona Apex	430	X	X	X	X		X		(1)		•	•	•	•	•							
Raxil MD	432	X	X	X					(1)	•	•	•	•	•	•							
Raxil PRO	433	X	X	X					•	•	•	•	•	•	•		(6)					
Raxil PRO Shield	435	X	X	X					(1)	•	•	•	•	•	•		(6)		**	•**		
Raxil WW	436	X	X	X					(1)	•	•	•	•	•	•					•**		
Sombrero	414					X														•		•
Stress Shield 600	439	X	X	X																•		
Thiram 75WP†	440					X			(1)	•												
Vibrance XL	448	X	X	X	X		X		(1)	•	•	•	•	•	•	•	(9)	•				
Vitaflo 280/ Vitaflo 220/ Vitaflo 5P/ Vitaflo	450	X	X	X	X	X	X***		(1)	•	•	•	•	•	•	(7)	(6)					

* For export markets only.

** Suppression only.

*** Vitafo 280 only.

† Available to commercial seed treaters only.

(1) Includes seed rots and blights caused by *Fusarium* species.

(2) Except oats.

(3) Includes rye.

(4) Except barley.

(5) Includes black cutworms, corn flea beetle, and white grubs.

(6) Barley leaf stripe control and net blotch suppression.

(7) Wheat only.

(9) Septoria leaf blotch on winter wheat.

(10) *Penicillium* 3-leaf dieback on corn.

Table 8. Seed Treatment Products for Oilseed Crops:

PRODUCTS	Page	CROPS					DISEASES					INSECTS	
		Canola	Mustard, Rapeseed	Flax	Sunflower	Safflower	Seed and Seedling Rots / Blights	<i>Fusarium</i> Seed Rot and Damping Off	Seed-borne Blackleg (canola)	Seed-borne <i>Alternaria</i>	Downy mildew (sunflower)	Flea Beetles	Cutworms
Acceleron Seed Treatment Package	389	X					•*					•	
Allegiance FL	424	X			X			•			•		
Belmont 2.7 FS	424	X			X			•			•		
Gaucha CS FL†	409	X	X				•	•	•	•		•	
Helix Vibrance	413	X	X				•	•	•	•		•	
Helix XTra†	413	X	X				•	•	•	•		•	
Lumidern	419											•	•
Prosper EverGo†/FX†	428	X	X				•	•	•	•		•	
Poncho 600†	428	X										•	
Rancona RS	431	X					•		•				
Sombrero	414	X	X**									•	
Thiram 75WP	440		X			X	•	•					
Vitaflo 280 / Vitaflo 220 / Vitaflo SP / Vitaflo	450			X			•						

† Available to commercial seed treaters only.

* Partial Suppression

** Condiment-type only

Table 9. Seed Treatment Products for Pulse Crops:

PRODUCTS	Page	CROPS							DISEASES										INSECTS									
		Beans	Chickpea	Fababean	Lentil	Lupin	Pea	Soybean	Damping-off caused by <i>Rhizoctonia solani</i>	Seed and Seedling Rots / Blights**	Pythium Seed Rot and Damping Off	Seed-borne ascochyta (chickpea, lentil, pea)	Seed-borne <i>Botrytis</i> spp.	Seed-borne anthracnose (beans)	Seed-borne bacterial brown spot (beans)	Seed-borne common blight (beans)	Seed-borne halo blight (beans)	Seed-borne Phomopsis (soybean)	Sudden death syndrome (Fusarium)	White mould (<i>Sclerotinia</i>)	Bean leaf beetle	European Chafer	Pea leaf weevil (pea only)	Potato leafhopper (dry beans only)	Root maggots	Seed corn maggot	Soybean aphid	Wireworms
Agrox B-2 / Agrox CD	390	X	X				X	X		•															•	•		
Agrox FL	391	X	X		X		X	X		•																		
Allegiance FL	424	X	X		X ⁽¹⁾		X	X			②																	
Apron Advance / Apron Maxx RTA / Apron Maxx RFC	392	X	X	X	X	X	X	X ⁽⁶⁾		•	•	•		•				②										
Belmont 2.7 FS	424	X	X		X ¹		X	X			②																	
Crown	395				X					•		③																
Cruiser 5FS [†]	396	X	X		X		X	X																•	•		•	•
Cruiser Maxx Beans [†]	397	X						X		•	•			•				②				•		•		•	•	•
Cruiser Maxx Pulses [†]	397		X		X		X			•	•	•											•					•
Cruiser Maxx Vibrance Beans [†]	397							X		•	•	•						②				•		•		•	•	•
EverGol Energy	407	X	X		X		X	X		•	•	•	*					•										
Heads Up Plant Protectant	412	X						X		•									• ⁽⁵⁾	•								
Kocide 2000	418	X	X							•					•	•	•											
Stress Shield 600	439	X	X	X	X		X	X												• ⁽⁵⁾			•	•		• ⁽⁵⁾	• ⁽⁵⁾	•
Thiram 75WP	440	X	X				X	X		•	•																	
Trilex AL	441	X	X		X		X	X		•	•	•*						•										
Trilex EverGol	442	X	X		X ⁽¹⁾		X			•	•	•*	•					•										
Vibrance Maxx RTA	447		X		X		X			•	•	•	•															
Vibrance Maxx RFC	447		X		X		X			•	•	•	•															
Vitaflo 280 / Vitaflo 220 / Vitaflo SP / Vitaflo	450	X	X		X***		X***	X		•	•	④						•										

* Suppression only.

** See product page for list of specific pathogens controlled.

*** Vitaflo 280 only.

† Available to commercial seed treaters only.

(1) Only for low-tannin lentils.

(2) Includes early season *Phytophthora* root rot.

(3) Except peas.

(4) Except chickpeas and lentils.

(5) Soybean.

(6) Except Apron Advance.

Table 10. Seed Treatment Products for Forages (Grasses and Legumes) and Special Crops:

PRODUCTS	Page	CROPS				DISEASES		
		Legumes (alfalfa, bird's-foot trefoil, clover, vetch, and sainfoin)	Grasses	Buckwheat, millet, sorghum (S)	Herbs and Spices	Seed and Seedling Rots / Blights	Pythium Seed Rot and Damping Off	wireworms (suppression)
Allegiance FL	424	X	X	S			•	
Belmont 2.7 FS	424	X	X	S			•	
Cruiser 5FS	396			X				•
Cruiser Maxx Cereals Seed Treatment / Cruiser Maxx Cereals Commercial [†]	399			X		(3)	•	•
Dividend XL RTA	405			X		•	•	
Proseed [†]	427			X**		(2)		
Thiram 75WP	440	X*	X			•	•	

[†] Available to commercial seed treaters only

* For alfalfa only

** Buckwheat and sorghum

(1) Includes early season

Phytophthora root rot.

(2) Includes *Fusarium* and *Rhizoctonia* spp.

(3) Includes *Fusarium*.

Table 11. Seed Treatments for Potatoes (includes post-harvest storage products and disinfectants):

PRODUCTS	Page	DISEASES							INSECTS					POST-HARVEST				
		Blackleg	Black Scurf	Silver Scurf	Common Scab (seed-borne)	Fusarium Seed Piece Decay	Verticillium Wilt		Rhizoctonia stem and stolon canker	Colorado Potato Beetle	Potato Flea Beetle	Potato Leafhopper	Aphids	Wireworms	Silver Scurf	Storage rots caused by <i>Fusarium</i> , <i>Phoma</i> , <i>Oospora</i> and <i>Rhizoctonia</i>	Bacterial Soft Rot	Bacterial Ring Rot
Aclara	498								•		•	•						
Confine	394													•**				•**
Cruiser Maxx D Potatoes	402		•	•					•		•	•			•			
General Storage Disinfectant	411																(1)	
Heads Up Plant Protectant	412		•															
Imidacloprid Products (Admire SPT, Alias 240 SC)	414								•	•	•	•						
Mancozeb Products (Potato ST 16, Solan MZ and Tuberseal)	420					•												
Maxim D	421		•**	•		•		•										
Maxim Liquid PSP Maxim PSP/Maxim MZ PSP	421		•	•		•												
Mertect SC	423													•	•			
Polyram 16D	426				•	•												
Senator PSP	437	•		•		•	•											
StorOx	438													•	(2)	•		
Titan ST	428								•	•	•	•	•					

* Before using any pesticide on potatoes, consult the list of Agricultural Pesticides Approved for Use from Simplot Canada and McCain Foods (Canada).

** Suppression only

(1) Not for use on potatoes. Use for disinfecting potato storages and equipment.

(2) *Fusarium* only

Seed Treatment Product Pages

Acceleron DX-910

Fungicide Group – U
(Refer to page 312)

Acceleron DX-910 Biological Seed Treatment is currently only available as a co-pack with *Helix Xtra* (see *Acceleron Seed Treatment Package* on page 399).

Company:

Becker Underwood Canada (PCP #29961)
Distributed by Monsanto Canada Inc.

Formulation:

0.18% *Bacillus subtilis*, strain MBI 600 (not less than 2.2×10^{10} viable spores per mL)
Container size – 1L to 1050 L.

Crops, Diseases, and Rate:

For partial suppression of seedling diseases (seed rot, pre and post emergent damping-off, seedling blight and root rot) caused by *Rhizoctonia* spp. and *Fusarium* spp. in canola, apply 160 mL of *Acceleron DX-910* per 100 kg of seed.

Application Information:

Apply *Acceleron DX-910* through commercial seed treating equipment. If used alone, slowly add *Acceleron DX-910* to sufficient water for good seed coverage and continue mixing until uniform. If used in combination, add the required amount of the chemical seed treatment first, then slowly add *Acceleron DX-910* to the solution and continue mixing until uniform. Do not store mixed solutions for longer than 72 hours before applying to the seed.

How it Works:

Bacillus subtilis, strain MBI 600 is a seed applied biological fungicide for the partial control of registered diseases. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

For control of seedling diseases (seed rot, pre and post emergent damping-off, seedling blight and root rot) caused by *Rhizoctonia* spp. and *Fusarium* spp. in canola, apply 160 mL of *Acceleron DX-910* tank-mixed with *Helix*, *Helix Xtra*, or *Prosper FL* at labeled rates per 100 kg of seed.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store away from food and feed to prevent contamination. Store in a dry area for up to 2 years. *Acceleron DX-910* treated canola seed can be stored without loss of germination or disease control. However, due to seed quality and seed storage conditions no claims are made to guarantee the germination of carryover seed.

Environment: DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:

Potential sensitizer.

For an explanation of the symbols used here see page 11.

Acceleron Seed Treatment Package

Fungicide Group – 3, 4, 12, and U
Insecticide Group – 4A

(Refer to page 312 and 456)

Acceleron Seed Treatment Package is a co-pack of *Acceleron*

DX-910 Biological Seed Treatment (*Bacillus subtilis*, page 388) and *Helix Xtra* (thiamethoxam insecticide, and difenoconazole, metalaxyl-M, and fludioxonil fungicides, page 413). Information listed is limited to crop, diseases, insects and rates. For other detailed information on the component products see the product pages listed above.

Company:

Distributed by Monsanto Canada Inc.

Acceleron: Becker Underwood Canada (PCP #30438)

Helix Xtra: Syngenta Canada (PCP #26638)

Formulation:

0.18% *Bacillus subtilis*, strain MBI 600 (not less than 2.2×10^{10} viable spores per mL)

Helix Xtra (PCP #26638): 20.7% thiamethoxam, 1.25% difenoconazole, 0.39% metalaxyl-M, 0.13% fludioxonil formulated as a liquid seed treatment.

Crops:

Canola

Diseases:

For partial suppression of seedling diseases (seed rot, pre and post emergent damping-off, seedling blight and root rot) caused by *Rhizoctonia* spp. and *Fusarium* spp.

Insects:

Early season control of flea beetles (28 to 35 days).

Rate:

Apply 160 mL of *Acceleron DX-910* tank-mixed with 1.5 L *Helix Xtra* per 100 kg of seed.

Hazard Rating:

Potential sensitizer.



Caution – Poison

For an explanation of the symbols used here see page 11.

Agrox B-2/Agrox CD

Note: These products are being discontinued, they will no longer be available for use after December 31, 2014.

**Fungicide Group – M4,
Insecticide Group – 1B**
(Refer to page 312 and 456)

Company:

Norac Concepts Inc.
(Agrox B-2 – PCP# 26956; Agrox CD – PCP# 26957)

Formulation:

Agrox B-2: 33.5% captan, 11% diazinon.

Agrox CD: 15% captan, 15% diazinon.

Products formulated as wettable powders.

Container sizes - Agrox B-2 - 2.0 kg; Agrox CD - 0.2, 0.6, 1 kg.

Crops:

Corn, bean, soybean, pea

Diseases:

Agrox B-2: Control of seedling blights and seed rot.

Agrox CD: Supplemental control of seedling blight and seed rot.

Insects:

Agrox B-2: Control of seed and root maggots.

Agrox CD: Control of seed corn maggots.

Rate and Application Information:

Agrox B-2:

Dry application: Apply 320 g for peas, beans, soybeans and 340 g for corn per 100 kg of seed.

Slurry machines: adjust to apply 1240 mL (1260 mL for corn) of slurry per 100 kg of seed and use 250 g of seed treatment per L of water.

Mixing in bucket (25 kg of seed): 84 g seed treatment per 500 mL of water per 25 kg of seed.

Agrox CD: Apply the contents of one 200 g container to 100 kg (4 units) of seed at planting or near planting time (50 g per 25 kg of seed).

How it Works:

The active ingredient captan is a phthalimide fungicide with multi-site contact activity. For more information refer to "Fungicide Modes of Action" on page 311. The diazinon component is an organophosphate insecticide with contact and stomach activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

Use *Agrox CD* only on seed previously treated with captan or thiram products. DO NOT use on seed already treated with an insecticide (other than methoxychlor or malathion). DO NOT use *Agrox B-2* on seed already treated with an insecticide.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled as follows: "This seed has been treated with diazinon and captan. Poisonous to humans and animals. DO NOT use for food or feed purposes. DO NOT sell to oil mills."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store in a cool, dry place. Use seed within one month of treatment.

Environment: DO NOT contaminate food, feed, or any body of water.

Hazard Rating:



Danger – Poison



Warning – contains the allergen soy.

For an explanation of the symbols used here see page 11.

Agrox FL

Fungicide Group – M4
(Refer to page 312)

Company:

Norac Concepts Inc. – PCP#12028

Formulation:

30% captan formulated as a flowable suspension seed treatment.

Container sizes - 20 L, 415 L, 1000 L returnable container.

Crops and Rates:

Crop:	Agrox FL (mL per 100 kg seed)
Beans (various)	280
Chickpea	280
Lentil	280
Peas (various)	280
Soybean	280
Corn (field)	120* to 200
Corn (sweet)	240* to 340

* Product is to be applied at this rate only by a professional applicator using equipment which will assure complete and uniform coverage.

Diseases Controlled:

Storage rot, seed decay, root rot, damping off, seedling blights.

Application Information:

For use prior to storage or as a seed treatment. Mix the recommended amount of *Agrox FL* with the amount of water required for the slurry treater equipment to be used. Seed treated by the slurry method should not be bagged or stacked until it has dried. A colourant must be added to this product to colour the treated seed.

How it Works:

The active ingredient captan is a phthalimide fungicide with multi-site protective activity. For more information refer to "Fungicide Modes of Action" on page 311.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled, "This seed has been treated with *Agrox FL*. Poisonous to man and animals. DO NOT use for food or feed. DO NOT sell to oil mills."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: DO NOT freeze. Product must be stored at ambient temperatures above 0°C and must not be stored with herbicides, feed, food or fertilizer.

Environment: DO NOT contaminate food, feed, or any body of water.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Allegiance FL (2013)

See Metalaxyl on page 424.

Apron Advance/ Apron Maxx RTA/ Apron Maxx RFC

Fungicide Group –
Apron Advance – 1, 4, 12
Apron Maxx RTA/RFC – 4, 12
(Refer to page 312)

Company:

Syngenta Canada (Apron Advance – PCP #30627, Apron Maxx RTA – PCP #27577, Apron Maxx RFC – PCP #28817)

Formulation:

Active ingredient:	Active Ingredient Formulation:		
	Apron Advance	Apron Maxx RTA	Apron Maxx RFC
Fludioxonil	25 g/L	0.73%	2.31%
Metalaxyl-M and S-isomer	20 g/L	1.10%	3.46%
Thiabendazole	150 g/L	–	–
Container sizes:	–	56.78L	56.78L

Crops and Diseases:

Crop:	Diseases Controlled by Apron Advance:	Diseases Controlled by Apron Maxx RTA and Apron Maxx RFC:
Chickpea	Seed rot, pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Rhizoctonia</i> spp. Seedling blights caused by <i>Pythium</i> and <i>Fusarium</i> spp.; Seed-borne ascochyta blight (<i>Ascochyta rubiei</i>); seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.	Seedling blights, damping-off, and seed rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp. Seed-borne ascochyta blight (<i>Ascochyta rubiei</i>); seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.
Lentil	Seed rot, pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Rhizoctonia</i> spp. Seedling root rot caused by <i>Fusarium</i> spp.; seed-borne ascochyta blight (<i>Ascochyta lentis</i>); seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.	Seedling blights, damping-off, and seed rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp. Seed-borne ascochyta blight (<i>Ascochyta lentis</i>); seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.
Dry pea	Seed rot, pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Rhizoctonia</i> spp. Seedling blights caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp.; seed-borne ascochyta and foot rot (<i>Ascochyta pinodes</i>)	Seedling blights, damping-off, and seed rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp. Seed-borne ascochyta and foot rot (<i>Ascochyta pinodes</i>)

Crop:	Diseases Controlled by <i>Apron Advance</i> :	Diseases Controlled by <i>Apron Maxx RTA</i> and <i>Apron Maxx RFC</i> :
Dry bean	Seed rot, pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Rhizoctonia</i> spp. Seedling blight caused by <i>Pythium</i> spp. Seed-borne anthracnose (<i>Colletotrichum</i> spp).	Seedling blights, damping-off, and seed rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp. Seed-borne anthracnose (<i>Colletotrichum</i> spp).
Lupin	Seed rot, pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Rhizoctonia</i> spp. Seed-borne anthracnose (<i>Colletotrichum</i> spp).	Seedling blights, damping-off, and seed rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp.
Fababean		
Soybean		Seedling blights, damping-off, and seed rots caused by <i>Pythium</i> , <i>Fusarium</i> , and <i>Rhizoctonia</i> spp. Seed rot and seedling blight caused by seed-borne <i>Phomopsis</i> spp. and early season root rot caused by <i>Phytophthora</i> spp.

Rates:

Apron Advance: 100 mL per 100 kg of seed.

Apron Maxx RTA: 325 mL per 100 kg of seed.

Apron Maxx RFC: 100 mL per 100 kg of seed.

Application Information:

Apron Advance is a seed treatment formulation for use in commercial seed treatment plants, and for on-farm treatment using auger treating only; do not use in hopper box or seed drill. *Apron Maxx RTA* is a ready-to-apply seed treatment formulation for use in commercial seed treatment plants and for on-farm treatment using standard gravity flow or mist type seed treatment equipment. Also used in treat-on-the-go air seeders. *Apron Maxx RFC* is a seed treatment formulation for use in commercial seed treatment plants, and for on-farm treatment using standard gravity flow or mist type seed treatment equipment which accurately meters, mixes and applies a flowable seed treatment. Also used in treat-on-the-go air seeders.

These products contain a pigment which will colour the treated seed. However, users are responsible for ensuring that the treated seed, when dried and ready for bagging, storage or seeding has an unnatural colour. If the pigment contained in the formulation does not colour the seed adequately, additional colourant must be added to the mixture while treating the seed. Ensure uniform coverage of the seed, as uneven seed coverage may not give the desired level of disease control. Treatment of highly damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. Allow the seed to dry before bagging, storing or seeding.

How it Works:

Fludioxonil is a phenylpyrrole fungicide with contact activity. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes, including *Pythium* damping-off. Thiabendazole is a benzimidazole fungicide with both contact and systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310. Experience has shown that strains of fungus resistant to metalaxyl-M may develop. Failure to control the disease will likely result in crop damage and/or yield losses. If disease appears in a treated field, consult the government extension specialist immediately.

Labelling: All seed treated with *Apron Maxx RTA* or *Apron Maxx RFC* must be labelled "This seed has been treated with fludioxonil and metalaxyl-M fungicides. DO NOT use for food, feed or oil purposes". All seed treated with *Apron Advance* must be labelled "This seed has been treated with thiabendazole, fludioxonil and metalaxyl-M and S-isomer fungicides. Do not use for food, feed or oil purposes".

Grazing: No restrictions listed.

Re-cropping: DO NOT plant any crop other than soybean, dry bean, chickpea, lentil or dry pea within 30 days to fields in which treated seed was planted.

Storage: Store away from feeds and feedstuffs. Store between 0 and 30°C. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up.

Environment: This product is toxic to fish and other aquatic organisms. Do not apply directly to aquatic habitats; do not contaminate water by cleaning of equipment or disposal of wastes. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned up.

Compatibility with rhizobia-based inoculants: These products are compatible with rhizobia-based inoculants. Check with inoculant manufacturer for details and refer to product labels prior to use. Mixing with inoculants may increase drying time while treating. Recalibrate the seed drill before planting treated seed.

Hazard Rating:

None listed.

Confine

Fungicide Group – 33

(Refer to page 312)

Company:

Winfield Solutions – PCP#29100

Canadian Agent: The Agronomy Company of Canada

Western Canada Distributor: Univar

Formulation:

45.8% mono and di-potassium salts of phosphorous acid

Container sizes - 9.46 L - 946.35 L.

Crops and Diseases:

Post-harvest treatment of potatoes for the suppression of late blight (*Phytophthora infestans*), pink rot (*Phytophthora erythroseptica*), and silver scurf (*Helminthosporium solani*) storage infection.

Rate and Application Information:

Dilute *Confine* at a 1:4.3 ratio with water (377 ml *Confine* + 1623 ml water). Apply 2 L of solution as a spray to 1000 kg of potatoes prior to storage. Ensure complete and even coverage.

How it Works:

The active ingredient mono- and di-potassium salts of phosphorous acid is a phosphonate fungicide with systemic activity to suppress pathogen inoculum. To be used as a preventative fungicide application on harvested tubers. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store this product away from food or feed.

Environment: DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of waste. DO NOT apply by air.

Hazard Rating:

None listed.

Crown

Fungicide Group – 1, 7

(Refer to page 312)

Company:

Manufactured by Chemtura Canada Co. – PCP# 23430

Formulation:

92 g per L carbathiin and 58 g per L thiabendazole, formulated as a liquid seed treatment. Container size - 10 L, 200 L, 1000 L.

Crops, Diseases and Rates:

Crop:	Diseases Controlled:	Application rates (mL per 100 kg seed):
Lentil	Seed-borne ascochyta caused by <i>Ascochyta lentis</i> ; seed rots and post-emergence damping off caused by <i>Botrytis</i> , <i>Fusarium</i> and <i>Rhizoctonia</i> .	300 to 600 mL *
Chickpea	Seed-borne ascochyta caused by <i>Ascochyta rabiei</i> .	300 to 600 mL **

* Use the higher rate to control seed-borne ascochyta in lentil.

** The lower rate will provide sufficient control of ascochyta in chickpea in most cases. Use the higher rate for smaller seed size varieties.

Application Information:

Commercial Treaters and On-Farm Auger Treating:

Crown is a ready to use formulation designed for commercial treaters and on-farm auger treating. *Crown* is added directly to the seed as it enters a mixing chamber or auger. It is important that the seed and chemical be mixed quickly and uniformly. See instructions supplied with the applicable treater system for information on proper application techniques. When a grain auger is used for treating, running the auger less than full is the key to adequate mixing. Augers used for handling treated seed should not be used to move seed for food, feed or oil processing.

Applications to Seed in a Hopper Box or Seed Drill:

Partially fill the hopper box or seed drill with a pre-measured amount of seed. Apply the proper amount of *Crown* evenly over the seed surface. DO NOT pour in one area. Mix with a paddle until all seed is of a uniform red colour, indicating adequate coverage. DO NOT MIX WITH HANDS. Repeat this procedure until the hopper box or seed drill is filled. Seed can be planted immediately after treatment without drying. Stir *Crown*-treated seed rigorously if the seeding operation has been interrupted for several hours or overnight.

How it Works:

The active ingredient carbathiin is a carboximide fungicide with systemic activity and the active ingredient thiabendazole is a benzimidazole fungicide with both contact and systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled "This seed has been treated with *Crown* seed protectant, which contains carbathiin and thiabendazole. DO NOT use for food, feed or oil processing."

Grazing: DO NOT graze or feed livestock on treated areas for 4 weeks after planting.

Re-cropping: No restrictions listed.

Storage: DO NOT store product above 35°C or below 0°C or in direct sunlight. Store in original container only, away from other pesticides, fertilizers, food or feed. Mix product prior to use by shaking the 10 L container.

Environment: Treated seed may be toxic to birds and other wildlife. Clean up any spilled seed. Ensure treated seed is properly incorporated at planting.

Compatibility with rhizobia-based inoculants: *Crown* is compatible with rhizobia and performs as a sticker for peat and granular inoculants. *Crown* and rhizobia can be applied simultaneously to seed through separate systems or applied sequentially. Read inoculant label before use.

Hazard Rating:

▽ Caution – Eye Irritant

For an explanation of the symbols used here see page 10

Cruiser 5FS

Insecticide Group – 4

(Refer to page 456)

Contains insecticide only. Available to commercial seed treaters only.

Company:

Syngenta Canada – PCP#27045

Formulation:

Cruiser 5FS: 47.6% thiamethoxam.
Container size - 23.4L and 56.78 L.

Crops, Insects and Rates:

Crops:	Insects Controlled:	Rate per 100 Kg of seed ¹
Wheat and Barley	Wireworms (suppression) ²	17 mL
	Wireworms (control) ²	33 to 50 mL
Corn	Seed corn maggot	83 to 166 mL
	Wireworms	83 mL
Soybean	Seed corn maggot	50 to 83 mL
	Wireworms, Soybean aphid	83 mL
Dry bean	Potato leafhopper ³ ; Seed corn maggot	50 to 83 mL
	Wireworms (suppression) ²	17 mL
	Wireworms (control) ²	33 to 50 mL
Pea, chickpea and lentil	Pea Leaf Weevil (pea only)	50 to 83 mL
	Wireworms (suppression) ²	17 mL
	Wireworms (control) ²	33 to 50 mL
Rye, millet, buckwheat, sorghum, and triticale	Wireworms (suppression) ²	17 mL
	Wireworms (control) ²	33 to 50 mL

¹ Use the higher rate for fields that have a history of moderate to severe insect pressure.

² Use lower rate for early season suppression of wireworms. However, if pressure is moderate to high or control is required, treat crops at higher control rate per 100 kg of seed.

³ Use the higher rate to replace one application of a foliar insecticide spray.

Application Information:

For use only in commercial seed treatment facilities with closed transfer systems that provide uniform seed coverage. *Cruiser 5FS* contains no colourant. A red colourant must be added when *Cruiser 5FS* is applied to grain. Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. Allow the seed to dry before bagging or storing in bulk containers.

How it Works:

Thiamethoxam is a seed treatment insecticide in the neonicotinoid class of chemistry that controls listed chewing and sucking insects through contact and systemic activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

For control of seed and soil-borne diseases, *Cruiser 5FS* can be mixed with fungicide seed treatments in a closed transfer system. Refer to label for details.

Restrictions:

Resistance management: Refer to page 455.

Labelling: All seed must be labelled "Seed treated with thiamethoxam insecticide. DO NOT use for food, feed or oil processing."

Grazing: DO NOT graze or feed livestock on treated areas for 45 days after planting.

Re-cropping: No restrictions listed.

Storage: Store away from food and feed. Ideal storage temperature is above freezing and below 30°C. If product should freeze, bring to room temperature, then ensure the contents are mixed well prior to application.

Environment: Products are toxic to fish and aquatic invertebrates. DO NOT apply directly to water or to areas where surface water is present. DO NOT contaminate food, feed, domestic or irrigation water supplies, lakes, streams and

ponds. If treated seed is spilled outdoors or accessible to birds, promptly clean up or bury to prevent ingestion.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Cruiser Maxx Beans/ Cruiser Maxx Vibrance Beans/ Cruiser Maxx Pulses

Available to commercial seed treaters only.

Cruiser Maxx Beans is a premix of *Cruiser 5FS* (page 396) and fludioxonil and metalaxyl-M (R- and S-isomers). *Cruiser Maxx Vibrance Beans* is a co-pack containing *Cruiser Maxx Beans* and *Vibrance 500FS* (page 446). *Cruiser Maxx Pulses* is a co-pack containing *Cruiser 5FS* (page 396) and *Apron Maxx RFC* (page 392). For more detailed information on the component products, consult individual pages in this guide as well as product labels.

Company:

Syngenta Canada

Formulation:

Cruiser Maxx Beans (PCP#28821) 22.6% thiamethoxam, 1.70% metalaxyl-M (R-and S-isomers), and 1.12% fludioxonil formulated as a liquid suspension seed treatment.

Cruiser Maxx Pulses: *Cruiser 5FS* (PCP#27045) 47.6% thiamethoxam formulated as a suspension and *Apron Maxx RTA* (PCP #27577) 0.73% fludioxonil plus 1.10% metalaxyl-M and S-isomers as a liquid seed treatment or *Apron Maxx*

Insecticide Group – 4
Fungicide Group – Cruiser
Maxx Pulses/Cruiser Maxx
Beans – 4, 12
Cruiser Maxx Vibrance Beans –
4, 7, 122

(Refer to page 312 and 456)

RTA (PCP#28817) 2.31% fludioxonil plus 3.46% metalaxyl-M formulated as a liquid seed treatment.

Cruiser Maxx Vibrance Beans: *Cruiser Maxx Beans* (PCP#28821) 22.6% thiamethoxam, 1.70% metalaxyl-M (R-and S-isomers), and 1.12% fludioxonil formulated as a suspension and *Vibrance 500FS* (PCP #30438) 500 g/L sedaxane formulated as a suspension.

Container size - *Cruiser Maxx Beans* 56.78 L, *Cruiser Maxx Vibrance Beans* – see sizes for *Cruiser Maxx Beans* and *Vibrance 500FS*, *Cruiser Maxx Pulses* – see sizes for *Cruiser 5FS* and *Apron Maxx RTA/Apron Maxx RFC*.

Crops, Diseases and Insects:

Product	Crop:	Diseases Controlled:	Insects Controlled:
Cruiser Maxx Vibrance Beans	Soybean	Seed decay, seedling blight and damping-off caused by <i>Rhizoctonia solani</i> and diseases controlled by Cruiser Maxx Beans.	See insects controlled by Cruiser Maxx Beans.
Cruiser Maxx Beans	Soybean ¹	Seed rot/pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp.; seedling blight caused by <i>Fusarium</i> spp., and <i>Pythium</i> spp.; seedling root rot caused by <i>Fusarium</i> spp.; seed rot and seedling blight caused by seed-borne <i>Phomopsis</i> spp.; early season root rot caused by <i>Phytophthora megasperma</i> var. <i>sojae</i> ²	Wireworm, seed corn maggot, European chafer, bean leaf beetle, and soybean aphid (early season protection)
	Dry Bean	Seed rot/pre-emergence damping-off, and post-emergence damping-off caused by <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Rhizoctonia</i> spp.; seedling blight caused by <i>Pythium</i> spp.; anthracnose caused by seed-borne <i>Colletotrichum</i> spp.	Wireworm, seed corn maggot, and potato leafhopper ³
Cruiser Maxx Pulses	Dry pea (including field pea)	Seed-borne ascochyta blight and foot rot caused by <i>Ascochyta pinodes</i> ; seed rot, damping-off, and seedling blight caused by <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., and <i>Fusarium</i> spp.	Pea leaf weevil and wireworm
	Lentil	Seed-borne ascochyta blight caused by <i>Ascochyta lentis</i> ; damping-off and seed rot caused by <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., and <i>Fusarium</i> spp.; seedling root rot caused by <i>Fusarium</i> spp.; seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.	Wireworm
	Chickpea	Seed-borne ascochyta blight caused by <i>Ascochyta rubiei</i> ; damping-off and seed rot caused by <i>Pythium</i> spp., <i>Rhizoctonia</i> spp., and <i>Fusarium</i> spp.; seedling blight caused by <i>Fusarium</i> spp., and <i>Pythium</i> spp.; seed rot and seedling blight caused by seed-borne <i>Botrytis</i> spp.	Wireworm

¹ Based on 6,600 soybean seeds per kg, Cruiser Maxx Beans delivers 85 µg of active ingredient (76 µg of thiamethoxam, 5.7 µg of metalaxyl-M and 3.8 µg of fludioxonil) per seed.

² Cruiser Maxx Beans provides early season protection against *Phytophthora* root rot for tolerant varieties of soybeans. If target fields have a history of high *Phytophthora* pressure, or susceptible varieties are to be treated, then tank-mix 195 mL of Cruiser Maxx Beans with 31 mL of Apron XL 1.5 per 100 kg of seed.

³ Replaces one application of a foliar insecticide spray.

Rate:

Cruiser Maxx Vibrance Beans: Cruiser Maxx Beans 195 mL + Vibrance 500FS at 5 to 10 mL per 100 kg of seed.

Cruiser Maxx Beans: 195 mL per 100 kg of seed.

Cruiser Maxx Pulses: Apron Maxx RTA 325 mL or Apron Maxx R/C 100 mL + Cruiser 5FS at 50 to 83 mL per 100 kg seed.

These should be tank mixed and applied by an approved commercial seed treating facility.

Application Information:

For use only in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment. No open transfer of seed treatments. All seed treated with this product must be conspicuously coloured at the time of treatment. Treatment of highly mechanically scarred or damaged seed or seed known

to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour.

How it Works:

Thiamethoxam is a seed treatment insecticide in the neonicotinoid class of chemistry that controls listed chewing and sucking insects through contact and systemic activity. For more information refer to "Insecticide Groups Based on modes of Action" on page 456. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes, including *Pythium* damping-off. The active ingredient fludioxonil is a phenylpyrrole fungicide with contact activity. The active ingredient Sedaxane is a succinate dehydrogenase inhibitor fungicide with systemic activity that inhibits fungal metabolism by binding to the succinate dehydrogenase enzyme to disrupt cellular respiration and energy generation. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Peas: When used at the low rate, *Cruiser Maxx Beans* may be tank-mixed with *Apron Maxx RTA*. Follow respective labels for diseases controlled and other instructions before tank-mixing.

Restrictions:

Resistance management: Refer to page 310.

Labelling: All seed must be labelled "This seed has been treated with the insecticide, thiamethoxam and metalaxyl M (including R- and S-isomer) and fludioxonil fungicides [and sedaxane fungicide if *Cruiser Maxx Vibrance Beans*]. DO NOT use for food, feed or oil processing."

Grazing: DO NOT graze or feed livestock on treated areas for 45 days after planting.

Re-cropping: DO NOT plant any crop other than soybeans, dry beans, chickpeas, lentils and dry peas (including field peas) within 45 days in which treated seeds were planted.

Storage: Store away from food and feed. Ideal storage temperature is above freezing and below 30°C. Repeated freeze-thawing of *Cruiser Maxx Beans* will not affect the physical integrity of the product. If product should freeze,

bring to room temperature, then ensure the contents are mixed well prior to application.

Environment: This product is toxic to fish and other aquatic organisms. DO NOT apply this product directly to aquatic habitats, estuaries or marine habitats. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. DO NOT contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

Compatibility with rhizobia-based inoculants: *Cruiser Maxx Beans* is compatible with rhizobium-based inoculants. Please check with inoculant manufacturers for details prior to use. Mixing with inoculants may increase drying time while treating extends the processing time. Recalibrate equipment before planting treated seed.

Hazard Rating:



Caution - Poison



Caution - Skin Irritant

For an explanation of the symbols used here see page 11.

Cruiser Maxx Vibrance Cereals/Cruiser Maxx Cereals/Cruiser Maxx Cereals Commercial

Cruiser Maxx Cereals Seed Treatment is a ready-to-use water-based formulation for on-farm seed treatment of small-grained cereals for disease and insect protection.

Cruiser Maxx Cereals Commercial is a ready-to-use water-based formulation for use in commercial seed treatment plants available to commercial seed treaters only.

Cruiser Maxx Vibrance Cereals OR *Cruiser Maxx Cereals Commercial* may be used with *Cruiser 5FS/350FS* seed treatment insecticide to achieve a high rate for control of wireworm under moderate to high insect pressure. This tank mix may be applied in COMMERCIAL SEED TREATMENT FACILITIES ONLY.

Insecticide Group - 4
Fungicide Group -
Cruiser Maxx Cereals/Cruiser
Maxx Cereals Commercial -
3, 4
Cruiser Maxx Vibrance
Cereals - 3, 4, 7

(Refer to page 312 and 456)

Company:

Syngenta Canada (*Cruiser Maxx Vibrance Cereals Seed Treatment* - PCP#30436; *Cruiser Maxx Cereals Seed Treatment* - PCP#29192; *Cruiser Maxx Cereals Commercial* - PCP#29127)

Formulation:

Cruiser Maxx Cereals Seed Treatment and *Cruiser Maxx Cereals Commercial*: 2.80% thiamethoxam, 3.36% difenoconazole, 0.56% metalaxyl-M (and S-isomer) formulated as a liquid suspension seed treatment.

Cruiser Maxx Vibrance Cereals Seed Treatment 30.7 g/L thiamethoxam, 36.9 g/L difenoconazole, 9.5 g/L metalaxyl-M (and S-isomer) and 8.0 g/L sedaxane formulated as a liquid suspension seed treatment.

Container size - 2x10L, 115 L, 450 L.

Crops, Diseases and Insects (Cruiser Maxx Cereals Seed Treatment, Cruiser Maxx Cereals Commercial, and Cruiser Maxx Vibrance Cereals):

Crop:	Diseases Controlled:	Diseases Suppressed ⁵	Insects:
Barley	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; covered smut (<i>Ustilago hordei</i>); false loose smut (<i>U. nigra</i>)	Common root rot (caused by <i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)	Suppression or control of wireworm ¹ (<i>Agriotes</i> , <i>Limonius</i> , and <i>Melanotus</i> spp.) and European chafer ² (<i>Rhizotrogus majalis</i>)
Winter Wheat	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>); dwarf bunt (<i>T. controversa</i>); loose smut (<i>Ustilago tritici</i>)		
Spring Wheat	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; common bunt ⁴ (<i>Tilletia tritici</i> , <i>T. laevis</i>); loose smut (<i>Ustilago tritici</i>)		
Rye	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; common bunt (<i>Tilletia caries</i>); dwarf bunt ⁴ (<i>T. controversa</i>)		Suppression or control of wireworm ¹ (<i>Agriotes</i> , <i>Limonius</i> , and <i>Melanotus</i> spp.)
Triticale	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; loose smut (<i>Ustilago tritici</i>)		
Oats	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; covered smut (<i>Ustilago kolleri</i>), loose smut (<i>U. avenae</i>)	Common root rot (<i>Cochliobolus</i> spp.)	Suppression of wireworm ⁵ (<i>Agriotes</i> , <i>Limonius</i> , and <i>Melanotus</i> spp.)

Additional Diseases (Cruiser Maxx Vibrance Cereals ONLY):

Crop:	Diseases Controlled:
Barley	Seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Rhizoctonia</i> spp. True loose smut (<i>Ustilago nuda</i>)
Winter Wheat	Seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Rhizoctonia</i> spp. Seed-borne <i>Septoria</i> (use 650 mL rate for control) <i>Septoria</i> leaf blotch (early season foliar disease control for first 4 weeks after planting; for full season control apply a foliar fungicide according to label directions)
Spring wheat	Seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Rhizoctonia</i> spp.
Oats	Seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Rhizoctonia</i> spp.

See footnotes on following page.

Additional Crops (Cruiser Maxx Cereals Seed Treatment, Cruiser Maxx Cereals Commercial ONLY):

Crop:	Diseases Controlled:	Insects:
Buckwheat	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i>	Wireworm (suppression) ¹ (<i>Agriotes</i> , <i>Limoni</i> us, and <i>Melanotus</i> spp.)
Millet (Pearl, Proso)	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i>	
Sorghum	General seed rots ³ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i>	

¹ These products provide suppression of wireworm activity at the 325 ml/100 kg seed rate; however, if pressure is moderate to high or control is required, use the 650 mL/100 kg seed rate to achieve 20 g thiamethoxam per 100 kg seed. Commercial seed treaters may tank mix Cruiser Maxx Cereals Commercial OR Cruiser Maxx Vibrance Cereals with Cruiser 5FS/350FS Seed Treatment insecticide to achieve a total use rate of 20 to 30 g of thiamethoxam per 100 kg seed. Consult labels for use rates and instructions.

² For control of European chafer activity on wheat and barley, Cruiser Maxx Cereals Commercial OR Cruiser Maxx Vibrance Cereals may be mixed with Cruiser 5FS/350FS Seed Treatment Insecticide to achieve a total use rate of 30 g of thiamethoxam per 100 kg seed. Consult labels for use rates and instructions. This tank mix may be applied in commercial seed treatment facilities ONLY.

³ General seed rots controlled include those caused by saprophytic organisms such as *Fusarium*, *Pythium*, *Penicillium* and *Aspergillus*.

⁴ Controls both seed- and soil-borne bunts (common, dwarf).

⁵ Suppression means consistent control at level which is not optimal but is still of commercial benefit.

⁶ When seed is treated for post-planting protection against registered pests, these products will also provide protection during post-treatment storage of the seed against damage from many storage insect pests. Seed treated with thiamethoxam has been tested and found to be effective against rusty grain beetle, saw-toothed grain beetle, red flour beetle, rice weevil, lesser grain borer, European corn borer and Indian meal moth. It is recommended that seed with existing populations of storage pests be fumigated prior to treating and storage of that seed.

Rate:

325 mL or 650 mL per 100 kg of seed. Use the higher rate if: a) There is a history of high disease pressures in the field, or b) where conditions favour seed-borne and soil-borne pathogens, or c) when controlling seed-borne *Septoria* or early season *Septoria* leaf blotch, or d) if control of wireworm is required or if wireworm populations are expected to be moderate to high; or e) for Cruiser Maxx Vibrance Cereals only, where high levels of seed-borne infections like smuts are high.

Application Information:

All seed treated with this product must be conspicuously coloured at the time of treatment. Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour.

Cruiser Maxx Cereals Commercial is for use only in commercial seed treatment facilities.

How it Works:

Thiamethoxam is a seed treatment insecticide in the neonicotinoid class of chemistry that controls listed chewing and sucking insects through contact and systemic activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456. The active ingredient difenoconazole is a triazole fungicide with broad-spectrum, systemic activity. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes, including *Pythium* damping-off. The active ingredient Sedaxane is a succinate dehydrogenase inhibitor fungicide with systemic activity that inhibits fungal metabolism by binding to the succinate dehydrogenase enzyme to disrupt cellular respiration and energy generation. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Cruiser Maxx Cereals Commercial or Cruiser Maxx Cereals Vibrance may be mixed with 325 mL of Dividend XL RTA if a) there is a history of high disease pressures in the field,

b) where field conditions favour seed-borne and soil-borne pathogens, or c) when controlling seed-borne *Septoria*. Consult each product and follow the most restrictive label precautions and limitations.

Restrictions:

Resistance management: Refer to page 310.

Thiamethoxam is a Group 4 Insecticide. Do not make any subsequent application of a Group 4 Insecticide (i.e., in-furrow or foliar application) following treatment with these products.

Experience has shown that strains of fungus resistant to metalaxyl-M (and S-isomer) may develop. Failure to control the disease will result in crop damage and/or yield losses. If disease appears in a treated field, consult the government extension specialist immediately.

Labelling: All seed must be labelled "This seed has been treated with the insecticide, thiamethoxam and difenoconazole and metalaxyl M (and S-isomer) fungicides [and sedaxane fungicide if *Cruiser Maxx Vibrance Cereals*]. DO NOT use for food, feed or oil processing."

Grazing: DO NOT graze or feed livestock on treated areas for 45 days after planting.

Re-cropping: No restrictions listed.

Storage: Store away from food and feed. Ideal storage temperature is above freezing and below 30°C. Repeated freeze-thawing of product will not affect the physical integrity of the product. If product should freeze, bring to room temperature, and then ensure the contents are mixed well prior to application.

Environment: Toxic to aquatic organisms. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. DO NOT contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

Hazard Rating:



Caution – Poison



Warning – contains the allergen sulfites.

For an explanation of the symbols used here see page 11.

Cruiser Maxx D Potatoes

Cruiser Maxx D Potatoes is a co-pack of *Maxim D Seed Treatment Fungicide* (fludioxonil and difenoconazole, page 421) and *Actara 240SC Insecticide* (thiamethoxam, page 498). Information listed is limited to crop, diseases, insects and rates. For other detailed information on the component products see the product pages listed above.

**Insecticide Group – 4A
Fungicide Group – 12**

(Refer to page 312 and 456)

Company:

Syngenta Canada

Formulation:

Cruiser Maxx Potatoes has two components:

Maxim D (PCP#30599): 19.4 g/L fludioxonil and 19.4 g/L difenoconazole formulated as a suspension.

Actara 240SC (PCP#28407): 240 g per L thiamethoxam formulated as soluble concentrate.

See the component products for other information. Use the most stringent restrictions for either product.

Crops, Diseases, and Insects:

Maxim D: Seed piece treatment fungicide for use on potato for the control of black scurf/stem and stolon canker (*Rhizoctonia solani*), silver scurf (*Helminthosporium solani*), and fusarium dry rot (*Fusarium* spp.).

Actara 240 SC: Seed piece treatment insecticide for use on potato for the control of Colorado potato beetle, aphids, potato leafhopper.

Rate and Application Information:

Maxim Liquid PSP: 130 mL per 100 kg of seed for control of black scurf/stem and stolon canker and silver scurf; 65-130 mL per 100 kg of seed for control of fusarium dry rot (use the high rate when disease pressure and/or resistance pressure is high).

Actara 240 SC: Choose the appropriate rate from the chart on label, based on seeding rate.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

DB-Red L

Fungicide Group – M3
(Refer to page 312)

Company:

Loveland Products Inc., distributed by United Agri Products – PCP# 27144

Formulation:

323 g per L maneb formulated as a liquid.
Container size - 9 L to 57 L.

Crops, Diseases and Rates:

Crop:	Diseases Controlled:	Rate per 100 Kg seed:
Wheat	Common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>), Root rot, and seedling blight (including <i>Fusarium</i>)	312 mL
Barley	Covered smut (<i>Ustilago hordei</i>), false loose smut (<i>U. nigra</i>), root rot, and seedling blight	396 mL
Oat	Covered smut (<i>Ustilago kolleri</i>), root rot, and seedling blight	552 mL
Rye	Common bunt (<i>Tilletia caries</i>), root rot, and seedling blight	260 mL

Application Information:

Designed as a ready-to-use seed treatment for on-the-farm and commercial use with a seed treater suitable for metering and mixing flowable seed treatments. Use only the recommended rate, lower amounts may not give the desired control. Higher amounts may cause seed injury. Uneven treating of seed may cause over treatment on some seed kernels and under treatment on other seed kernels. Seed should be well cured, dry and cleaned before treatment. For maximum benefits, avoid deep seeding and

exceptionally early sowing under poor growing conditions. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Dispose of all excess treated seeds by seeding away from bodies of water.

How it Works:

The ingredient maneb is a dithiocarbamate fungicide with multi-site contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: If treated seed is stored, label container "This seed has been treated with DB-Red L seed treatment. DO NOT use for food, feed or processing."

Grazing: DO NOT graze or feed livestock on treated areas.

Re-cropping: No restrictions listed.

Storage: Store away from heat source and in a cool, dry area. If product becomes frozen, thaw and shake or agitate. Excess treated seed should not be stored past planting time. DO NOT contaminate feed or food stuffs.

Environment: This product is toxic to fish, birds and other wildlife. DO NOT contaminate ponds, lakes or streams.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

DCT

Note: This product is being discontinued; will no longer be available for use after December 31, 2014.

Fungicide Group – 1,M4 Insecticide Group – 1B

(Refer to page 312 and 456)

Company:

Norac Concepts Inc. – PCP#14986

Formulation:

6% diazinon, 18% captan, and 14% thiophanate-methyl formulated as a water soluble powder.

Container size - 0.4 kg, 10 kg.

Crops and Diseases:

Field bean: Control of seedling blight, root rot, seed-borne anthracnose.

This product will not control anthracnose if seed is severely infected.

Sweet corn: Control of seed borne *Penicillium oxalicum* and *Penicillium* spp.

Insects:

Control of root maggots on beans and sweet corn.

Rates:

For slurry machines: 520 g per 1 L water. Use 1 L of slurry per 100 kg of seed.

For mixing in bucket: 130 g per 0.35 L water. Use 0.35 L of slurry per 25 kg of seed. Mix well to keep powder suspended in water. Dry the seed before seeding or bagging.

For Sweet corn: 125 g of product per 25 kg of seed.

Application Information:

May be applied as a water-based slurry through standard slurry or commercial seed treating equipment. For best results, seed should be planted within a week after treating. This product will not control anthracnose if seed is severely infected. Treatment of mechanically scarred or damaged seed, or of seed known to be of low germ, low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment.

How it Works:

The active ingredient captan is a phthalimide fungicide with multi-site protective activity. The active ingredient thiophanate-methyl is a methyl-benzimidazole carbamate fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311. The diazinon component is an organophosphate insecticide with contact and stomach activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed should be coloured and labelled "Poisonous to man and animals. Seed treated with diazinon, captan and thiophanate-methyl for control of insects and seed-borne diseases. DO NOT use for food or feed purposes."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: DO NOT store treated seed for more than one month before planting.

Environment: DO NOT contaminate any body of water.

Hazard Rating:



Danger – Poison



Warning – contains the allergen soy.

For an explanation of the symbols used here see page 11.

Dividend XL RTA

Fungicide Group – 3, 4
(Refer to page 312)

Company:

Syngenta Canada – PCP# 25777

Formulation:

3.37% difenoconazole, 0.27% metalaxyl-M formulated as a flowable seed treatment.

Container size - 2 x 10 L, 115 L and 450 L.

Crops, Diseases, and Rates:

Crop:	Rate per 100 Kg seed: ¹	Diseases Controlled:	Diseases Suppressed ⁴ :
Barley	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off; seed-borne <i>Septoria</i> ³ ; false loose smut (<i>Ustilago nigra</i>); covered smut (<i>U. hordei</i>)	Common root rot (<i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)
Corn	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off	—
	325 mL	Penicillium three leaf dieback	
Oat	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off; covered smut (<i>Ustilago kolleri</i>); loose smut (<i>U. avenae</i>)	Common root rot (<i>Cochliobolus</i> spp.)
Rye	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off; seed-borne <i>Septoria</i> ³ ; common bunt ⁶ (<i>Tilletia caries</i>); dwarf bunt ⁶ (<i>T. controversa</i>)	Common root rot (<i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)
Triticale	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off; loose smut (<i>Ustilago tritici</i>)	Common root rot (<i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)

See footnotes on following page.

Continued...

Crops, Diseases, and Rates *continued*:

Crop:	Rate per 100 Kg seed: ¹	Diseases Controlled:	Diseases Suppressed ⁴ :
Spring Wheat	325 to 650 mL	General seed rots ² ; seedling blight and root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off; seed-borne <i>Septoria</i> ³ ; common bunt ⁶ (<i>Tilletia tritici</i> , <i>T. laevis</i>); loose smut (<i>Ustilago tritici</i>)	Common root rot (<i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)
Winter Wheat	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off; seed-borne <i>Septoria</i> ³ ; <i>Septoria</i> leaf blotch ^{3,5} ; common bunt ⁶ ; (<i>Tilletia tritici</i> , <i>T. laevis</i>); dwarf bunt ⁶ (<i>T. controversa</i>); loose smut (<i>Ustilago tritici</i>)	Common root rot (<i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)
Buckwheat, millet, sorghum	325 to 650 mL	General seed rots ² ; seedling blight and seedling root rot caused by seed- and soil-borne <i>Fusarium</i> and soil-borne <i>Pythium</i> ; <i>Pythium</i> and <i>Fusarium</i> damping-off	—

¹ Use the higher rate where conditions favour seed- or soil-borne pathogens or if field has history of high disease pressure.

² General seed rots controlled include those caused by saprophytic organisms such as *Penicillium*, *Aspergillus*, *Fusarium*, and *Pythium*. Use the 325 mL rate for control of these diseases.

³ Use the 650 mL rate for control of these diseases.

⁴ Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

⁵ Early season foliar disease control for first 4 weeks after planting. For full season control apply a foliar fungicide according to label directions.

⁶ Controls both seed- and soil-borne bunts (common, dwarf).

Application Information:

Dividend XL RTA is a ready-to-apply formulation for use in commercial seed treatment plants, and for on-farm treatment using standard gravity flow or mist-type seed treatment equipment which accurately meters and mixes a flowable seed treatment. *Dividend XL RTA* may also be used in a treat-on-the-go air seeder. The equipment must provide uniform coverage of product on the seed. Consult the manufacturer of the application equipment for suitability for this application and for instructions on operation and calibration of the equipment.

Uneven seed coverage may not give the desired level of disease control.

Water Volume: *Dividend XL RTA* does not require addition of water for application. However, when using the Flexi-Coil Seed Treatment Unit, *Dividend XL RTA* must be diluted with water to reach 99 mL per 10 kg of seed. The recommended dilution rate is 1 part *Dividend XL RTA* to 2 parts water. This is equal to 33 mL of *Dividend XL RTA* plus 66 mL of water in order to achieve the total liquid volume requirement of 99 mL per 10 kg seed.

How it Works:

The active ingredient difenoconazole is a triazole fungicide with broad-spectrum, systemic activity. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes class, including *Pythium* damping-off. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

For the control of true loose smut (*Ustilago nuda*) in barley, tank mix *Dividend XL RTA* with *Charter*, *Raxil 250FL* or *Baytan 30*. Read the label directions for each product and follow the more restrictive label precautions and limitations.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed should be labelled "This seed has been treated with *Dividend XL RTA*; DO NOT use for food, feed, or oil purposes."

Grazing: DO NOT graze, feed green forage or cut for hay within 35 days of planting treated cereal grain seeds.

Re-cropping: DO NOT plant any crop other than cereal grains listed on this label within 30 days to fields in which treated seeds were planted.

Storage: Keep in heated storage. Product will freeze at -18°C. Store away from feeds and food stuffs. DO NOT carry-over treated seed.

Environment: This product is toxic to fish and other aquatic invertebrates. DO NOT apply directly to water or to areas where surface water is present. DO NOT contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. If treated seed is spilled outdoors or in areas accessible to birds, promptly clean up or bury to prevent ingestion.

Hazard Rating:
None listed.

EverGol Energy

Fungicide Group – 3, 4, 7
(Refer to page 312)

Company:

Bayer CropScience – PCP#30364

Formulation:

38.4 g per L penflufen, 76.8 g per L prothioconazole, 61.4 g per L metalaxyl formulated as a suspension.

Container size - 33.75 L.

Crops, Diseases, and Rates:

Crop:	Rate per 100 Kg seed: ¹	Diseases Controlled:	Diseases Suppressed:
Wheat, barley, oat, rye, triticale, pearl millet, and proso millet	65 mL	Seed rot/pre-emergence damping-off caused by seed-borne <i>Fusarium</i> spp., <i>Aspergillus</i> spp. and <i>Cochliobolus sativus</i> ; seed rot/pre-emergence damping-off caused by soil-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and <i>Pythium</i> spp.; post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp. and seed-borne <i>Aspergillus</i> spp.; seedling blight caused by seed-borne <i>Aspergillus</i> spp.; true loose smut (<i>Ustilago nuda</i>); covered smut (<i>U. hordei</i> , <i>U. koleri</i>); false loose smut (<i>U. nigra</i>); loose smut (<i>U. tritici</i> , <i>U. avenae</i>); common bunt (<i>Tilletia caries</i> , <i>T. laevis</i>); leaf stripe (<i>Pyrenophora graminea</i>)	Root rot caused by soil-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and seed-borne <i>Fusarium</i> spp.; crown rot caused by seed-borne <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Penicillium</i> spp.
Corn	65 mL	Seed rot/pre-emergence damping-off caused by <i>Rhizoctonia solani</i> , soil-borne <i>Fusarium</i> spp. and <i>Pythium</i> spp., seed-borne <i>Fusarium</i> spp., <i>Cladosporium</i> spp., and <i>Aspergillus</i> spp.; post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp.	Seed rot/pre-emergence damping-off caused by <i>Penicillium</i> spp.
Soybean, bean, pea, lentil	65 mL	Seed rot/pre-emergence damping off caused by <i>Rhizoctonia solani</i> , <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Phomopsis longicolla</i> ; post-emergence damping-off caused by <i>R. solani</i> , <i>Fusarium</i> spp., and <i>Pythium</i> spp.; early-season root rot and seedling blight caused by <i>R. solani</i> and <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Botrytis cinerea</i>	

Continued...

Crops, Diseases, and Rates *continued*:

Crop:	Rate per 100 Kg seed: ¹	Diseases Controlled:	Diseases Suppressed:
Chickpea	65 mL	Seed rot/pre-emergence damping off caused by <i>Rhizoctonia solani</i> , <i>Fusarium</i> spp., <i>Pythium</i> spp., and <i>Phomopsis longicolla</i> ; post-emergence damping-off caused by <i>R. solani</i> , <i>Fusarium</i> spp., and <i>Pythium</i> spp.; early-season root rot and seedling blight caused by <i>R. solani</i> and <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Botrytis cinerea</i>	Seed rot/pre-emergence damping-off caused by seed-borne <i>Ascochyta rubiei</i>

Application Information:

EverGol Energy is designed for commercial or on-farm treating with conventional seed treating equipment which can accurately control application rates and provide a good distribution of the chemical into the seed in the mixing chamber. This product is recommended to be diluted with water or another suitable liquid just prior to application to ensure uniform coverage on the seed during the application process. Uniform application to seed is necessary to ensure optimum product performance. This product contains no dye and an appropriate seed colourant must be applied.

How it Works:

The active ingredient penflufen is a carboxamide (SDHI) fungicide with systemic activity. The active ingredient prothioconazole is a demethylation inhibitor with broad-spectrum systemic activity. The active ingredient metalaxyl is an acylalanine fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Insecticide Seed Treatments: *Stress Shield 600* in wheat, barley, oat, and soybean; *Poncho 600 FS* in corn

Fungicide Seed Treatment: *Allegiance FL* for control of early-season *Phytophthora* in soybean

Follow the label directions for each product and use the most restrictive precautions and limitations for either product.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labeled "This seed has been treated with *EverGol Energy*, which contains penflufen, prothioconazole and metalaxyl. When handling treated seed wear a long-sleeved shirt, long pants and chemical-resistant gloves. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs."

Grazing: No restrictions listed

Re-cropping: Registered crops for *EverGol Energy*, as well as canola, mustard, rapeseed, borage, flax, crambe and potato, may be replanted at any time. For all other crops, do not plant-back within 30 days of seeding with *EverGol Energy*-treated seed.

Storage: To prevent contamination store this product away from food or feed. Store in cool, dry area. DO NOT store in direct sunlight. DO NOT allow prolonged storage in temperatures that exceed 40°C or go below -10°C.

Environment: Toxic to aquatic organisms and non-target terrestrial plants. DO NOT discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans or other water. Dispose of all excess treated seed. Left over seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. DO NOT leave exposed treated seed on soil surface. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface. The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or depth to the water is shallow.

Hazard Rating:

None listed.

Gaucha CS FL

Gaucha CS FL is available to commercial seed treaters only.

Fungicide Group – 7, M3
Insecticide Group – 4A
(Refer to page 312 and 456)

Company:

Bayer CropScience – PCP# 27174

Formulation:

285.7 g per L imidacloprid, 47.6 g per L carbathiin, 95.3 g per L thiram formulated as a suspension.

Container sizes - 10, 100, 1000 L.

Crops, Diseases, Insects and Rates:

Crop:	Rate (ml) per 100 Kg seed ¹	Insects Controlled ² :	Diseases Controlled:
Canola, rapeseed, and mustard (including oil-seed mustard)	1400 to 2100 mL	Early season control of flea beetles.	Seed rot, damping off, seedling blight and early season root rot caused by <i>Rhizoctonia</i> , <i>Pythium</i> and <i>Alternaria</i> spp.; Seed-borne blackleg (<i>Leptosphaeria maculans</i>) ³

¹ In areas where flea beetle populations are often high, use the higher rates.

² Under high insect pressure, a foliar insecticide may also be required. Monitor crop regularly for insect infestation levels.

³ In canola and rapeseed only.

Application Information:

For use in commercial seed treaters only. Seed treatment must be thoroughly agitated to ensure uniform mixing of product prior to and during application. DO NOT apply direct heat to container. These products DO NOT contain colourant. A blue colourant must be added when products are applied to oilseeds. Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour.

How it Works:

Imidacloprid is a chloronicotynyl insecticide with systemic activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456. Carbathiin is a carboximide fungicide with systemic activity and thiram is a dithiocarbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled as follows: "This

seed has been treated with Gaucha CS FL seed protectant, which contains imidacloprid, carbathiin and thiram. Do not use for food, feed or oil processing. Store away from feeds and other foodstuffs. Wear a long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seeds."

Grazing: DO NOT graze or feed livestock on treated areas for 4 weeks after planting.

Re-cropping: No restrictions listed.

Storage: Protect products from freezing. Keep above 10°C prior to and during application. DO NOT store in direct sunlight or above 35°C. Treated seed stored for periods in excess of 9 months should be tested for germination before planting. DO NOT store treated seed above 25°C or in direct sunlight.

Environment: These products are highly toxic to birds and aquatic invertebrates. DO NOT apply directly to water or to areas where surface water is present. DO NOT contaminate water when disposing of equipment wash water. Cover or incorporate spilled treated seeds. Leftover treated seed should be double sown around the headlands, or buried away from water sources.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Gemini

Fungicide Group – 3, M4

(Refer to page 312)

Company:

BASF Canada – PCP# 27826

Formulation:

1.25% triticonazole and 12.5% thiram formulated as a liquid flowable seed treatment. Container size – 2 x 6 L; 200 L.

Crops, Rates and Diseases:

Crop:	Rate (ml) per 100 Kg seed	Diseases Controlled:	Diseases Suppressed*:
Wheat	360 mL	Seed rot caused by <i>Fusarium</i> spp.; seedling blights caused by seed and soil-borne <i>Fusarium</i> spp.; loose smut (<i>Ustilago tritici</i>); common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>); <i>Pythium</i> damping off	<i>Fusarium</i> crown and root rot; seedling blight and common root rot caused by <i>Cochliobolus</i> sp.
Barley	360 mL	Seed rot caused by <i>Fusarium</i> spp.; seedling blights caused by seed and soil-borne <i>Fusarium</i> spp.; true loose smut; (<i>Ustilago nuda</i>); covered smut (<i>U. hordei</i>); false loose smut (<i>U. nigra</i>); <i>Pythium</i> damping off	<i>Fusarium</i> crown and root rot; seedling blight and common root rot caused by <i>Cochliobolus</i> sp.
Oats	360 mL	Loose smut (<i>Ustilago avenae</i>), covered smut (<i>U. kolleri</i>)	—

* Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

Application Information:

Gemini is a ready-to-use seed treatment for use in commercial seed treatment plants and for use in on-farm standard gravity flow or mist type treatment machines. *Gemini* can also be used in on-the-go air seeder treatment systems. Agitate or shake well prior to usage.

Water volume: *Gemini* does not require the addition of water for application. When using the Flexi-Coil "on-the-go" seed treatment unit, please contact BASF AgSolutions at 1-877-371-2273 for details on product rates.

How it Works:

The active ingredient triticonazole is a triazole fungicide with broad-spectrum systemic activity. Thiram is a dithiocarbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled "This seed is treated with *Gemini* seed treatment fungicide containing

triticonazole and thiram. DO NOT use for food, feed or oil processing".

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store in original container with lid tightly closed. Store away from children, animals, food, feed stuffs, fertilizers and seed. Protect from frost and freezing. DO NOT store treated seed for more than 18 months. Store treated seed in cool, dry conditions.

Environment: DO NOT feed treated seed to, or otherwise expose, wildlife or domestic birds. DO NOT contaminate domestic or irrigation water supplies, lakes, streams, ponds or any body of water with the chemical, used containers, treated seed or bags. DO NOT contaminate water by cleaning of equipment or disposal of wastes.

Hazard Rating:



Caution – Poison



Caution – Eye irritant and potential skin sensitizer

For an explanation of the symbols used here see page 11.

General Storage Disinfectant

Company:

Ag-Services Inc (Distributed by United Agri Products) –
PCP#14957

Formulation:

10% dimethyl benzyl ammonium chloride, formulated as a liquid. Container size - 20 L.

Crops:

Use for disinfecting storage areas and equipment. Not for direct use on potatoes.

Diseases:

Control of bacterial ring rot and other decay organisms in potato storage.

Rate:

60 mL per 10 L water.

Application Information:

Apply only when storage areas are empty. To disinfect walls and floors of potato storages, clean surfaces thoroughly with a broom or vacuum to remove all dirt and debris. Wash with detergent prior to disinfecting. Then, wash, mop or spray thoroughly with disinfectant solution (60 mL disinfectant in 10 L of water). Allow treated surfaces to remain wet for at least 20 minutes. The same strength solution can be used to disinfect used bags, potato planters and other machinery after all dirt has been removed. Soak bags for at least 1 hour.

Equipment: All handling and planting equipment should be cleaned and treated on a regular basis (daily when preparing seed and seed pieces). Dirt should be removed through washing with detergent prior to disinfection. Treat equipment by mopping and brushing methods.

Storage walls and ceilings: Use 600 mL of disinfectant in 100 L of water. Wash areas with detergent prior to disinfecting. Spray areas using a high pressure jet (up to 4250 kPa pressure) to penetrate cracks, etc. in floors. Spray storage air ducts with a solution of 1.2 L of disinfectant in 100 L of water. Sub-surface air ducts, flumes and plenums should be thoroughly cleaned prior to disinfection.

Tank Mixes:

DO NOT mix with soaps, detergents, foaming agents or surfactants.

Hazard Rating:



Caution – Corrosive

Other precaution: Corrosive, causes severe eye and skin damage. DO NOT get in eyes, on skin or on clothing. Avoid contamination of food. DO NOT breathe mist of diluted chemical created from pressure washer applications. Wear gloves, goggles, rubber boots, wet suit, and mist respirator when using pressure washer system.

For an explanation of the symbols used here see page 11.

Heads Up Plant Protectant

Fungicide Group – NC
(Refer to page 312)

Company:

Heads Up Plant Protectants, Inc.– PCP# 29827

Distributed by Engage Agro Corporation

Formulation:

63.02% saponins of *Chenopodium quinoa* formulated as a soluble powder.

Container size - 25 g pouches

Crops and Diseases:

Crop:	Diseases Suppressed:
Potato (cut or whole tubers)	Rhizoctonia canker and black scurf (<i>Rhizoctonia solani</i>)
Soybean	Root rot and post-emergence damping-off (<i>Rhizoctonia solani</i>); white mould (<i>Sclerotinia sclerotiorum</i>); sudden death syndrome (<i>Fusarium virguliforme</i>)
Dry bean	White mould (<i>Sclerotinia sclerotiorum</i>)

Rate Information:

Mix 1 gram of product per 1 L of water. Apply 1 L of solution for every 100-264 kg of potato seed or for every 163 kg of soybean or dry bean seed.

Application Information:

Treat soybean or dry bean seed by dipping, spraying or dribbling the solution into a rotation auger conveyor or some other approved seed treatment device. Spray application to seeds within an enclosed spray device to ensure thorough coverage.

For seed potatoes, product must be applied to germination seed potatoes, as indicated by obvious sprouting activity coming from potato eyes. This sprouting activity can be from peeking to full sprout length, but before green leaves appear.

How it Works:

The active ingredient saponins of *Chenopodium quinoa* is made from plant sources. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store this product away from food or feed. Store above -12°C.

Environment: DO NOT contaminate irrigation or drinking water supplies or aquatic habitats. This product is toxic to aquatic organisms.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Helix Vibrance

Available to commercial seed treaters only.

Fungicide Group – 3, 4, 12, 7
Insecticide Group – 4A

(Refer to page 312 and 456)

Helix Vibrance is a co-pack of **Helix Xtra** (thiamethoxam insecticide, and difenoconazole, metalaxyl-M, and fludioxonil fungicides, page 413) and **Vibrance 500FS** (sedaxane, page 446). Information listed is limited to crop, diseases, insects and rates. For other detailed information on the component products see the product pages listed above.

Company:

Syngenta Canada

Formulation:

Helix Xtra (PCP #26638): 20.7% thiamethoxam, 1.25% difenoconazole, 0.39% metalaxyl-M, 0.13% fludioxonil formulated as a liquid seed treatment.

Vibrance 500FL (PCP #30438): 500 g/L sedaxane formulated as a suspension.

Container size - 105 L to Bulk.

Crops:

Canola, Oriental mustard (both oilseed and condiment types).

Diseases Controlled:

Seed-borne blackleg (*Leptosphaeria maculans*), seed-borne *Alternaria*, and the seedling disease complex (damping off, seedling blight, seed rot, root rot) caused by *Pythium*, *Fusarium* and *Rhizoctonia* spp. Early season control of flea beetles (28 to 35 days).

Rate:

1.5 L **Helix Xtra** and 5 to 10 mL **Vibrance 500FS** per 100 kg seed.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Helix Xtra

Available to commercial seed treaters only.

Fungicide Group – 3, 4, 12
Insecticide Group – 4A

(Refer to page 312 and 456)

Company:

Syngenta Canada – PCP#26638

Formulation:

20.7% thiamethoxam, 1.25% difenoconazole, 0.39% metalaxyl-M, 0.13% fludioxonil formulated as a liquid seed treatment.

Container size - 105 L to Bulk.

Crops:

Canola, Oriental mustard (both oilseed and condiment types).

Diseases Controlled:

Seed-borne blackleg (*Leptosphaeria maculans*), seed-borne *Alternaria*, and the seedling disease complex (damping off, seedling blight, seed rot, root rot) caused by *Pythium*, *Fusarium* and *Rhizoctonia* spp.

Insects Controlled:

Early season control of flea beetles (28 to 35 days).

Rate:

1.5 L per 100 kg seed.

Application Information:

For use only in commercial seed treatment facilities with closed transfer systems. *Helix Xtra* is a ready to use liquid product and contains an appropriate colourant. Apply using standard commercial seed treatment equipment that provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect or disease control. Consult the manufacturer of the seed treating equipment for advice on the operation and calibration of the equipment. Maintain constant agitation during the seed treatment process. Allow the seed to dry before bagging.

How it Works:

The active ingredient thiamethoxam is a systemic insecticide from the neonicotinoid chemical class. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456. The active ingredient difenoconazole is a triazole fungicide with broad-spectrum systemic activity. The active ingredient metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes class, including Pythium damping off. The active ingredient fludioxonil is a phenylpyrrole chemistry and has contact and locally systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 399.

Labelling: Treated seed must be labelled "This seed has been treated with *Helix Xtra* which contains insecticide (thiamethoxam) and fungicides (difenoconazole, metalaxyl-M and S-isomer, and fludioxonil). Wear long-sleeve shirt, long pants, and chemical-resistant gloves when handling treated seed. Do not use for food, feed or oil processing. Store away from food and feed."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store in a well-ventilated, secure area. Store away from food and feed.

Environment: DO NOT apply directly to water, or to areas where surface water is present. In cleaning of equipment or disposing of wastes, DO NOT contaminate water used for human or animal consumption or by wildlife and aquatic life or for irrigation purposes. If treated seed is spilled outdoors, promptly clean up.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Imidacloprid

Insecticide Group – 4A
(Refer to page 456)

Admire SPT / Alias 240 SC / Sombrero 600 FS

Company:

Bayer CropScience (*Admire SPT* – PCP#27702)
MANA Canada (*Alias 240 SC* – PCP#28475 and *Sombrero 600 FS* – PCP#30505)

Formulation:

Admire and *Alias*: 240 g per L imidacloprid.
Container sizes - 1 L, 3.78 L. Contains insecticide only.
Sombrero: 600 g per L imidacloprid.
Contains insecticide only.

Crops and Insects:

Product:	Crop:	Rate:	Insects Controlled:
<i>Admire</i> and <i>Alias</i>	Potato	26 to 39 ml per 100 kg of potato seed tubers. The higher rate is recommended when extended length of control is needed. DO NOT apply more than 0.47 L per acre per year.	Colorado potato beetle, potato flea beetle, potato leafhopper and aphids (including green peach, buckthorn, foxglove and potato aphid)
<i>Sombrero</i>	Canola, mustard (condiment-type only) and rapeseed	667 ml per 100 kg seed to 1333 mL per 100 kg seed.	Flea beetles
	Corn	21.3 mL product per 80,000 seeds.	Corn flea beetle
		80 mL product per 80,000 seeds.	Wireworms

Application Information:

May be applied when potato pieces are being cut. Apply specified dosage as a diluted spray onto seed-pieces using a shielded spray system that is well contained and will prevent the loss of any liquid. DO NOT dilute with any more than 3 parts water to 1 part *Admire SPT/ Alias* 240 SC. DO NOT dilute *Sombrero* beyond 6%. Agitate or stir spray solution as needed. Complete coverage of the seed piece is required for optimal insect control. As part of the seed cutting and treating process, application of a fungicide registered for potato seed treatment or an inert absorbent ingredient is recommended.

NOTE: A colourant must be added to *Sombrero* to colour seed in accordance with the Pest Control Products Act and the Seeds Act Regulations. A blue colourant must be added when this product is applied to an oilseed.

How it Works:

Imidacloprid is a chloronicotinyl insecticide with systemic activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 455.

Labelling: All bags containing *Sombrero*-treated seed must be labeled or tagged as followed: "This seed has been treated with *Sombrero* 600I'S, which contains imidacloprid. Do not use for feed, food, or oil processing. Store away from feeds and other foodstuffs".

Grazing: Cover crops that are used as a rotational crop without a plant-back interval following treatment should not be grazed or harvested for food or feed. Do not graze or feed livestock on areas treated with *Sombrero* for four weeks after planting. Mustard greens grown or harvested

from *Sombrero*-treated seed must not be used for human consumption.

Re-cropping: Use a minimum plant-back interval of 30 days for cereals, 9 months for peas and beans, and 12 months for all other food and feed crops. Green manure and other cover crops not intended for human or animal consumption do not require a plant-back interval following treatment. DO NOT graze or harvest cover crops for food or feed. It is not recommended that this product be used in fields treated with imidacloprid during the previous season. DO NOT apply any subsequent application of imidacloprid in furrow or foliar application, or any other Group 4 insecticide following *Admire SPT/ Alias* 240 SC treatment.

Storage: Store product in cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of reach of children. Carry-over of *Sombrero*-treated seed is not recommended (treated canola, rapeseed or mustard (condiment-type only) seed stored for periods in excess of 6 months may decrease at a faster rate than untreated seed). Test seed germination if stored for more than 6 months. Do not store *Sombrero*-treated seed above 25C or in direct sunlight.

Environment: DO NOT plant treated seed pieces when rainfall is forecast for the next 48 hours. DO NOT plant treated seed pieces within 15 metres of well-head or aquatic systems, including marshes, ponds, ditches, streams, lakes, etc. This product is toxic to wildlife. Keep out of lakes, streams, ponds, or other aquatic systems. DO NOT contaminate water when disposing of equipment wash waters. Leftover treated seed should be double sown around the headland, or buried away from water sources such as lakes, streams, ponds or other aquatic systems.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Insure Cereal

Fungicide Group – 3, 4, 11

(Refer to page 312)

Company:

BASF Canada – PCP#30685

Formulation:

17 g per L pyraclostrobin, 17 g per L triticonazole, 10 g per L metalaxyl (both R and S isomers) formulated as a liquid suspension. Container size - 2 x 9.8 L jug

Crops, Rates and Diseases:

Crop:	Rate (ml) per 100 Kg seed	Diseases Controlled:	Diseases Suppressed:
Barley	300 mL	Seed rots and pre-emergence damping-off caused by <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and <i>Pythium</i> spp.; post-emergence damping-off caused by <i>Pythium</i> spp.; seedling blight and root rot caused by <i>Fusarium</i> spp. and <i>Pythium</i> spp.; true loose smut (<i>Ustilago nuda</i>); covered smut (<i>U. hordei</i>); false loose smut (<i>U. nigra</i>)	Seedling blight and root rot caused by <i>Cochliobolus sativus</i>
Oat	300 mL	Seed rots and pre-emergence damping-off caused by <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and <i>Pythium</i> spp.; post-emergence damping-off caused by <i>Pythium</i> spp.; seedling blight and root rot caused by <i>Fusarium</i> spp. and <i>Pythium</i> spp.; loose smut (<i>Ustilago avenae</i>); covered smut (<i>U. kolleri</i>)	Seedling blight and root rot caused by <i>Cochliobolus sativus</i>
Wheat, rye, triticale	300 mL	Seed rots and pre-emergence damping-off caused by <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and <i>Pythium</i> spp.; post-emergence damping-off caused by <i>Pythium</i> spp.; seedling blight and root rot caused by <i>Fusarium</i> spp. and <i>Pythium</i> spp.; loose smut (<i>Ustilago tritici</i>); common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>)	Seedling blight and root rot caused by <i>Cochliobolus sativus</i>

Application Information:

Insure Cereal is a ready to use seed treatment formulation. This product is for used in commercial seed plant, in on-farm standard gravity flow or mist type treatment machines, and in on-the-go air seeder treatment systems. Agitate or shake well prior to usage. Uneven seed coverage may result in poor levels of disease control. Seed should be well conditioned and cleaned prior to treating. Treated seed should not require drying after treatment.

How it Works:

The active ingredient pyraclostrobin is a strobilurin fungicide with broad spectrum contact and systemic activity. The active ingredient triticonazole is a triazole fungicide that provides systemic broad spectrum activity. The active ingredient

metalaxyl is an acylalanine fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labeled "This seed has been treated with *Insure Cereal* containing fungicides pyraclostrobin, triticonazole and metalaxyl. Workers handling or planting treated seed must wear long-sleeved shirt, long

pants, chemical-resistant gloves, shoes and socks, and respiratory protection (i.e. NIOSH/MSHA/BHSE approved respirator or fresh air hood). Respiratory protection is not required when workers are in a closed cab tractor. A closed cab is a chemical resistant barrier that completely surrounds the occupant of the cab and prevents contact with the pesticide or treated surfaces outside the cab. DO NOT use for food, feed or oil processing. Store away from feed or food stuff. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store in original containers with lid tightly closed. Store away from children, animals, feed stuffs, fertilizers and seed. Protect from frost and freezing. DO NOT store treated seed for more than 18 months. Store treated seed in cool, dry conditions.

Environment: Ensure proper soil incorporation of the seeds. DO NOT feed treated seed to, or otherwise expose, wildlife or domestic birds. DO NOT contaminate domestic or irrigation water supplies, lakes, streams, ponds or any body of water with the chemical, used containers, treated seed or bags. DO NOT contaminate water by cleaning of equipment or disposal of wastes.

Hazard Rating:



Caution – Poison



Caution – Eye irritant and potential skin sensitizer

For an explanation of the symbols used here see page 11.

Kocide 2000

Fungicide Group – M1

(Refer to page 312)

Company:

E.I. du Pont Canada Company – (PCP# 27348)

Formulation:

35% copper as copper hydroxide formulated as a dry flowable.

Container size - 10 to 25 kg.

Crops, Diseases and Timing:

User requested minor use label expansion (URMULE) for use as a seed treatment on beans. For suppression of the following seed borne diseases; common blight (*Xanthomonas axonopodis* pv. *phaseoli*), halo blight (*Pseudomonas syringae* pv. *phaseolicola*) and bacterial brown spot (*Pseudomonas syringae* pv. *syringae*), in dry edible beans and succulent beans.

Rate:

For every 100 kg seed to be treated, add 113 g Kocide 2000 to 200 ml of warm water and stir until dissolved. Ensure even coverage and allow treated seed to dry before planting.

Application Information:

Kocide 2000 used as a seed treatment may cause some delay in seed germination. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment.

All seed treated with this product must be conspicuously coloured at the time of treatment. When handling contaminated equipment or treated seed, wear long sleeved coveralls over normal work clothing, chemical resistant gloves and goggles; use an approved respirator with a canister approved for pesticides to minimize exposure to dust when handling, mixing leading product, bagging seed,

sewing bags of treated seed or when transferring treated seed to a storage bin.

Refer to the label for additional application instructions and/or use precautions regarding this URMULE.

How it Works:

The active ingredient copper hydroxide is an inorganic fungicides with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labeled as follows: "This seed has been treated with Kocide 2000; DO NOT use treated seed for food, feed or oil processing."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store in cool, dry, ventilated area, away from feed or food. Keep away from heat, fire and sparks.

Environment: DO NOT apply or allow to drift onto streams or any body of water.

Hazard Rating:



Caution – Poison



Warning – Eye irritant

For an explanation of the symbols used here see page 11.

Lumiderm

Insecticide Group – 28

(Refer to page 456)

Company:

E.I. du Pont Canada Company – PCP#30894

Formulation:

625 g per L cyantraniliprole formulated as a suspension.

Container size – 100 L, 1000 L, Bulk

Crops, Rates and Insects:

Crop:	Rate per 100 Kg seed:	Insects Controlled:
Canola, Rapeseed, Oilseed Mustard	960 to 1600 mL	Flea beetles
	480 to 960 mL	Cutworms

* The application rates for flea beetles will also provide early season protection from cutworm feeding damage. Use higher rates in areas with high pest pressure, or where extended early season control is required.

Application Information:

For use in closed treatment systems at commercial seed treatment facilities only. The product contains no colourant. An appropriate colourant must be added when the product is applied. Seed must be conspicuously coloured at the time of treatment.

How it Works:

The active ingredient is a cyantraniliprole is a systemic insecticide from the diamides chemical class. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

Seed Treatments: *Prosper EverGol Seed Treatment* or *Helix Xtra Seed Treatment*.

Restrictions:

Resistance management: Refer to page 455.

Labelling: Treated seed must be labeled "This seed has been treated with *Lumiderm Insecticide Seed Treatment* which contain cyantraniliprole. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs. Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed. This product is toxic to aquatic organisms. Dispose of all excess treated seed. Left over treated seed may be double-sown around the

headland or buried away from water sources in accordance with local requirements. Dispose of seed packaging in accordance with local requirements. Cover or incorporate spilled treated seeds."

Grazing: No restrictions listed.

Re-cropping: Registered crops for *Lumiderm* may be replanted at any time. For cereal, pulse and forage crops DO NOT plant back within 30 days of seeding with *Lumiderm* treated seed.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed. To prevent contamination store this product away from food or feed.

Environment: Toxic to aquatic organisms and bees. When this product is applied and used according to label directions, risk to bees is expected to be negligible. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:



Warning – Poison



Caution – Eye irritant and potential skin sensitizer

For an explanation of the symbols used here see page 11.

Mancozeb

Potato ST 16 / Solan MZ / Tuberseal

Fungicide Group – M3
(Refer to page 312)

Company:

Norac Concepts Inc. (Solan MZ – PCP#29377, Tuberseal – PCP#17042)

Wilbur-Ellis Co., distributed by United Agri Products (Potato ST 16 – PCP#24734)

Formulation:

16% mancozeb formulated as a powder. Container size - 10 kg (Tuberseal); 20 kg (Solan MZ and Potato ST 16).

Crops and Disease:

Control of *Fusarium* seed piece decay in potatoes.

Rates:

500 g per 100 kg seed.

Application Information:

Apply product before planting; thoroughly coat surface of whole or cut seed with dust. If treated whole seed is cut, make a second application to protect cut surfaces. Plant as soon as possible after treating. If cut seed is not planted within two days of treating, store in a ventilated location to allow cut surfaces to dry.

How it Works:

The active ingredient mancozeb is a dithiocarbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed pieces should be labelled "Poisonous to man and animals. This seed has been treated with mancozeb for the control of fusarium decay. Do not use for food or feed purposes."

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store product in a cool, dry, well-ventilated place. Keep away from fire and sparks.

Environment: DO NOT contaminate feed or food. DO NOT contaminate any body of water.

Hazard Rating:

None listed.

Maxim D/Maxim Liquid PSP/ Maxim PSP/Maxim MZ PSP

Fungicide Group –
Maxim D – 12, 3
Maxim Liquid PSP,
Maxim PSP – 12
Maxim MZ PSP – 12, M4
(Refer to page 312)

Company:

Syngenta Canada (*Maxim D* – PCP #30599; *Maxim Liquid PSP* – PCP #29110; *Maxim PSP* – PCP#26647; *Maxim MZ PSP* – PCP#27965)

Formulation:

Maxim D: 19.4 g per L fludioxonil and 19.4 g per L difenoconazole formulated as a suspension.

Maxim Liquid PSP: 40.3% fludioxonil net contents 1 L to 55 L.

Maxim PSP: 0.5% fludioxonil formulated as a dry powder. Container size 10 kg.

Maxim MZ PSP: 0.5% fludioxonil plus 5.7% mancozeb formulated as a dry powder. Container size – 10 kg, 20 kg, 22.7 kg.

Crops and Diseases Controlled:

Seed piece treatment fungicide for use on potato for the control of black scurf/stem and stolon canker (*Rhizoctonia solani*), silver scurf (*Helminthosporium solani*), and fusarium dry rot (*Fusarium* spp.). These products do not control bacterial diseases present within the seed.

Rate:

Maxim D: 130 mL per 100 kg of seed for control of black scurf/stem and stolon canker and silver scurf; 65-130 mL per 100 kg of seed for control of fusarium dry rot (use the high rate when disease pressure and/or resistance pressure is high); 65 mL per 100 kg of seed for suppression of black scurf and control of stem and stolon canker (*Rhizoctonia solani*).

Maxim Liquid PSP: 5.2 mL per 100 kg of seed.

Maxim PSP; *Maxim MZ PSP*: Use 500 g per 100 kg of cut seed. One 10 kg bag treats 2000 kg of seed.

Application Information:

Maxim D: Shake or mix well before using. Apply using standard seed treatment equipment that provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. When applying at a rate of 130 mL/100 kg seed, *Maxim D* may be applied undiluted or

as a water-based slurry. DO NOT exceed a maximum slurry volume of 260 mL per 100 kg seed. When applying at rates lower than 130 mL of *Maxim D* per 100 kg seed, add sufficient water to allow for a slurry volume of at least 130 mL per 100 kg seed. Follow manufacturer's application instructions for the seed treatment equipment used.

Maxim Liquid PSP: Shake or mix well before using. Apply as a water-based slurry utilizing standard slurry seed treatment equipment which provides uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of disease control. Thoroughly mix the recommended amount of seed treatment into the required amount of water for the slurry treater and dilution rate to be used. Follow the manufacturer's application instructions for the seed treatment equipment being used. Maintain constant agitation of the slurry during the treatment. Consult the label for more information on seed piece treatment application procedures.

Maxim PSP; *Maxim MZ PSP*: Apply using appropriate treater designed for treating potatoes or by dust attachment over belt. Cut pieces should be treated immediately after cutting. If treated seed pieces are bagged, they should be stored for 2 to 3 days in open crates before bagging. For optimum protection against silver scurf, ensure that seed tubers are completely free of soil. Total skin coverage is essential.

How it Works:

Fludioxonil is a phenylpyrrole fungicide with contact activity. Mancozeb is a dithiocarbamate fungicide with contact activity. The active ingredient difenoconazole is a triazole fungicide with broad-spectrum systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

For control of potato diseases and insects, *Maxim Liquid PSP* fungicide seed protectant can be tank-mixed with *Actara 240* insecticide. Consult each label for pests controlled, precautions, and specific application instructions. For control of Colorado potato beetle, aphids and potato leafhopper, *Maxim D* can be tank-mixed with *Cruiser Potato Seed Piece Treatment*. Refer to product labels for specific application instructions and precautions. Follow the most restrictive label precautions and limitations.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Ideal storage temperature for the product is above freezing and below 30°C. Store in a dry place. Avoid contamination of feed.

Environment: This product is toxic to fish and aquatic invertebrates. DO NOT apply directly to water, or to areas where surface water is present. In cleaning of equipment or disposing of wastes, DO NOT contaminate water used for human or animal consumption or by wildlife and aquatic life or for irrigation purposes.

Hazard Rating:



Caution – Poison

Other precautions: Keep out of reach of children. May be harmful if swallowed. Avoid inhalation of dust, vapour and/or spray mist. Wash hands and face after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Treated seed must not be used for food, feed or oil processing. Treat seed in a well-ventilated area. When handling, contaminated equipment or treated seed, wear long-sleeved coveralls over long pants and long-sleeved shirt and chemical-resistant gloves plus goggles.

For an explanation of the symbols used here see page 11.

Maxim Quattro

Available to commercial seed treaters only.

Company:

Syngenta Canada – PCP#29871

Formulation:

26.5% thiabendazole, 3.32% fludioxonil, 2.65% metalaxyl-M and S-isomer, 1.33% azoxystrobin formulated as a liquid suspension seed treatment.

Container size - 5 L to Bulk.

Crops and Diseases:

Control of seed- and soil-borne *Pythium*, *Rhizoctonia*, and *Fusarium* (including *F. graminearum* and *F. verticillioides*) and weakly pathogenic *Aspergillus* and *Penicillium* causing seed rot/pre-emergence damping-off, post-emergence damping-off, and seedling blight on corn.

Rate:

67 mL per 100 kg seed.

Application Information:

For use by a commercial seed treater only. Mix with water to form a slurry seed treatment. Contains no colourant; an appropriate colourant must be added to slurry before treating seed. Maintain constant agitation of slurry. Allow seed to dry before bagging. Treatment of highly mechanically dam-

Fungicide Group – 1, 4, 11, 12

(Refer to page 312)

aged, poor quality or low vigour seed may result in reduced germination and/or reduced seed and seedling vigour. If seed lot quality is unknown conduct a germination test prior to treating.

How it Works:

The active ingredient thiabendazole is a benzimidazole fungicide with contact and systemic activity. The active ingredient fludioxonil is a phenylpyrrole fungicide with contact and locally systemic activity. The active ingredient metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes class, including *Pythium* damping off. The active ingredient azoxystrobin is a methoxyacrylate (strobilurin) fungicide with broad spectrum activity to be used as a preventative and curative fungicide. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled "This seed has been treated with thiabendazole, fludioxonil, metalaxyl-M and S-isomer, and azoxystrobin. Use chemical resistant-

gloves when handling treated seed. DO NOT use for food, feed or oil processing. Store away from feed and foodstuffs. DO NOT graze corn or cut for forage within 30 days of planting."

Grazing: DO NOT graze or feed livestock on treated areas within 30 days of planting.

Re-cropping: No restrictions listed.

Storage: Store product between 0°C and 30°C. Repeated freeze-thawing of the product will not compromise its integrity. If the product should freeze, bring the product back to room temperature and ensure thorough mixing before use. Store away from food and feed.

Environment: DO NOT apply directly to water, or to areas where surface water is present. In cleaning of equipment or

disposing of wastes, DO NOT contaminate water used for human or animal consumption or by wildlife and aquatic life or for irrigation purposes. If treated seed is spilled outdoors, promptly clean up.

Hazard Rating:

 Caution – Potential skin sensitizer

For an explanation of the symbols used here see page 11.

Mertect SC

Fungicide Group – 1
(Refer to page 312)

Company:

Syngenta Canada – PCP#13975

Formulation:

500 g per L thiabendazole formulated as a water dispersible suspension. Container size - 4 x 5 L.

Crops and Diseases:

Post-harvest control of storage rots caused by *Fusarium*, *Phoma*, *Helminthosporium*, *Oospora* and *Rhizoctonia* spp. on potato.

Rate and Water Volume:

7.5 L per 170 L of water. Spray 2 L of this suspension per 1 metric tonne of potatoes.

Application Information:

Post-harvest treatment. Shake well before using. DO NOT allow suspension to stand without continuous agitation. Potatoes must rotate along conveyor line to ensure complete coverage. Prior to treating potatoes destined for export, confirm with authorities that treated potatoes will be allowed to enter importing country.

How it Works:

The active ingredient thiabendazole is a benzimidazole fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Minimum storage temperature 0°C.

Environment: Toxic to aquatic organisms. Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. Do not discharge effluent containing this product into sewer systems, lake, streams, ponds, estuaries, oceans, and other waters.

Hazard Rating:

None listed.

Metalaxyl

Allegiance FL/Belmont 2.7FS

Fungicide Group – 4
(Refer to page 312)

Company:

Bayer CropScience (*Allegiance FL* – PCP#26674)
Chemtura Canada Co. (*Belmont 2.7 FS* – PCP#30246)

Formulation:

317 g per L metalaxyl formulated as a liquid seed treatment.
Allegiance FL container size – 4 X 3.79 L.
Belmont 2.7 FS container size – 500 mL, 10 L and 200 L.

Crops, Diseases and Rates (for crops processed in Canada):

Crop	Disease Controlled:	Application Rates* (mL per 100 kg seed treated):	Water Volume (mL required to make up a total volume of 500 mL):
Chickpea, dry pea	Seed rots and seedling blights caused by <i>Pythium</i> spp.	16 to 110	484 to 390
Canola (rapeseed)	Seed rots and seedling blights caused by <i>Pythium</i> spp.	32 to 110	468 to 390
Alfalfa, bean, clover, corn, sainfoin, vetch	Seed rots and seedling blights caused by <i>Pythium</i> spp.	46 to 110	454 to 390
Grasses (forage)	Seed rots and seedling blights caused by <i>Pythium</i> spp.	46 to 93	454 to 407
Grasses (turf)	Seed rots and seedling blights caused by <i>Pythium</i> spp.	93	407
Soybean	Seed rots and seedling blights caused by <i>Pythium</i> spp., early season <i>Phytophthora</i>	46 to 93	454 to 407
Sunflower	Seed rots and seedling blights caused by <i>Pythium</i> spp., downy mildew	110 to 189**	390 to 311
Low tannin lentil***	Seed rots and seedling blights caused by <i>Pythium</i> spp.	16	484

* Use the high rate if planting into cold, wet soils, if the seed is of poor quality, or if disease pressure is expected to be high.

** High rate is for downy mildew control.

*** For use on low tannin lentils destined export or seed production only.

Crops, Diseases and Rates (for crops processed in Canada) *continued*:

Crop	Diseases Controlled:	Application Rates** (mL per 100 kg seed treated):	Water Volume(mL required to make up a total volume of 500 to 620 mL):
Corn	Downy mildew (<i>Sclerophthora macrospora</i>)	189 to 620	311 to 0
Pea	Downy mildew (<i>Peronospora viciae</i>)	146	354
Sunflower	Downy mildew (<i>Plasmopora halstedii</i>)	620	0
Cereals (wheat, barley, oats, rye)	Seed rots and seedling blights caused by <i>Pythium</i> spp.	46 to 110	454 to 390
Sorghum	Seed rots and seedling blights caused by <i>Pythium</i> spp.	93 to 110	407 to 390
Sorghum	Downy mildew (<i>Perenosclerospora sorghi</i>)	189	311
Bird's-foot trefoil	Seed rots and seedling blights caused by <i>Pythium</i> spp.	46 to 110	454 to 390
Low-tannin lentil	Seed rots and seedling blights caused by <i>Pythium</i> spp.	16	484

* Use the high rate if planting into cold, wet soils, if the seed is of poor quality, or if disease pressure is expected to be high.

Application Information:

Mix with water to form a slurry seed treatment. Contains no colourant; an appropriate colourant must be added to slurry before treating seed. Maintain constant agitation of slurry. Allow seed to dry before bagging. Treatment of highly mechanically damaged, poor quality or low vigour seed may result in reduced germination and/or reduced seed and seedling vigour. If seed lot quality is unknown conduct a germination test prior to treating.

How it Works:

The active ingredient metalaxyl is an acylalanine fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled as follows; "This seed has been treated with *Allegiance FL* or *Belmont 2.7 FS* seed protectant which contains metalaxyl. DO NOT use for feed, food or oil processing." All bags containing seed for export must be labelled "FOR EXPORT ONLY."

Grazing: DO NOT graze or feed livestock on treated areas for 4 weeks after planting.

Re-cropping: No restrictions listed.

Storage: DO NOT store above 35°C or below 0°C. Store in original container, away from pesticides, food or feed.

Environment: Treated seed may be toxic to birds and other wildlife. Clean up any spilled seeds and ensure seed is properly incorporated at planting.

Hazard Rating:



Warning – Skin and eye irritant

For an explanation of the symbols used here see page 11.

Polyram 16D

Fungicide Group – M3
(Refer to page 312)

Company:

BASF Canada – PCP# 25867

Formulation:

16% metiram formulated as dust. Container size - 20 kg.

Crops and Diseases:

Control of *Fusarium* seed piece decay; seed-borne common scab (*Streptomyces* spp.) on potato.

Rates:

450 to 650 g per 100 kg seed. One container treats 2,222 to 1,538 kg of seed.

Application Information:

Apply to entire surface of seed pieces after cutting. Treated potato seed should not be allowed to stand in hot sun or drying wind. Seed should not be planted into hot, dry soil or into cold, wet soil. Plant as soon as possible after treating. If not planted immediately, provide sufficient ventilation to allow cut surface to dry. If not cutting seed, treat at same rate for control of seed-borne common scab, apply a second application to control fusarium seed piece decay.

How it Works:

The active ingredient metiram is a dithiocarbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store product in cool, dry, locked, well-ventilated area without floor drain. Keep away from heat, fire or sparks. DO NOT allow product to become wet or overheated as this will reduce its effectiveness and may create flammable vapours. Protect from freezing. DO NOT ship or store near food, feed, seed or fertilizers.

Environment: DO NOT contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

Hazard Ratings:

None listed.

Proseed

Fungicide Group – 12
(Refer to page 312)

Proseed is available to commercial seed treaters only

Company:

Syngenta Canada – PCP#29814

Formulation:

40.3% fludioxonil formulated as a seed treatment solution.
Container size - 1 L to bulk.

Crops and Diseases:

Protection of wheat, barley, oat, rye, corn, triticale, sorghum, and buckwheat seed and seedlings from seed- and soil-borne fungi which cause seed decay, damping-off, and seedling blights including *Fusarium* spp. and *Rhizoctonia* spp.

Rate:

5.2 to 10.4 ml per 100 kg of seed.

Note: use higher rates for crop/cultivars that are more susceptible to pathogens or when higher disease pressure is expected.

Application Information:

For use in commercial seed treatments facilities only. Apply as a water-based slurry using standardized equipment to ensure uniform seed coverage. Maintain constant agitation of the slurry during treatment. Allow seed to dry before bagging. This product contains no colourant. A red colourant must be added when this product is applied to grain.

How it Works:

Fludioxonil is a phenylpyrrole fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Proseed may be mixed with *Apron XL* seed treatment for control of diseases caused by *Pythium* spp. and *Phytophthora* spp. provided *Apron XL* is registered on that crop.

Follow the label directions for each product and use the most restrictive precautions and limitations for either product.

Restrictions:

Resistance management: Refer to page 310.

Labelling: All bags containing treated seed for sale or use in Canada must be labelled or tagged as follows: "This seed has been treated with fludioxonil fungicide (or fludioxonil and metalaxyl-M fungicide if appropriate). DO NOT use for feed, food or oil purposes. Store away from feeds and foodstuffs. DO NOT graze treated crops, or cut for forage, within 30 days of planting."

Grazing: DO NOT graze or feed livestock on treated areas for 30 days after planting.

Re-cropping: No restrictions listed.

Storage: Store in temperatures above freezing and below 30°C.

Environment: This product is toxic to fish and other aquatic organisms. DO NOT apply this product directly to water, or to areas where surface water is present. In cleaning of equipment or disposing of wastes, do not contaminate water used for human or animal consumption, or by wildlife and aquatic life, or for irrigation purposes.

Hazard Ratings:

None listed.

Prosper EverGol/ Prosper FX/ Poncho 600 FS/Titan ST

Fungicide Group –
Prosper EverGol/
Prosper FX - 4,7,11
Insecticide Group – 4
(Refer to page 312 and 456)

Prosper EverGol, Prosper FX and Poncho 600 FS available to commercial seed treaters only. Titan ST is available for on-farm seed treatment. Poncho 600 FS and Titan ST contain insecticide only.

Company:

Bayer CropScience (*Prosper EverGol* – PCP#30363; *Prosper FX* – PCP#29159; *Poncho 600 FS* – PCP#27453; and *Titan ST 600 FS* – PCP#27449)

Formulation:

Active ingredient:	Active ingredient strength (g per L):			
	<i>Prosper EverGol</i>	<i>Prosper FX</i>	<i>Poncho 600 FS</i>	<i>Titan ST</i>
Container sizes:	3.8L-1000L, bulk	56.8L, 100L, 1000L, bulk	1L, 3.8L, 100L, 113L, 200L, 1000L	1L, 3.8L, 10L, 200L, 1000L
Clothianidin	290	285.7	600	600
Carbathiin	-	50	-	-
Penflufen	10.7	-	-	-
Trifloxystrobin	7.15	7.14	-	-
Metalaxyl	7.15	5.36	-	-

Crops, Insects, Diseases and Rates:

Product:	Crop:	Rate per 100 kg of seed:	Insects Controlled:	Diseases Controlled:
<i>Prosper EverGol</i>	Canola, Rapeseed, Mustard (oilseed and condiment)	1400 mL	Flea beetles	Seed rot, damping off, seedling blight and early season root rot caused by <i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , and seed-borne <i>Alternaria</i> spp.; seed-borne blackleg (<i>Leptosphaeria maculans</i>).
<i>Prosper FX</i>	Canola, Rapeseed	1400 mL		
<i>Poncho 600 FS</i>	Canola, Rapeseed	250, 333 or 666 mL ¹	Flea beetles	-
	Corn	33.3 to 66.6 mL of product per 80,000 units of seed (0.25 to 0.5 mg a.i. per kernel)	Wireworm; seed corn maggot; black cutworms; corn flea beetle; white grubs	-

See footnotes on following page.

Continued...

Crops, Insects, Diseases and Rates *continued*:

Product:	Crop:	Rate per 100 kg of seed:	Insects Controlled:	Diseases Controlled:
<i>Titan ST</i>	Potatoes	20.8 mL per 100 kg potato seed pieces	Wireworms	—
		10.4 to 20.8 mL per 100 kg potato seed pieces	Aphid (potato, green peach, foxglove and buckthorn aphids), Colorado potato beetle, potato leafhopper, potato flea beetle (overwintered adults and suppression of second generation)	—

¹ Increasing rates for low, moderate and severe flea beetle pressure.

Application Information:

Prosper Evergol, *Prosper FX* and *Poncho 600 FS* are for use in commercial seed treatment facilities with closed transfer systems only. *Prosper FX* and *Poncho 600 FS* DO NOT contain a colourant. An appropriate colour must be added when these products are applied. Seed treatment must be thoroughly agitated to ensure uniform mixing of product prior to and during application. Treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour.

Titan ST is a seed piece treatment. Apply specified rate as a diluted spray onto seed pieces using a well contained, shielded spray system to prevent the loss of any liquid. Apply only in areas with adequate ventilation or in areas equipped to remove spray mist or dust. Agitate or stir spray solution as needed. For optimal insect control good coverage of seed pieces is required. DO NOT dilute with any more than 6 parts water to 1 part *Titan ST*. Plant seed pieces as soon as possible after cutting and treating.

How it Works:

Clothianidin is a chloronicotinyl insecticide with systemic activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456. Carbothiin is a carboxamide fungicide with systemic activity; penflufen is a carboxamide (SDHI) fungicide with systemic activity; tri-floxystrobin is a strobilurin fungicide with broad spectrum preventative activity; and metalaxyl is an acylalanine fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Subsequent applications: DO NOT make any subsequent application of a group 4 insecticide (in furrow or foliar application) following treatment with any of these products.

Labelling: Treated seed must be labelled as follows: "This seed has been treated with clothianidin and/or carbothiin, penflufen and metalaxyl. DO NOT use for food, feed or oil processing. Store away from feeds and other foodstuffs."

Grazing: *Prosper FX* – Do not graze or feed livestock on seeded areas for four weeks after planting.

Re-cropping: For *Prosper Evergol*, *Prosper FX* and *Poncho 600 FS*, corn and canola may be replanted at any time. For *Titan ST*, corn, and canola and potatoes may be replanted at any time. For all products, a 1-year plant back interval is required for leafy, root and tuber vegetables. A 30-day plant back is required for cereals, grasses, nongrass animal feeds, soybeans and dry beans.

Storage: Protect products from freezing. DO NOT contaminate water, food or feed by storage, disposal or by cleaning of equipment. Store in a cool place. DO NOT store in direct sunlight. Store away from food or feed. DO NOT store treated seed above 25°C or in direct sunlight. Treated seed stored for periods in excess of 9 months should be tested for germination before planting.

Environment: These products are toxic to aquatic invertebrates. DO NOT apply directly to water or to areas where surface water is present. DO NOT contaminate water when disposing of equipment wash waters. These products are toxic to birds and mammals. Any spilled or exposed seeds should be incorporated into the soil or otherwise cleaned up from the soil surface.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Rancona Apex

Fungicide Group - 3

(Refer to page 312)

Company:

Manufactured by Chemtura Canada Co - PCP#29176

Formulation:

4.61 g per L ipconazole formulated as a micro-dispersion liquid. Container sizes - 10 L, 200 L, 1000 L.

Crops, Diseases and Rates:

Crop:	Rate (mL) per 100 kg of seed:	Diseases Controlled:	Diseases Suppressed*:
Barley	325	General seed rots (including those caused by saprophytic organisms such as <i>Penicillium</i> and <i>Aspergillus</i>); seed rot, damping off and seedling blight (seed- and soil-borne <i>Fusarium</i>); seed rot and seedling blight (seed and soil-borne <i>Cochliobolus sativus</i>); true loose smut (<i>Ustilago nuda</i>), covered smut (<i>U. hordei</i>), false loose smut (<i>U. nigra</i>), leaf stripe (<i>Pyrenophora graminea</i>)	Common root rot (<i>Cochliobolus sativus</i>) Crown and foot rot (<i>Fusarium</i>)
	325 to 433	True loose smut (<i>Ustilago nuda</i>) control. Use the higher rate for highly infected seed lots only.	
Wheat	325	Loose smut (<i>Ustilago tritici</i>); common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>); seed rot and seedling blight caused by <i>C. sativus</i> , <i>Fusarium</i> , <i>Aspergillus</i> , and <i>Penicillium</i> .	
Oat	325	Loose smut (<i>Ustilago avenae</i>); covered smut (<i>U. kolleri</i>); seed rot and seedling blight caused by <i>Fusarium</i> , <i>C. sativus</i> , <i>Aspergillus</i> and <i>Penicillium</i> .	
Rye	325	Seed rot and seedling blight caused by <i>Fusarium</i> , <i>C. sativus</i> , <i>Aspergillus</i> and <i>Penicillium</i> .	
Triticale	325	Seed rot and seedling blight caused by <i>Fusarium</i> , <i>C. sativus</i> , <i>Aspergillus</i> and <i>Penicillium</i> .	

* Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

Application Information:

Rancona Apex is designed to be used undiluted in commercial seed treaters and by dripper applicators. Undiluted *Rancona Apex* can be used at temperatures down to -20°C.

How it Works:

The active ingredient ipconazole is a demethylation inhibitor with systemic and contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

RANCONA Apex Fungicide may be tank mixed with *Apron XL LS* and applied to wheat (spring or winter), barley, oats or rye. Apply *RANCONA Apex Fungicide* at any rate recommended in the table above, and apply *Apron XL LS* at 2.7 ml/100Kg.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled as follows "This seed has been treated with *Rancona Apex* liquid seed protectant containing ipconazole. Do not use for feed, food, or oil processing."

Grazing: DO NOT graze or feed livestock on treated area for 30 days after planting.

Re-cropping: No restrictions listed.

Storage: DO NOT store in direct sunlight or above 35°C. DO NOT freeze.

Environment: DO NOT contaminate ponds, lakes or streams.

Hazard Rating:

None listed.

Rancona RS

Fungicide Group – 3, 7
(Refer to page 312)

Company:

Manufactured by Chemtura Canada Co – PCP#30217

Formulation:

9.38 g per L ipconazole and 87.5 g per L carbathiin formulated as a liquid suspension seed treatment. Container sizes - 10 L, 1000 L.

Crops, Diseases and Rates:

Crop:	Rate (mL) per 100 kg of seed:	Diseases Controlled:	Diseases Suppressed*:
Canola, rape-seed	800	Seed rot, damping off, and seedling blight caused by <i>Rhizoctonia</i> , and <i>Fusarium</i> spp. Seed-borne blackleg (<i>Leptosphaeria maculans</i>)	Root rot caused by <i>Rhizoctonia</i> and <i>Fusarium</i> spp.

Application Information:

Rancona RS is ready to use and may be applied to seed as purchased. However, dilution with water may help to achieve more uniform seed coverage when using some types of treaters and/or when treating under dry and/or hot conditions.

How it Works:

The active ingredient ipconazole is a demethylation inhibitor with systemic and contact activity and carbathiin is a carboximide fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled as follows "This seed has been treated with carbathiin and ipconazole. Do not use treated seed for feed, food, or oil processing."

Grazing: DO NOT graze or feed livestock on treated area for four weeks after planting.

Re-cropping: No restrictions listed.

Storage: Store this product away from food or feed.

Environment: DO NOT contaminate ponds, lakes or streams.

Hazard Rating:

None listed.

Raxil MD

Fungicide Group – 3, 4

(Refer to page 312)

Company:

Bayer CropScience – PCP#27692

Formulation:

5.0 g per L tebuconazole and 6.6 g per L metalaxyl formulated as a flowable seed treatment. Container size - 10 L, 200 L, 1,000 L.

Crops, Rates and Diseases:

Crop:	Undiluted Use Rate:		Diseases Controlled:	Diseases Suppressed*:
	Rate per 100 kg of Seed	Amount of Seed per 10 L Jug		
Wheat	300 mL	3320 kg	Loose smut (<i>Ustilago tritici</i>), common bunt or stinking smut (<i>Tilletia tritici</i> , <i>T. laevis</i>); seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Fusarium</i> spp.; damping-off caused by <i>Pythium</i> spp.; seed-borne <i>Septoria nodorum</i> .	Root and crown rot caused by seed- and soil-borne <i>Fusarium</i> spp.; common root rot caused by seed- and soil-borne <i>Cochliobolus sativus</i> ; seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>C. sativus</i> ; seedling blight caused by seed-borne <i>C. sativus</i> .
Barley	300 mL	3320 kg	True loose smut (<i>Ustilago nuda</i>), covered smut (<i>U. hordei</i>), false loose smut (<i>U. nigra</i>); seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Fusarium</i> spp.; damping-off caused by <i>Pythium</i> spp.; barley leaf stripe (<i>Pyrenophora graminis</i>).	
Oat	300 mL	3320 kg	Covered smut (<i>Ustilago kolleri</i>), loose smut (<i>U. avenae</i>); seed rot and pre-emergent damping-off caused by seed- and soil-borne <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Fusarium</i> spp.; damping-off caused by <i>Pythium</i> spp.	

* Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

Application Information:

Raxil MD is a ready to use formulation designed for commercial or on-farm treatment with conventional seed treating equipment which can accurately control application rates and provide good distribution of the chemical onto the seed in the mixing chamber. Uniform application to seed is necessary to ensure seed safety and best disease control. See manufacturer's instructions supplied with the treater system for information on proper application technique.

Uniform coverage at the correct rate is important for satisfactory results. Under-treatment may lead to loss of efficacy and over-treatment could reduce germination. Seed may be planted immediately after treating.

How it Works:

The active ingredient tebuconazole is a systemic triazole fungicide with broad-spectrum activity. The active ingredient metalaxyl is an acylalane fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled "This seed has been treated with *Raxil MD* Fungicide which contains tebuconazole and metalaxyl; DO NOT use for food, feed or oil processing."

Grazing: DO NOT graze or feed livestock on treated areas for 4 weeks after planting.

Re-cropping: No restrictions listed.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Store in a cool, dry place and avoid excessive heat.

Environment: DO NOT contaminate water, food, or feed by storage, disposal, or by cleaning of equipment.

Hazard Rating:



Danger – Skin and eye irritant

For an explanation of the symbols used here see page 11.

Raxil PRO

Fungicide Group – 3, 4
(Refer to page 312)

Company:

Bayer CropScience – PCP#30102

Formulation:

3.0 g per L tebuconazole, 15.4 g per L prothioconazole and 6.2 g per L metalaxyl formulated as a suspension. Container sizes - 10 L, 58.5 L, 175.5 L, 1000 L.

Crops, Diseases and Rates:

Crop:	Rate per 100 kg of seed:	Diseases Controlled:	Diseases Suppressed*:
Wheat, barley, oat	325 mL	Seed rot/pre-emergence damping-off caused by seed-borne <i>Fusarium</i> spp., <i>Aspergillus</i> spp. and <i>Cochliobolus sativus</i> ; seed rot/pre-emergence damping-off caused by soil-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and <i>Pythium</i> spp.; post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp. and seed-borne <i>Aspergillus</i> spp.; seedling blight caused by seed-borne <i>Aspergillus</i> spp.; true loose smut (<i>Ustilago nuda</i>); covered smut (<i>U. hordei</i> , <i>U. kollerii</i>); false loose smut (<i>U. nigra</i>); loose smut (<i>U. tritici</i> , <i>U. avenae</i>); common bunt (<i>Tilletia caries</i> , <i>T. laevis</i>); barley leaf stripe (<i>Pyrenophora graminea</i>)	Root rot caused by soil-borne <i>Fusarium</i> spp., <i>Cochliobolus sativus</i> , and seed-borne <i>Fusarium</i> spp.; crown rot caused by seed-borne <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Penicillium</i> spp.

Application Information:

Raxil PRO is a ready-to-use treatment formulation for use in commercial seed treatment operations and for on-farm treatment with conventional seed treating which can accurately meter, mix and apply flowable seed treatment formulations.

How it Works:

The active ingredient tebuconazole is a triazole demethylation inhibitor (DMI) fungicide with systemic activity. The active ingredient prothioconazole is a demethylation inhibitor with broad-spectrum systemic activity. The active ingredient metalaxyl is an acylalanine fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Insecticide Seed Treatment: May be mixed with *Stress Shield 600*.

Follow the label directions for each product and use the most restrictive precautions and limitations for either product.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labeled "This seed has been treated with *Raxil PRO*, which contains tebuconazole, prothioconazole and metalaxyl. When handling treated seed wear chemical-resistant gloves. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs."


Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed. Store in a cool, dry area and avoid excessive heat.

Environment: Toxic to aquatic organisms. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes. Dispose of all excess treated seed. Left over seed may be double-sown around the headland or buried away from water sources. DO NOT leave exposed treated seed on soil surface. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface. The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable and/or the depth to the water is shallow.

Hazard Rating:

 Caution – Skin irritant

For an explanation of the symbols used here see page 11.

Raxil PRO Shield

Fungicide Group – 3, 4
Insecticide Group – 4

(Refer to page 312 and 456)

Raxil PRO Shield is a co-pack of *Raxil PRO* (tebuconazole, prothioconazole and metalaxyl fungicides, page 433) and *Stress Shield 600* (imidacloprid insecticide, page 439). For other detailed information on the component products see the product pages listed above.

Company:

Bayer CropScience (*Raxil PRO* – PCP#30102; *Stress Shield 600* – PCP#30668)

Formulation:

Raxil PRO: 3.0 g per L tebuconazole, 15.4 g per L prothioconazole and 6.2 g per L metalaxyl formulated as a suspension. Container size – 10 L, 58.5 L, 175.5 L, 1000 L.

Stress Shield 600: 600 g per L imidacloprid formulated as a suspension. Container size – 27 L.

Crops:

Wheat, barley and oat.

Diseases and Insects:

Diseases Controlled: Seed rot/pre-emergence damping-off caused by seed-borne *Fusarium* spp., *Aspergillus* spp. and *Cochliobolus sativus*; seed rot/pre-emergence damping-off caused by soil-borne *Fusarium* spp., *Cochliobolus sativus*, and *Pythium* spp.; post-emergence damping-off caused by soil-borne *Fusarium* spp. and seed-borne *Aspergillus* spp.;

seedling blight caused by seed-borne *Aspergillus* spp.; true loose smut (*Ustilago nuda*); covered smut (*U. hordei*, *U. kolleri*); false loose smut (*U. nigra*); loose smut (*U. tritici*, *U. avenae*); common bunt (*Tilletia caries*, *T. laevis*); barley leaf stripe (*Pyrenophora graminea*).

Disease Suppressed: Root rot caused by soil-borne *Fusarium* spp., *Cochliobolus sativus*, and seed-borne *Fusarium* spp.; crown rot caused by seed-borne *Fusarium* spp.; seedling blight caused by seed-borne *Penicillium* spp.

Insects Controlled: Wireworm

Rates:

325 mL *Raxil PRO* and 17 to 50 mL of *Stress Shield 600* per 100 kg of seed.

Hazard Rating:



Warning – Poison (*Stress Shield 600*)



Caution – Skin irritant (*Raxil PRO*)

For an explanation of the symbols used here see page 11.

Raxil WW

**Fungicide Group – 3, 4
Insecticide Group – 4A**

(Refer to page 312 and 456)

Company:

Bayer CropScience

Formulation:

Raxil WW has two components: *Raxil MD* (PCP# 27692) 5.0 g per L tebuconazole and 6.6 g per L metalaxyl formulated as a flowable seed treatment and *Stress Shield* (PCP# 29610) 480 g per L imidacloprid formulated as a suspension.

Crops, Diseases, Insects, and Rates:

Crops:	Diseases:	Insects:	Rate of <i>Raxil MD</i> per 100 kg seed:	Rate of <i>Stress Shield</i> per 100 kg seed:
Wheat (durum, spring, winter), Barley, Oats	For diseases controlled and/or suppressed, refer to <i>Raxil MD</i> (page 432).	Wireworm	300 ml	21 to 63 ml*

* Use the higher rate when infestation pressures are expected to be heavy. DO NOT apply any subsequent application of a Group 4 Insecticide (i.e. in-furrow or foliar application) following treatment with *Stress Shield*.

How it Works:

The active ingredient tebuconazole is a triazole fungicide with broad-spectrum systemic activity. The active ingredient metalaxyl is an acylalane fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311. Imidacloprid is a chloronicotinyl insecticide with systemic activity. For more information refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

None registered.

Restrictions:

See *Raxil MD* (page 432) and consult individual labels for *Raxil MD* and *Stress Shield* for additional information regarding use restrictions.

Resistance management: Refer to page 310.

Labelling: All bags containing treated seed must be labelled or tagged as follows: "This seed has been treated with *Raxil WW* (*Raxil MD* and *Stress Shield*) seed protectant which

contains tebuconazole and metalaxyl fungicides and imidacloprid insecticide. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs."

Grazing: DO NOT graze or feed livestock on treated areas for four weeks after planting.

Re-cropping: No restrictions listed.

Storage: Low temperature storage is not recommended. Prior to and during application, treatments must be thoroughly agitated to ensure uniform mixing of the product. Due to the viscosity of the material, it should be kept above 10 °C prior to and during application. DO NOT apply direct heat to container.

Environment: DO NOT contaminate water, food, or feed by storage, disposal, or by cleaning of equipment. Cover or incorporate spilled treated seeds. Left over treated seed should be double-sown around the headland, or buried away from water sources.

Hazard Rating:



Danger – Skin and eye irritant

For an explanation of the symbols used here see page 11.

Senator PSPT

Fungicide Group – 1

(Refer to page 312)

Company:

Nippon Soda Company Ltd. – PCP#26236

Distributed by Engage Agro Corporation

Formulation:

10% thiophanate-methyl formulated as dust.

Container size - 10 kg.

Crops and Diseases:

Control of *Fusarium* rot, *Verticillium* wilt, silver scurf (*Helminthosporium solani*) of potato. Also aids in control of seed piece decay and blackleg infections of potato.

Rate:

Use 500 g per 100 kg cut seed (one 10 kg bag treats 2,000 kg seed).

Application Information:

Seed piece treatment. Apply in a convenient container or by dust attachment over belt. Cut pieces should be treated within 6 hours of cutting. For optimum control of silver scurf, ensure that seed tubers are completely free of soil. Total skin coverage is essential. If planting is to be delayed more than 1 to 2 days, the treated pieces should be stored for 2 to 3 days in open crates before bagging. This product contains no colourant; an appropriate colourant must be added when this product is applied.

How it Works:

The active ingredient thiophanate-methyl is a benzimidazole fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: No restrictions listed.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store product in a dry place. Avoid contamination of feed or food stuffs.

Environment: DO NOT contaminate domestic or irrigation water supplies, lakes, streams and ponds.

Hazard Rating:

None listed.

StorOx

Fungicide Group – U

(Refer to page 312)

Company:

Manufactured by BioSafe Systems Inc. – PCP# 27432
Distributed in Western Canada by Brenntag Canada

Formulation:

27% hydrogen peroxide. Container size - 10 - 220 L.

Crops and Diseases:

Control of Fusarium tuber rot, bacterial soft rot and silver scurf in potato.

Rate and Application Information:

Prior to storage and in storage treatment for harvested potato tubers.

As a spray treatment for newly harvested potatoes before storage: 100 mL of StorOx per 10 L water. Spray diluted solution on tuber to runoff to achieve full and even coverage. Use 4.15 to 8.3 L water per tonne of potatoes.

As application to potatoes in storage as a direct injection into humidification water: 100 mL StorOx per 10 L water. Apply diluted product for at least 20 minutes per day, based on a humidification airflow rate of 0.6 cfm.

Tank Mixes:

May be used in conjunction with a growth inhibitor during humidification. Should not be combined or mixed with pesticides or fertilizer.

How it Works:

Hydrogen peroxide is an inorganic compound with contact activity against fungi and bacteria. For more information refer to "Fungicide Modes of Action" on page 311.

Restrictions:

Storage: Store in cool, well ventilated area away from direct sunlight. Since StorOx is a strong oxidizing agent, contact with combustibles may cause fire.

Environmental: DO NOT discharge effluent containing StorOx into lakes, streams, ponds or other bodies of water. DO NOT permit this product to enter surface or ground water.

Hazard Rating:



Danger – Corrosive to eyes



Warning – Skin irritant

Other Precautions: This product is corrosive to metal surfaces; rinse all application equipment thoroughly with water after use. Do not enter treated storage bins until the hydrogen peroxide air concentrations are below exposure levels established by occupational health and safety authorities.

For an explanation of the symbols used here see page 11.

Stress Shield 600

Insecticide Group – 4

(Refer to page 456)

Company:

Bayer CropScience – PCP#30668

Formulation:

600 g per L imidacloprid formulated as a suspension.
Contains insecticide only.

Container size - 27 L.

Crops, Insects, and Rates:

Crop:	Rate per 100 Kg seed ¹ :	Insects Controlled:
Wheat, barley, oat	17 to 50 mL ²	Wireworm
Bean	104 mL	Potato leafhopper, wireworm
Soybean	104 to 208 mL ³	Soybean aphid, bean leaf beetle, seedcorn maggot, wireworm
Field pea	104 mL	Wireworm
	104 to 208 mL	Pea leaf weevil
Fababean	104 mL	Pea leaf weevil, wireworm
Chickpea, lentil	104 mL	Wireworm

¹ DO NOT apply any subsequent applications of a Group 4 Insecticide (i.e. in-furrow or foliar application) following treatment with *Stress Shield 600*.

² For fields with a history of moderate to high wireworm pressure, treat crops at 34 to 50 mL per 100 kg seed. Use the higher rate when infestation pressures are expected to be heavy.

³ Use the higher rate for early seeding, when insect populations are expected to be high, or for extended control period of aphids

Application Information:

For use in commercial and on-farm seed treatment equipment. Mix thoroughly before use or use entire container at one time. Apply *Stress Shield 600* through slurry applicator seed treaters which provide uniform seed coverage. Uneven or incomplete seed coverage may not give the desired level of insect control. Maintain constant agitation of the slurry during application. Allow the seed to dry before bagging or storing in bulk containers. All seed must be conspicuously coloured at the time of treatment in accordance with Seed Act and Regulations. Seed treated with *Stress Shield 600* may reduce seed flow in the seed drill. Recalibration of the seed drill may be required to obtain correct seeding rate before planting. *Stress Shield 600* can be used as an over-treatment. If this product is being used alone a dye must be added to the solution.

How it Works:

The active ingredient imidacloprid is a chloronicotinyl insecticide with systemic activity. For more information

refer to "Insecticide Groups Based on Modes of Action" on page 456.

Tank Mixes:

Fungicide Seed Treatments: *Raxil MD*, or *EverGol Energy* in cereals. *Trilex AL*, *Allegiance*, *EverGol Energy*, *Apron Maxx R1C*, or *Apron Maxx RTA* in pulses.

Follow the label directions for each product and use the most restrictive precautions and limitations for either product.

Bayer CropScience also supports the tank-mix of *Trilex EverGol* with *Stress Shield 600*. Apply mixes according to the most restrictive use limitation.

Restrictions:

Resistance management: Refer to page 455.

Labelling: Treated seed must be labeled "This seed has been treated with *Stress Shield 600* seed protectant which

contain imidacloprid. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs. Toxic to birds. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface."

Grazing: DO NOT graze or feed livestock on treated areas for four weeks after planting.

Re-cropping: No restrictions listed.

Storage: Store product in original container only, away from other pesticides, fertilizer, food, or feed. Keep container closed. DO NOT store in direct sunlight nor above 35°C.

Environment: Toxic to birds and aquatic organisms. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Thiram 75WP

Fungicide Group – M3
(Refer to page 312)

Company:

Manufactured by Chemtura Canada Co.: *Thiram 75WP* – PCP#27556

Formulation:

Thiram 75WP: 75% thiram formulated as wettable powder. Container sizes - 5 kg, 25 kg.

Diseases Controlled:

Seed decay, seedling blight, damping off.

Crops and Rates:

Crop:	Rate per 100 kg Seed:	Amount of Seed one 5 Kg Package Treats:
Mustard, grasses, alfalfa	360 g	1,389 kg
Dry bean, pea, soybean	100 to 140 g	3,571 to 5,000 kg
Field corn	120 g	4,166 kg
Sweet corn	220 g	2,272 kg
Safflower	200 g	2,500 kg

Application Information:

To apply as slurry treatment, pre-mix *Thiram 75WP* in water (as indicated below) and apply through commercial seed treating equipment.

How it Works:

The active ingredient thiram is a dithiocarbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed should be labelled "DO NOT use for food or feed. This seed has been treated with thiram".

Grazing: DO NOT graze treated areas or feed clippings from treated areas to livestock.

Re-cropping: No restrictions listed.

Storage: Store in a cool, dry, ventilated place, away from feeds and foods. Keep away from flame, sparks and heat.

Environment: Treated seed may be harmful to birds if ingested. DO NOT contaminate any body of water.

Compatibility with rhizobia-based inoculants: *Thiram* is compatible with rhizobia, however, some restrictions may apply in storage length with *Thiram*-treated seed inoculated with rhizobia. Contact rhizobia manufacturer on use patterns with their rhizobia strain.

Hazard Rating:



Warning – Poison

Other precautions: May irritate eyes, nose, throat and skin. May cause allergic eczema in sensitive individuals. Do not consume alcoholic beverages 24 hours before or after working with thiram.

For an explanation of the symbols used here see page 11.

Trilex AL

Fungicide Group – 11, 4

(Refer to page 312)

Company:

Bayer CropScience (PCP#29160)

Formulation:

13.5 g per L trifloxystrobin and 10.8 g per L metalaxyl formulated as a suspension concentrate.

Crops and Disease:

Protection of bean, chickpea, pea, lentil and soybean seed and seedlings from certain seed and soil-borne diseases caused by *Rhizoctonia solani*, *Fusarium* spp., and *Pythium* spp. Also protects from seed rot / pre-emergence damping-off, post-emergence damping-off, and seedling blight caused by *Botrytis cinerea* on lentil; and seed decay / pre-emergence damping-off caused by *Phomopsis longicolla* on soybean. It also provides suppression of seed-borne *Ascochyta* blight caused by *Ascochyta* spp on lentil, pea and chickpea.

Rate:

370 ml per 100 kg of seed.

Application Information:

The ready-to-use formulation is designed for commercial or on-farm treating with conventional seed treating equipment that can accurately control application rates and provide a good distribution of the chemical onto the seed in the mixing chamber. Uniform application on seed is necessary to ensure seed safety and best disease protection. If *Trilex AL* is diluted with water by greater than 10% by volume, ensure agitation of the mixture prior to application to seed.

How it Works:

Trifloxystrobin is a strobilurin fungicide with broad spectrum preventative activity. Metalaxyl is an acylalanine fungicide with systemic activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: All bags containing treated seed for sale or use in Canada must be labelled or tagged as follows: "This seed has been treated with *Trilex AL* containing trifloxystrobin and metalaxyl. Use chemical-resistant gloves when handling treated seed. Do not use for feed, food or oil processing. Store away from feeds and other foodstuffs."

Grazing: DO NOT graze or feed livestock on treated areas for 4 weeks after planting.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Store in a cool, dry place and avoid excessive heat. Keep container closed.

Environment: Toxic to aquatic organisms. DO NOT discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans or other waters. Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. Residues of this product demonstrate the properties and characteristics associated with chemicals detected in groundwater. The use of this product may result in contamination of groundwater particularly in areas where soils are permeable (e.g., sandy soil) and/or depth to the water table is shallow.

Compatibility with rhizobia-based inoculants: *Trilex AL* is compatible with rhizobia-based inoculants. Check with inoculant manufacturers for further details prior to use.

Hazard Rating:

None listed.

Trilex EverGol

Fungicide Group – 4, 7, 11

(Refer to page 312)

Trilex EverGol is a co-pack of *Trilex Component A* (penflufen and trifloxystrobin) and *Trilex Component B* (metalaxyl).

Company:

Bayer CropScience (*Trilex Component A* – PCP#30644; *Trilex Component B* – PCP#30645)

Formulation:

Trilex Component A: 154 g per L penflufen and 154 g per L trifloxystrobin formulated as a suspension.

Container size – 1.5L *Trilex Component A* or in bulk package 8L

Trilex Component B: 317 g per L metalaxyl formulated as a suspension.

Container size – 0.96L *Trilex Component B* or in bulk package 5.12 L.

Crops, Diseases and Rates:

Crop:	Rate per 100 kg of seed:		Diseases Controlled:		Diseases Suppressed:	
	<i>Trilex Component A</i>	<i>Trilex Component B</i>	<i>Trilex Component A</i>	<i>Trilex Component B</i>	<i>Trilex Component A</i>	<i>Trilex Component B</i>
Bean	25 mL	16 mL	Seed decay / pre-emergence damping-off and post-emergence damping-off caused by <i>Rhizoctonia solani</i> , <i>Fusarium</i> spp., and <i>Botrytis cinerea</i> ; seedling blight caused by <i>B. cinerea</i>	Seed rots and seedling blights caused by <i>Pythium</i> spp.	-	-
	25 to 32 mL	-	-	-	Seedborne <i>Ascochyta</i> blight caused by <i>Ascochyta</i> spp.	-
Chickpea, pea	25 mL	16 mL	Seed decay / pre-emergence damping-off and post-emergence damping-off caused by <i>Rhizoctonia solani</i> , <i>Fusarium</i> spp., and <i>Botrytis cinerea</i> ; seedling blight caused by <i>B. cinerea</i>	Seed rots and seedling blights caused by <i>Pythium</i> spp.	-	-
	25 to 32 mL	-	-	-	Seedborne <i>Ascochyta</i> blight caused by <i>Ascochyta</i> spp.	-

See footnotes on following page.

Continued...

Crops, Diseases, and Rates *continued*:

Crop:	Rate per 100 kg of seed:		Diseases Controlled:		Diseases Suppressed:	
	Trilex Component A	Trilex Component B	Trilex Component A	Trilex Component B	Trilex Component A	Trilex Component B
Lentil	25 mL	16 mL ¹	Seed decay / pre-emergence damping-off and post-emergence damping-off caused by <i>Rhizoctonia solani</i> , <i>Fusarium</i> spp., and <i>Botrytis cinerea</i> ; seedling blight caused by <i>B. cinerea</i>	<i>Pythium</i> seed rot	-	-
	25 to 32 mL	-	-	-	Seedborne <i>Ascochyta</i> blight caused by <i>Ascochyta</i> spp.	-

¹ Trilex Component B is for use on low tannin lentils destined for export or seed production only and no foliar application of metalaxyl can be used on low-tannin lentils grown from metalaxyl-treated seed.

Application Information:

Trilex Component A is a seed treatment formulation for use in commercial seed treatment operations, and for on-farm treating with conventional seed treating equipment which can accurately meter and apply flowable seed treatment formulations. This product is recommended to be diluted with water or another suitable liquid just prior to application to ensure uniform coverage on the seed during the application process. Uniform application to seed is necessary to ensure optimum performance. Allow seeds to dry before bagging, storing or seeding.

Trilex Component B should be mixed with water to form a slurry seed treatment. Mix 500 mL of slurry per 100 kg of seed to be treated. The slurry should be applied as a spray into the mixing chamber of the seed treating equipment to ensure good coverage. When preparing the slurry the following procedure should be used: 1) partially fill the mixing tank with water; 2) add the required quantity of **Trilex Component B** onto the water surface; 3) allow product to disperse and then switch on agitation; 4) top up with extra water to required volume and maintain agitation during use; and 5) add colourant last.

How it Works:

Trilex Component A: The active ingredient penflufen is a carboxamide (SDHI) fungicide with systemic activity. The active ingredient trifloxystrobin is a strobilurin fungicide with broad spectrum preventative activity.

Trilex Component B: The active ingredient metalaxyl is an acylalanine fungicide with systemic activity.

For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Fungicide Seed Treatments: In addition to **Trilex Component B**, **Trilex Component A** may be mixed with **Allegiance FL**.

Follow the label directions for each product and use the most restrictive precautions and limitations for either product.

Bayer CropScience also supports the tank-mix of Stress Shield 600 with Trilex EverGol. Apply mixes according to the most restrictive use limitations.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labeled "This seed has been treated with **Trilex Component A** (containing penflufen and trifloxystrobin) and **Trilex Component B** (containing metalaxyl). Wear long-sleeved shirt, long pants and chemical-resistant gloves when handling treated seed. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs."

Grazing: DO NOT graze or feed livestock on treated areas for four weeks after planting.

Re-cropping: Registered crops for **Trilex Component A**, as well as canola, mustard, rapeseed, soybean, alfalfa, corn and cereal grains, may be replanted at any time. For all other crops, DO NOT plant back within 30 days of seeding with **Trilex Component A**-treated seed.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Keep container closed. Store in a cool, dry area. Do not store in direct sunlight. Do not store **Trilex Component A** above 40°C or below -10°C. Do not store **Trilex Component B** above 35°C or below 0°C.

Environment: Toxic to aquatic organisms. Treated seed may be toxic to birds and other wildlife. DO NOT discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans or other waters. Dispose of all excess treated seed. Left over seed may be double-sown around the headland or buried away from water sources. DO NOT leave exposed treated seed on soil surface. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable and/or the depth to the water is shallow.

Hazard Rating:



Warning – Skin and eye irritant (*Trilex Component B*)

For an explanation of the symbols used here see page 11.

Triticonazole

Charter RTU / Armour RTU / Armour

Fungicide Group – 3

(Refer to page 312)

Company:

BASF Canada (*Charter RTU* – PCP#29400)

Loveland Products (formerly Viterra) (*Armour RTU* – PCP#30226, *Armour* – PCP#29296)

200 L drums.

Armour RTU – 16.8 g per L triticonazole formulated as a liquid flowable seed treatment. Container size - 9.3 L.

Armour – 25 g per L triticonazole formulated as a liquid flowable seed treatment. Container size - 10 L (containing 6.2 L *Armour*).

Formulation:

Charter RTU – 16.8 g per L triticonazole formulated as a liquid flowable seed treatment. Container size - 9.3 L and

Crops, Diseases and Rates:

Crop:	Rate per 100 kg Seed: <i>Charter RTU, Armour RTU</i> ¹	Rate per 100 kg Seed: <i>Armour</i>	Diseases Controlled:	Diseases Suppressed ² :
Wheat	300 mL	200 mL	Seed rot caused by <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Fusarium</i> spp.; loose smut (<i>Ustilago tritici</i>); common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>)	<i>Fusarium</i> crown and root rot; common root rot (caused by <i>Cochliobolus</i>); seedling blight caused by <i>Cochliobolus sativus</i>
Barley	300 mL	200 mL	Seed rot caused by <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Fusarium</i> spp.; true loose smut (<i>Ustilago nuda</i>); covered smut (<i>U. hordei</i>); false loose smut (<i>U. nigra</i>)	
Oat	300 mL	200 mL	Seed rot caused by <i>Fusarium</i> spp.; seedling blight caused by seed-borne <i>Fusarium</i> spp.; loose smut (<i>Ustilago avenae</i>); covered smut (<i>U. kolleri</i>)	

¹ *Charter RTU* and *Armour RTU* are ready to use formulations. No water or dye is required to be added.

² Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

Application Information:

Charter RTU and *Armour RTU* are ready to use seed treatment formulations and *Armour* is a concentrated seed treatment formulation. These products are for use in commercial seed plants, in on-farm standard gravity flow or mist type treatment machines, and in on-the-go air seeder treatment systems. Agitate or shake well prior to usage. Uneven seed coverage may result in poor levels of disease control. Seed should be well conditioned and cleaned before treating. Treated seed should not require drying after treatment.

Charter RTU and Armour RTU Water Volume: *Charter RTU* and *Armour RTU* are ready to use formulations and no additional water is required. Consult the seed treatment application equipment manufacturer for specific application instructions for use of various seed treatment application machines.

Armour Water Volume: Concentrate requires the addition of water. Mix 2 parts seed treatment with 1 part water. Consult the seed treatment application equipment manufacturer for specific application instructions for use of various seed treatment application machines.

How it Works:

The active ingredient triticonazole is a triazole fungicide that provides systemic broad spectrum protection against seed- and soil-borne diseases. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed should be labelled "This seed has been treated with (*Charter RTU*, *Armour RTU*, or *Armour*) Seed Treatment Fungicide containing triticonazole, a Group 3 fungicide. DO NOT use for food, feed or oil processing". Store treated seed under cool, dry conditions.

Grazing: No restrictions listed.

Re-cropping: No restrictions listed.

Storage: Store product in the original container. Keep container closed. Any seed treated with this product must be either thoroughly stained with a conspicuous color or coated with a material that renders it conspicuous. DO NOT store treated seed for more than 18 months. Treated seed stored for more than 6 months should be tested for germination before use. Thoroughly clean auger after handling treated seed before using same auger for handling commercial or feed grains.

Environment: DO NOT feed treated seed to, or otherwise expose, wildlife or domestic birds. DO NOT contaminate domestic or irrigation water supplies, lakes, streams, ponds or any body of water with the chemical, used containers, treated seed or bags. DO NOT contaminate water by cleaning of equipment or disposal of wastes.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Vibrance 500FS

Fungicide Group – 7

(Refer to page 312)

Vibrance 500FS Seed Treatment is currently only available as a co-pack (see **Cruiser Maxx Vibrance Beans**, **Helix Vibrance**, **Vibrance Maxx RTA/RFC**, **Vibrance XL Co-Pack**).

Company:

Syngenta Canada (PCP #30438)

Formulation:

500 g per L sedaxane formulated as a suspension.

Container size – 1L to 1050 L

Crops and Diseases:

Crop:	Diseases Controlled:
Barley	True loose smut caused by <i>Ustilago nuda</i> ; seed decay, seedling blight and damping-off caused by <i>Rhizoctonia solani</i>
Wheat	True loose smut caused by <i>Ustilago tritici</i> ; seed decay, seedling blight and damping-off caused by <i>Rhizoctonia solani</i>
Oats, rye, and triticale	Seed decay, seedling blight and damping-off caused by <i>Rhizoctonia solani</i>
Canola, soybeans, chickpea, lentil and pea	Seed decay, seedling blight and damping-off caused by <i>Rhizoctonia solani</i>

Rate:

5 to 10 mL product per 100 kg of seed. Use the low rate for control of pre-emergent damping-off, seedling decay, or seedling blight. Use the high rate for extended control of post-emergent damping-off and seedling blight or high disease pressure or high levels of seed-borne infections like smut.

Application Information:

Vibrance 500FS is for use on-farm and in closed transfer commercial seed treatment facilities. No open transfer is permitted for commercial seed treatment of barley, wheat, oats, rye and triticale. No on-farm seed treatment is permitted for canola and soybean.

Note: treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour.

This product contains no colourant. An appropriate colourant must be added when this product is applied to seed. Regulations pertaining to the "Seeds Act" must be strictly adhered to when using this product. Users are responsible for ensuring that the treated seed, when dried and ready for bagging, has an unnatural colour.

How it Works:

Sedaxane is a succinate dehydrogenase inhibitor fungicide with preventative and systemic activity that inhibits fungal metabolism by binding to the succinate dehydrogenase enzyme to disrupt cellular respiration and energy generation. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Small-grain cereals (Wheat, Barley, Oats, Rye, and Triticale): **Vibrance 500FS** may be mixed with **Dividend XL RTA Fungicide** or **Dividend Extreme Fungicide** for broad spectrum disease control of cereals. For protection from various insects, **Vibrance 500FS** may be mixed with **Cruiser 5FS** or **350FS Seed Treatment Insecticides** in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. This tank-mix option is only valid for those crops common to the registered labels of both products. For combined protection from labeled insect pests and soil and seed-borne pathogens, **Vibrance 500FS** may be mixed with **Cruiser Maxx Cereals Seed Treatment** or **Cruiser Maxx Cereals Commercial Seed Treatment**. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

Canola: For additional disease control and protection from labeled insects, **Vibrance 500FS** may be mixed with **Helix Xtra Seed Treatment** in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

Soybeans: **Vibrance 500FS** may be mixed with **Apron Maxx RTA** or **Apron Maxx RFC Seed Treatments** and **Dynasty 100FS Fungicide** in commercial seed treatment facilities to control a broad spectrum of diseases. For protection from various insect pests, **Vibrance 500FS** may be mixed

with *Cruiser 5FS* or *350FS Seed Treatment Insecticides* in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. For combined protection from labeled insect and soil- and seed-borne pathogens, *Vibrance 500FS* may be mixed with *Cruiser Maxx Beans Seed Treatment* under the same "closed system" conditions as noted above. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

Restrictions:

Resistance management: Refer to page 310.

Labelling: All seed must be labelled "This seed has been treated with sedaxane fungicide. Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed. Do not graze or feed livestock on seeded area for 45 days after planting. Do not use for food, feed or oil processing. Store away from food and feed".

Grazing: DO NOT graze or feed livestock on treated areas for 45 days after planting.

Re-cropping: DO NOT plant any crop other than those on the product label within 60 days to fields in which seed treated with *Vibrance 500FS* seed treatment were planted.

Storage: Store away from food and feed.

Compatibility with rhizobia-based inoculants: *Vibrance 500FS* is compatible with Rhizobium based inoculants. Please check with inoculant manufactures for details prior to use. Note: Mixing with inoculants may increase drying time while treating extending the processing time.

Environment: Toxic to aquatic organisms. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. Treated seed is toxic to small wild animals. Any spilled or exposed seed must be incorporated into the soil or otherwise cleaned up from the soil surface.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Vibrance Maxx RTA Vibrance Maxx RFC

Fungicide Group – 4, 7, 12
(Refer to page 312)

Vibrance Maxx RTA is a co-pack of *Apron Maxx RTA* (page 392) and *Vibrance 500 FS* (page 446). *Vibrance Maxx RFC* is a co-pack of *Apron Maxx RFC* (page 392) and *Vibrance 500 FS* (page 446). For other detailed information on the component products see the product pages listed above.

Company:

Syngenta Canada

Formulation:

Vibrance Maxx RTA: *Apron Maxx RTA* (PCP# 27577): 0.73% fludioxonil plus 1.10% metalaxyl- M formulated as a liquid seed treatment and *Vibrance 500 FS* (PCP #30438) 500 g/L sedaxane formulated as a suspension.

Container size – 2 x 10 L + 2 x 300 ml; 115 L + 3.33 L; 450 L + 4 X 3.33L

Vibrance Maxx RFC: *Apron Maxx RFC* (PCP# 28817) 2.31% fludioxonil plus 3.46% metalaxyl- M and S-isomers as a liquid seed treatment and *Vibrance 500 FS* (PCP #30438) 500 g/L sedaxane formulated as a suspension.

Container size – 56.78 L + 1.45 L

Crops:

Peas, lentil, chickpeas

Diseases and Insects:

Diseases Contolled:

Seedling blights, damping-off, and seed rots caused by *Pythium*, *Fusarium*, and *Rhizoctonia* spp. Seed-borne ascochyta blight (*Ascochyta* spp.); seed rot and seedling blight caused by seed-borne *Botrytis* spp.

Rates:

Vibrance Maxx RTA: 325 mL *Apron Maxx RTA* and 5 to 10 mL *Vibrance 500 FS* per 100 kg of seed.

Vibrance Maxx RFC: 100 mL *Apron Maxx RFC* and 5 to 10 mL *Vibrance 500 FS* per 100 kg of seed.

Vibrance XL/ Vibrance XL Co-Pack

Fungicide Group – 3, 4, 7
(Refer to page 312)

Vibrance XL Co-Pack is a co-pack of **Dividend XL RTA** (difenoconazole and metalaxyl-M, page 405) and **Vibrance 500FS** (sedaxane, page 446). Information listed is limited to crop, diseases, insects and rates. For other detailed information on the component products see the product pages listed above.

Company:

Syngenta Canada

Formulation:

Vibrance XL (PCP #30437): 66.2 g per L difenoconazole, 16.5 g per L metalaxyl-M (and S-isomer), and 13.8 g per L sedaxane formulated as a suspension. Container size – 2 x 5.54 L

Vibrance XL Co-Pack has two components:

Dividend XL RTA (PCP#25777): 3.37% difenoconazole, 0.27% metalaxyl-M formulated as a flowable seed treatment. **Vibrance 500FS** (PCP#30438): 500 g per L sedaxane formulated as a suspension. See the component products for other information. Use the most stringent restrictions for either product. Container size – 115 L **Dividend XL RTA** + 1.77 L **Vibrance 500FS** or 450 L **Dividend XL RTA** + 4x1.77 L **Vibrance 500FS**

Crops and Diseases:

Crop:	Diseases Controlled:	Diseases Suppressed ²
Barley	General seed rots ¹ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp or soil-borne <i>Pythium</i> ; seed-borne <i>Septoria</i> ² ; covered smut (<i>Ustilago hordei</i>), false loose smut (<i>U. nigra</i>), and true loose smut (<i>U. nuda</i>).	Common root rot (caused by <i>Cochliobolus</i> spp.); <i>Fusarium</i> crown and foot rot; take-all (<i>Gaeumannomyces graminis</i> var. <i>tritici</i>)
Winter Wheat and Spring Wheat	General seed rots ¹ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp or soil-borne <i>Pythium</i> ; loose smut (<i>Ustilago tritici</i>), seed-borne and soil-borne common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>).	
Winter Wheat only	Dwarf bunt (<i>Tilletia controversa</i>); seed-borne <i>Septoria</i> and early season foliar disease control of <i>Septoria</i> leaf blotch for first 4 weeks after planting (for full season control apply a foliar fungicide according to label directions).	
Rye	General seed rots ¹ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp or soil-borne <i>Pythium</i> ; seed-borne <i>Septoria</i> ² ; seed-borne and soil-borne common bunt (<i>Tilletia caries</i>) and dwarf bunt (<i>T. controversa</i>).	
Triticale	General seed rots ¹ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp or soil-borne <i>Pythium</i> ; loose smut (<i>Ustilago tritici</i>).	Common root rot (<i>Cochliobolus</i> spp.)
Oats	General seed rots ¹ ; seedling blight, root rot, and damping-off caused by seed- and soil-borne <i>Fusarium</i> or <i>Rhizoctonia</i> spp or soil-borne <i>Pythium</i> ; covered smut (<i>Ustilago kolleri</i>) and loose smut (<i>U. avenae</i>)	

¹ General seed rots controlled include those caused by saprophytic organisms such as *Fusarium*, *Pythium*, *Penicillium* and *Aspergillus*.

² Suppression means consistent control at a level which is not optimal but is still of commercial benefit.

Rate:

Vibrance XL: 180 to 360 mL product per 100 kg of seed. Use the high rate if there is a history of high disease pressure in the field, when field conditions favour seed-borne and

soil-borne pathogens, when planting seed with high levels of seed-borne infections like smut, for extended control of post-emergent damping-off, or when controlling seed-borne *Septoria* or early season *Septoria* leaf blotch.

Application Information:

Vibrance XL is for use on-farm and commercial seed treatment facilities. No open transfer is permitted for commercial seed treatment of barley, wheat, oats, rye and triticale.

Uneven or incomplete seed coverage may not give the desired level of disease control. In order to ensure uniform coverage with the 5.54 L *Vibrance XL* containers, add water to the 10 L level and mix thoroughly to form a slurry. Adjust application rate accordingly.

Note: treatment of highly mechanically scarred or damaged seed or seed known to be of low vigour and poor quality may result in reduced germination and/or reduction of seed and seedling vigour.

This product contains a pigment that will adequately colour treated seed. However, ensure that treated seed has an unnatural colour after treatment, when dried and ready for bagging. Regulations pertaining to the *Seeds Act* must be strictly adhered to when using this product. An appropriate colourant must be added when this product if the pigment does not colour the seed adequately.

How it Works:

Sedaxane is a succinate dehydrogenase inhibitor fungicide with preventative and systemic activity that inhibits fungal metabolism by binding to the succinate dehydrogenase enzyme to disrupt cellular respiration and energy generation. The active ingredient difenoconazole is a triazole fungicide with broad-spectrum, systemic activity. Metalaxyl-M is an acylalanine fungicide with systemic activity against diseases caused by the Oomycetes class, including *Pythium* damping-off. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

Small-grain cereals (Wheat, Barley, Oats, Rye, and Triticale): *Vibrance 500FS* may be mixed with *Dividend XL RIA Fungicide* or *Dividend Extreme Fungicide* for broad spectrum disease control of cereals. For protection from various insects, *Vibrance 500FS* may be mixed with *Cruiser 5FS* or *350FS Seed Treatment Insecticides* in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. This tank-mix option is only valid for those crops common to the registered labels of both products. For combined protection from labeled insect pests and soil and seed-borne pathogens, *Vibrance 500FS* may be mixed with *Cruiser Maxx Cereals Seed Treatment* or *Cruiser Maxx Cereals Commercial Seed Treatment*. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

Canola: For additional disease control and protection from labeled insects, *Vibrance 500FS* may be mixed with *Helix Xtra Seed Treatment* in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment

only. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

Soybeans: *Vibrance 500FS* may be mixed with *Apron Maxx RIA* or *Apron Maxx RFC Seed Treatments* and *Dynasty 100FS Fungicide* in commercial seed treatment facilities to control a broad spectrum of diseases. For protection from various insect pests, *Vibrance 500FS* may be mixed with *Cruiser 5FS* or *350FS Seed Treatment Insecticides* in commercial seed treatment facilities with closed transfer including closed mixing, loading, calibrating, and closed treatment equipment only. For combined protection from labeled insect and soil- and seed-borne pathogens, *Vibrance 500FS* may be mixed with *Cruiser Maxx Beans Seed Treatment* under the same "closed system" conditions as noted above. Consult each product label for registered use rates and follow all label use instructions. Read the label directions for each product and follow the most restrictive label precautions and limitations.

Restrictions:

Resistance management: Refer to page 310. Experience has shown that strains of fungus resistant to metalaxyl-M (and S-isomer) may develop. Failure to control the disease will result in crop damage and/or yield losses. If disease appears in a treated field, consult the government extension specialist immediately.

Labelling: All seed must be labelled "This seed has been treated with difenconazole, metalaxyl-M (and S-isomer), and sedaxane fungicides. Wear long-sleeved shirt, long pants, and chemical-resistant gloves when handling treated seed. Do not graze or feed livestock on seeded area for 45 days after planting. Do not use for food, feed or oil processing. Store away from food and feed".

Grazing: DO NOT graze or feed livestock on treated areas for 45 days after planting.

Re-cropping: DO NOT plant any crop other than those on the product label within 60 days to fields in which seed treated with *Vibrance XL* were planted.

Storage: Store away from food and feed. Ideal storage is above freezing and below 30C.

Environment: Toxic to aquatic organisms. Do not contaminate food, feed, domestic or irrigation water supplies, lakes, streams and ponds. Treated seed is toxic to small wild animals. Any spilled or exposed seed must be incorporated into the soil or otherwise cleaned up from the soil surface.

Hazard Rating:



Caution - Poison

For an explanation of the symbols used here see page 11.

Vitaflo Brands

Fungicide Group – 7, M3
(Refer to page 312)

Company:

Manufactured by Chemtura Canada Co
(Vitaflo 280 PCP#11423; Vitaflo 220 PCP#21174)

Manufactured by Interprovincial Cooperative Limited
(Vitaflo SP Fungicide PCP#30381)

Manufactured for Loveland Products (formerly Viterra)
by Interprovincial Cooperative Limited (Vitaflo Fungicide
PCP#30380)

Formulation:

Vitaflo 280/Vitaflo Fungicide/Vitaflo SP Fungicide: 15.59% carbathiin and 13.25% thiram formulated as a liquid suspension.

Container sizes - 10 L, 55L, 100 L, 200 L, 1000 L.

Vitaflo 220: 220 g/L carbathiin and 200 g/L thiram formulated as a liquid suspension. Container size - 10 L.

Crops, Diseases and Rates:

Crop:	Vitaflo 280/ Vitaflo Fungicide/ Vitaflo SP Fungicide Rate per 100 kg Seed:	Vitaflo 220 Rate per 100 kg Seed:	Diseases Controlled:	Diseases Suppressed ⁴ :
Barley	230 to 330 ¹ mL	250 mL	False loose smut (<i>Ustilago nigra</i>); true loose smut (<i>U. nuda</i>); covered smut (<i>U. hordei</i>); leaf stripe (<i>Pyrenophora graminea</i>); seed rot and seedling blight caused by <i>Pythium</i> spp., <i>Penicillium</i> spp., <i>Fusarium</i> spp. and <i>Cochliobolus sativus</i> . Seed rot due to storage fungi <i>Aspergillus</i> and <i>Alternaria</i> spp.	Suppression of root rot caused by <i>Cochliobolus sativus</i> and <i>Fusarium</i> spp.; suppression of net blotch (<i>Pyrenophora teres</i>)
Wheat	230 to 330 ¹ mL	250 mL	Loose smut (<i>Ustilago tritici</i>); common bunt (<i>Tilletia tritici</i> , <i>T. laevis</i>); seed-borne dwarf bunt (<i>T. controversa</i>); seed-borne <i>Septoria</i> ; seed rot and seedling blight caused by <i>Cochliobolus sativus</i> , <i>Fusarium</i> , <i>Pythium</i> , <i>Aspergillus</i> , <i>Penicillium</i> and <i>Alternaria</i> spp. Seed rot due to storage fungi <i>Aspergillus</i> and <i>Alternaria</i> spp.	Suppression of root rot caused by <i>Cochliobolus sativus</i> and <i>Fusarium</i> spp.
Oat	330 mL	250 mL	Loose smut (<i>Ustilago avenae</i>); covered smut (<i>U. kolleri</i>); seed rot seedling blight caused by <i>Fusarium</i> , <i>Pythium</i> , <i>Penicillium</i> . Seed rot due to storage fungi <i>Aspergillus</i> and <i>Alternaria</i> spp.	Suppression of root rot caused by <i>Cochliobolus sativus</i>
Rye	230 to 330 ¹ mL	250 mL	Stem smut (<i>Urocystis occulta</i>); damping off; seed rot and seedling blight caused by <i>Fusarium</i> , <i>Pythium</i> , <i>Penicillium</i> and <i>Cochliobolus sativus</i> . Seed rot due to storage fungi <i>Aspergillus</i> and <i>Alternaria</i> spp.	Suppression of root rot caused by <i>Cochliobolus sativus</i> and <i>Fusarium</i> spp.
Triticale	200 mL	—	Seed rot, damping off and seedling blight caused by <i>Fusarium</i> , <i>Pythium</i> , <i>Penicillium</i> , and <i>Cochliobolus sativus</i> .	—
Dry bean	260 mL	195 mL	Early season seed rot, seedling blight, and root rot caused by <i>Rhizoctonia solani</i> ; Seed-borne anthracnose (<i>Colletotrichum lindemuthianum</i>). Will not protect from wind borne spores. This product will not control anthracnose if seed is severely infected.	—

See footnotes on following page.

Continued...

Crops, Diseases and Rates *continued*:

Crop:	Vitaflo 280/ Vitaflo Fungicide/ Vitaflo SP Fungicide Rate per 100 kg Seed:	Vitaflo 220 Rate per 100 kg Seed:	Diseases Controlled:	Diseases Suppressed ⁴ :
Corn (Field & Sweet)	280 560 to 748 ² mL	210 420 to 565 mL	Seed rot and damping off caused by <i>Fusarium</i> , <i>Pythium</i> , and <i>Penicillium</i> spp. and seed-borne head smut (<i>Sporisorium holci-sorghii</i>). Will not control soil-borne head smut.	—
Flax	525 mL	395 mL	Seed rot, root rot and seedling blight caused by <i>Rhizoctonia solani</i> and <i>Fusarium</i> spp. DO NOT use slurry treatment for flax, use dilution, according to Vitaflo220 Operating Manual, available from dealers.	—
Lentil	330 mL	—	Seed rot, seedling blight, and early season root rot caused by <i>Botrytis cinerea</i> , <i>Rhizoctonia solani</i> , <i>Fusarium</i> and <i>Pythium</i> spp.	—
Pea	260 to 330 ³ mL	—	Seed rot and seedling blight caused by <i>Ascochyta</i> (<i>Mycosphaerella</i>), <i>Rhizoctonia solani</i> , <i>Fusarium</i> and <i>Pythium</i> spp.	—
Soybean	260 mL	195 mL	Seed rot and seedling blight caused by <i>Rhizoctonia solani</i> , <i>Phomopsis</i> and <i>Fusarium</i> spp.	—

¹ The low rate will give partial control of true loose smut in wheat/barley and stem smut in rye. Use the high rate for septoria, seed rot and seedling blight, and suppression of root rot. Also use the high rate (330 mL per 100 kg) for dwarf bunt control in winter wheat.

² Use 560-750 mL per 100 kg seed to control head smut of corn.

³ Use high rate (330 mL per 100 kg) on pea seed to control ascochyta (*Mycosphaerella pinodes*).

⁴ Suppression means consistent control at a level which is not optimal but is still of economic benefit.

Application Information:

Designed to be used undiluted in commercial seed treaters. Undiluted product can be used at temperatures down to -20°C. Centrifugal pumps are not recommended for pumping product. Use of a Vitaflo Pump is recommended. If containers have been in storage, some settling may occur and require agitation.

How it Works:

The active ingredient carbathiin is a carboximide fungicide with systemic activity and the active ingredient thiram is a dithiocarbamate fungicide with contact activity. For more information refer to "Fungicide Modes of Action" on page 311.

Tank Mixes:

None registered.

Restrictions:

Resistance management: Refer to page 310.

Labelling: Treated seed must be labelled as follows "This seed has been treated with Vitaflo 280, Vitaflo 220, Vitaflo Fungicide, or Vitaflo SP Fungicide liquid seed protectant containing carbathiin and thiram. Do not use for feed, food, or oil processing."

Grazing: DO NOT graze or feed livestock on treated area for four weeks after planting except for the following crops: Soybean - DO NOT graze or feed livestock on forage and hay on treated areas; Bean - DO NOT graze or feed on bean forage for 60 days; Barley, oat, wheat - DO NOT graze or feed on treated area for 6 weeks.

Re-cropping: No restrictions listed.

Storage: DO NOT store product in direct sunlight or above 35°C. Will not freeze even at extreme temperatures. If containers have been stored for several months, shake well before using. DO NOT store dry beans, peas, lentils,

or soybeans treated with *Vitaflo 280/Vitaflo Fungicide/Vitaflo SP Fungicide*. Wheat, barley, rye, oats, triticale and flax seed treated with *Vitaflo 280/Vitaflo Fungicide/Vitaflo SP Fungicide* can be stored up to 18 months without reduction in germination. Corn seed treated with *Vitaflo 280/Vitaflo Fungicide/Vitaflo SP Fungicide* can be stored up to one year without reduction in germination.

Environment: DO NOT contaminate ponds, lakes or streams.

Compatibility with rhizobia-based inoculants: *Vitaflo 280, Vitaflo 220, Vitaflo Fungicide, and Vitaflo SP Fungicide* are compatible with rhizobia. Do not tank mix *Vitaflo 280, Vitaflo Fungicide, or Vitaflo SP Fungicide* and rhizobia. Always check with rhizobia manufacturers on any restrictions that may exist with seed treatments.

Hazard Rating:



Warning – Eye Irritant



Caution – Skin Irritant

For an explanation of the symbols used here see page 11.

Insect Control

Additional Resources

For additional information on monitoring, economic thresholds and biological control of insects in field crops, as well as information on insect management in commodities other than those covered in this guide see the WCCP Guide to Integrated Control of Plant Pests at:

<http://www.westernforum.org/wccp%20guidelines.html>

Insect Management Decisions

Crop rotations, cultivar selections, and seeding dates can be chosen to reduce the risk of injury from some insects that may be of higher risk to a crop. Management of insects with insecticides should only be considered when numbers or damage exceed economic thresholds. To select an insecticide, verify the registered products for the insect and field crop in the following tables. Consideration should then be given to the preharvest intervals, how the product will be applied, restrictions, precautions and the hazard rating.

Preharvest Interval

The **preharvest interval** is the number of days that must pass between the last application of a pesticide and harvest. Harvest is the cutting of the crop or removal of the produce from the plant. It includes direct-combining, cutting (swathing) or grazing; it does not include swath-combining or baling for hay.



Field Scouting

Field scouting is the regular examination of fields to accurately assess the kind and the number of insects, plant pathogens and weeds present and the amount of damage being done. Scouting should be done weekly during the growing season and more frequently when infestations approach economic levels or when weather conditions favour the rapid development of specific pests.

To properly scout for insect pests, you must know when they occur, where they live, what they look like, and how to find and count them. The number of locations to assess in a field will depend on the field size, and any specific pests that may be of concern. Generally a minimum of 5 sites should be sampled, however some insects may require more sites to be sampled to accurately make management decisions. There are several possible scouting patterns that can be used when checking fields. These options are based on pest distribution and field configuration.

Pattern 1: Used when pests are uniformly distributed.

This scouting pattern typically looks like an X, Z or W, excluding field edges. Pests that fit this pattern include aphids, bertha armyworm and diamondback moth.



Pattern 2: Used when pests are at the edges of fields.

Scout by walking along field edges, fence lines or ditches. Some examples of when you would include more focused scouting along field edges are to estimate early-season populations of flea beetles, Colorado potato beetles and grasshoppers.

In each area examined, use of a sweep net, if possible, is a good way to determine what potential pests and beneficial insects may be present. This should be followed by examining some plants and the soil surface. More specific counts of a particular type of insect or plant damage may be necessary if they are abundant during the more general scouting.

Economic Thresholds

Monitoring methods, typical symptoms and economic thresholds or nominal thresholds for the more common crop pests are described in the field scouting section for each commodity. The density of the insects that cause damage equal to the cost of preventing the damage is called the **economic injury level**. The **economic threshold** is the density of insects at which control measures should be applied to prevent an increasing population from reaching the economic injury level. Note that factors such as moisture, temperature conditions and stage of crop growth, can increase or decrease the impact of insects on crop production. In some instances, **nominal thresholds** are presented; these decision guidelines are based on experience rather than research quantifying the impact of the insects on the crop.

Estimating Percent Defoliation

Many economic thresholds for insects are based on percent defoliation of the plants they are feeding on. The following figure may assist in determining the percent defoliation. Although the photo below is of sunflower leaves, this figure can be used to estimate % defoliation for many crops.




Photo courtesy of North Dakota State University Extension Service

Field Hazard of Insecticides to Bees

The following table indicates the maximum time required for the insecticides listed in the Guide to be degraded by weather

to a low hazard level for bees. **These times are to be used as general guidelines only.** Most of these insecticides have not been tested for bee toxicity under Western Canadian conditions and environmental conditions influence the rate at which pesticides degrade.

INSECTICIDE	HAZARD RATING ^a		RESIDUE HAZARD (DAYS) ^b
	HONEY BEE	LEAFCUTTER BEE	
Admire/ Alias/ Grapple	1	1-2	<1 - >1
Assail	1-2	-	<1
Beleaf	3	-	<1
Cygon/ Lagon	1	1	3 - 7
Coragen	3	-	-
Decis	1-2	1-2	<1 - 1
Delegate	1-2	1	-
Dibrom	1-2	1-2	<1-4.5
Dipel	3	3	none
Eco bran	3	3	N/A
Fulfill	2-3	2	<1
Lannate	1-2	1-2	<1 - 1.5
Lorsban/ Pyrinex/ Nufos/ Citadel/ Warhawk/ MPOWER Krypton	1	1	2-6
Malathion	1-2	1	2 (Honey Bee), 6 (Leafcutter Bee)
Matador/ Silencer	1	1	>1
Movento	1	-	-
Oberon	1-2	1	-
Orthene	1	1	2.5 - 3
Pounce/ Perm-UP	1	1	<1 - 5
Rimon	1-2	2	1
Ripcord/ UP-Cyde	1	1	<1 - >3
Sevin	1-2	1-2	<1 - 7
Success /Entrust	1-2	1	<1 - >1
Thiodan/ Thionex	1-2	1	<1 - 3

^a **HAZARD RATING** 1 = Very poisonous to bees; do not apply to crops or weeds in bloom unless bees are kept off for the period that residue on the crop is a hazard. 2 = Moderately poisonous to bees; avoid direct application to bees, but may be applied with minimum hazard in late evening when bees are not foraging. 3 = Not very poisonous to bees; may be applied with minimum hazard to bees.

^b Residue hazard represents the average time in days that residues poisonous to honey bees will remain on foliage (may vary with formulation and weather). Unusually low temperatures following spray application may cause residues to remain toxic longer than under warmer conditions. Morning dew can also make residues more toxic to foraging bees.

A more extensive list of hazard ratings of insecticides to bees and duration of toxicity can be found at the Western Committee on Crop Pests website at: <http://www.westernforum.org/WCCP%20Guidelines.html>

Reducing Bee Losses from Insecticides

Careless use of insecticides can kill bees and other beneficial insects such as pollinators, predatory and parasitic biological control insects. Help to reduce insecticide poisoning of bees by:

1. **Avoid applying insecticides that are toxic to bees on crops in bloom.** Any field with even a small amount of bloom, whether it is the main crop, cover crop, or weeds will probably have foraging bees visiting the flowers. If at all possible, apply insecticides before or after the crop has gone into bloom. Control all flowering weeds prior to insecticide application.
2. **Apply insecticides when bees are least active.** The highest level of bee activity occurs during the day. Apply insecticides in late evening or early morning when the bees are not foraging. As a general rule, evening applications are less hazardous to bees than morning applications.
3. **Avoid insecticide drift.** To avoid insecticides drifting into non-target locations, do not apply insecticides during windy conditions. Choose nozzles with a low drift rating. As a general rule, ground applications of insecticide are less prone to drift than aerial applications.
4. **Contact the beekeeper before spraying.** Communication and cooperation between the insecticide applicator and the beekeeper can usually prevent bee losses. Notifying the beekeeper in advance (i.e. 48 hours) of applying insecticides will allow the beekeeper to move or protect the colonies from insecticide damage.
5. **If possible, use insecticides and/or insecticide formulations which are the least hazardous to bees.** The following table "Field Hazard of Insecticides to Bees" will help in selecting the least hazardous insecticide. In general, dusts are more hazardous to bees than sprays. Wettable powders are more hazardous than emulsifiable concentrates (EC) or water-soluble formulations. Granular insecticides and spreadable bran bait insecticides are generally the least hazardous to bees.

Insecticide Poisoning in Humans

Organophosphate (OP) and carbamate insecticides (identified on the Insecticide Groups chart page 456) can pose a serious risk to unprotected persons. Poisonings can occur while mixing, loading and/or during the application of these products without the appropriate protective equipment or measures. These pesticides are readily absorbed

through the skin or the lungs, and can act as nervous system toxins. Overexposure can produce symptoms such as headache, nausea, pupil dilation and excessive sweating and salivation. Higher doses may cause breathing difficulties, muscle twitching, weakness and spasms. Very high doses have caused respiratory failure and death.

Both OP and carbamate pesticides inhibit an enzyme called cholinesterase. Measurements of cholinesterase in the blood before and during the application season can indicate harmful exposures to OPs and carbamates. **Persons who intend to mix, load and/or apply these types of pesticides repeatedly during a season, need a baseline and repeat measurements. Consult your doctor before the spraying season to arrange for these measurements.**

Degree of Risk and Hazard Rating:

(see page 11 for full description)

Resistance of Insects to Insecticides

Repeated use of the same insecticide, or insecticides with the same mode of action, against a particular insect in a given area may result in the effectiveness of the insecticide being reduced.

To delay or prevent resistance of insects to insecticides:

1. Integrate different control methods (cultural, biological, chemical) into insect control programs whenever possible.
2. Use insecticides only when the economic threshold for a pest has been surpassed and natural controls fail to limit economic damage, and
3. Rotate between insecticides with different modes of action, particularly if several applications are made in a season.
4. Keep accurate records of insecticides used for each of your fields.

Insecticides can be classified according to their similarity in chemical structure (chemical group in the table below), and by mode of action (the process by which the insecticide kills the insect). The "Group" column in the following table separates insecticides based on their mode of action. By selecting products with different modes of action for an insecticide rotation program, risk of insecticide resistance can be reduced.

Insecticide Groups Based on Modes of Action

GROUP	CHEMICAL GROUP	TRADE NAME	ACTIVE INGREDIENT	MODE OF ENTRY
1A	Carbamates	Sevin	carbaryl	contact/ ingestion (Sevin)
		Eco Bran		ingestion (Eco Bran)
		Lannate	methomyl	contact/ ingestion
		Vydate	oxamyl	contact/ ingestion
1B	Organophosphates	Malathion	malathion	contact
		Agrox B-2, Agrox CD	diazinon	ingestion
		Orthene	acephate	contact/ ingestion
		Dibrom	naled	contact/ ingestion
		Imidan	phosmet	
		Lorsban, Pyrinex, Nufos, Citadel, Warhawk, MPOWER Krypton, Pyrifos	chlorpyrifos	contact/ ingestion/ inhalation
		Lagon, Cygon 480 EC, Cygon 480-AG	dimethoate	contact/ ingestion
		Thimet	phorate	ingestion
2A	Cyclodiene organochlorines	Thiodan, Thionex	endosulfan	contact/ ingestion
3A	Pyrethroids	Decis	deltamethrin	contact/ ingestion
		Ripcord, UP-Cyde	cypermethrin	contact/ ingestion
		Matador, Silencer	lambda-cyhalothrin	contact/ ingestion
		Ambush, Pounce, Perm-UP	permethrin	contact/ ingestion
		Tempo	cyfluthrin	
4A	Neonicotinoids	Helix, Cruiser	thiamethoxam	ingestion
		Actara 240SC		ingestion
		Actara 25WG		contact/ ingestion
		Admire, Alias, Grapple, Grapple ₂ , Gaucho, Raxil ProShield, Sombrero, Stress Shield 600	imidacloprid	contact/ ingestion (flowable formulations) ingestion (seed treatments)
		Assail	acetamiprid	contact/ ingestion
5	Spinosyns	Prosper, Poncho, Titan, Clutch	clothianidin	ingestion
9B	Selective homopteran feeding blockers	Success, Entrust	spinosad	contact/ ingestion
		Delegate	spinetoram	contact/ ingestion
9C		Fulfill	pymetrozine	ingestion mainly, some contact activity
		Beleaf	flonicamid	contact/ ingestion
11	Microbial disruptors of insect midgut membranes	Dipel	<i>Bacillus thuringiensis</i> var. <i>Kurstaki</i>	ingestion
15	Benzoylureas	Rimon 10 EC	novaluron	ingestion/ contact
23	Tetronic and tetramic acid derivatives	Movento	spirotetramat	
		Oberon	spiromesifen	contact
24A		Phostoxin	aluminum phosphide	inhalation (fumigant)
28	Diamides	Coragen	chlorantraniliprole	ingestion/ contact
		Lumiderm	cyantraniliprole	ingestion

A more detailed table showing insecticides organized by mode (site) of action, and specific information on the mode (site) of action for the different groups can be found in Appendix III of PMRA Regulatory Directive DIR99-06 found at: <http://www.hc-sc.gc.ca/pmra-arla/english/pdf/dir/dir9906-e.pdf>

Field Scouting in Alfalfa

SAP OR FLUID FEEDERS

Lygus bugs/Alfalfa plant bug

Typical Damage: Field blooms poorly or not at all. Flower buds blasted, whitish, and dry; flowers dropping off before fully open. Collapsed seed.

When and How to Monitor: Look for plant bugs when monitoring alfalfa in June through mid-August. Make five 180° sweeps with a 15-inch (40 cm) insect net through alfalfa canopy at each sampling site. Record total number of plant and lygus bugs (both nymphs and adults) captured. Calculate average number per sweep.

Economic Threshold: Hay: Control not recommended. Seed alfalfa at bud and early bloom: 8 lygus bugs/sweep; 4 alfalfa plant bugs/sweep; or 5 bugs if the plant bug population is a combination of lygus bugs and alfalfa plant bug. If insecticides are used, attempt to spray before the onset of bloom. Protecting insect pollinators in seed production fields is very important.

Potato Leafhopper

Leafhoppers are most severe in new seedlings and in regrowth under hot dry weather.

When and How to Monitor: Take 20 180° sweeps from 5 areas of the field. Avoid field edges. Determine the average number of potato leafhoppers per sweep.

Economic Threshold: For 9cm stem height = 0.2 adult leafhoppers per sweep; 15 cm stem height = 0.5 adults per sweep; 25 cm stem height = 1 adult or nymph per sweep; 36 cm stem height = 2 adults or nymphs per sweep.

Pea Aphid

Typical Damage: Suck juices from plants; stunt growth; cause premature drying.

When and How to Monitor: Look for when monitoring in July through August. Take 5 sweeps at each location. Monitor fields closely during periods of slow plant growth.

Economic Threshold: 100-200 aphids/180°sweep when crop is moisture stressed, or until mid-August.

DEFOLIATORS

Alfalfa Weevil

Typical Damage: Feed on developing buds and leaves. Stunt growth.

When and How to Monitor: Start scouting fields in mid-May. Look for shot holes initially, then clipping along the edges of leaves and pinhole damage. For determining if levels are at threshold in hay crops, collect 30 stems in an M-shaped pattern, place them inside a white pail and beat them against the side to knock off larvae. Do not include younger first and second instar larvae (3 mm or less) in the counts. Determine the average height of the crop as well.

Economic Threshold: Alfalfa Hay: One of the best control strategies is to cut fields for hay early. If early cutting of the hay crop is not possible, treatment thresholds are based on the following measurements of plant height and levels of larvae: <30 cm – 1 larva/stem; <40 cm – 2 larvae/stem; 3 larvae per stem requires immediate action regardless of height of crop. On regrowth for second crop, 2 or more active larvae per crown (4 to 8 larvae/ft²) will require insecticide application.

Alfalfa Seed: 20 to 30 3rd or 4th instar larvae/sweep (90°=straight sweep) or 35 to 50% of foliage tips showing damage. In some instances it may be practical to just treat hotspots and not entire fields.

Alfalfa Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Sap or Fluid Feeders					
Lygus bugs	Rimon 10EC (seed production only) (SB)	0.338 L	14	G	>5,000
	Assail (seed production only) (N)	35-69 g	1	G	1,064
	Matador /Silencer (P)	34 ml	Do not apply within 3 days of livestock foraging.	A or G (Matador) G (Silencer)	64-110
	Decis 5EC (seed production only) (P)	80-100 ml	20	G	395
	Malathion 500 (OP)	0.80-1.21 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Dibrom (OP)	0.44-0.88	4	A or G	345

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Lygus bugs <i>continued</i>	Cygon 480 EC /Cygon 480-AG (OP) (seed and forage production)	0.17 L	2	A or G	60-450
	Lagon /Cygon 480 EC /Cygon 480-AG (OP) (seed production only)	0.44 L	28	A or G	60-450
Alfalfa plant bug	Assail (seed production only) (N)	35-69 g	1	G	1,064
	Cygon 480 EC (OP) (seed and forage production)	0.17 L	2	A or G	60-450
	Lagon /Cygon 480 EC /Cygon 480-AG (OP) (seed production only)	0.44 L	28	A or G	60-450
Potato leafhopper	Matador / Silencer (P)	34 ml	Do not apply within 3 days of livestock foraging.	A or G (Matador) G (Silencer)	64-110
	Sevin XLR Plus (C)	1.01-1.6 L	2	A or G	649
	Malathion 500 (OP)	0.80-1.21 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Dibrom (OP)	0.44-0.88 L	4	A or G	345
	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.17 L	2	A or G	60-450
Spittlebugs	Malathion 85E (OP) (adults)	0.445 – 0.544 L	7	A or G	5,500
Pea Aphid	Beleaf (seed production only) (HFB)	49 – 65 g		G	
	Matador / Silencer (P)	34 ml	Do not apply within 3 days of livestock foraging.	A or G (Matador) G (Silencer)	64-110
	Malathion 500 (OP)	0.80-1.21 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Dibrom (OP)	0.44-0.88 L	4	A or G	345
	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.17 L	2	A or G	60-450
Spider mites	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Oberon	0.202 – 0.405 L		A or G	>2,000
Defoliators					
Grasshoppers	Spreadable Bran Baits				
	Nolo Bait (M)	Minimum of 0.45 kg		A or G	
	Eco bran (C)	0.8-1.6 kg	2	G	N/A
	Sprays				
	Matador /Silencer (P)	25-34 ml (Ground) 34 ml (Aerial)	Do not apply within 3 days of livestock foraging.	A or G	64-110

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Grasshoppers continued	Sevin XLR Plus (C)	0.50-1.01 L	2	A or G	649
	Malathion 500 (OP)	0.80-1.21 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Lagon / Cygon 480 EC / Cygon 480-AG (OP)	0.22 L (nymphs) 0.34-0.36 L (adults)	2-7 (Lagon, Cygon 480 EC) 28 (Cygon 480-AG)	A or G	60-450
Alfalfa weevil	If alfalfa has reached the bud or early bloom stage, immediate cutting will kill many alfalfa weevil larvae				
	Coragen (D) (suppression only)	152-202 ml	0	G	>5,000
	Matador / Silencer (P)	34 ml	Do not apply within 3 days of livestock foraging.	A or G (Matador) G (Silencer)	64-110
	Decis 5EC (seed crops only) (P)	80 – 100 ml	20	G	395
	Malathion 500 (OP)	0.80-1.21 L	7	A or G	4302
	Malathion 85E (larvae only) (OP)	0.445-0.544 L	7	A or G	5,500
	Imidan (OP)	0.65 kg	7	G	285
	Lagon / Cygon 480 EC / Cygon 480-AG (OP) (reduction only)	0.17 L	2	A or G	60-450
Blister beetle	Sevin XLR Plus (C)	1.01-1.6 L	2	A or G	649
True Armyworm	Sevin XLR Plus (C)	1.01-2.12 L	2	A or G	649
Beet webworm	Sevin XLR Plus (C)	1.01-2.12 L	2	A or G	649
Alfalfa looper	Dibrom (OP)	0.44-0.88 L	4	A or G	345
Alfalfa Caterpillar	Sevin XLR Plus (C)	1.01-2.12L	2	A or G	649
Leafminers					
Alfalfa blotch leafminer	Malathion 85E (OP)	0.544 L	7	A or G	5,500
	Imidan (OP)	0.65 kg	7	G	285
	Lagon / Cygon 480 EC / Cygon 480-AG (OP)	0.22 L	2	A or G	60-450

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: M=microbials, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀

Barley-See Small Grain Cereals

Scouting for insects in Beans (Dry Edible Beans)

BELOWGROUND FEEDERS AND CUTWORMS

Seedcorn Maggot

Typical Damage: Seedcorn maggot attacks bean seed, preventing sprouting or weakening seedlings. The yellowish white maggot is found burrowing in the seeds or emerging stem. Seedcorn maggots are usually most severe in wet, cold seasons and on high organic matter soils.

Cutworms

When and How to Monitor: To find cutworms, dig in the soil to a depth of 2.5 to 5 cm at the base of recently damaged plants.

Nominal Threshold: Treatment is warranted when one cutworm or more is found per metre of row and the larvae are still small (less than 2 cm long).

SAP FEEDERS

Leafhoppers

Typical Damage: Foliage becomes dwarfed, crinkled, and

curled. Small triangular brown areas appear at the tips of leaves, gradually spreading around the entire leaf margin.

When and How to Monitor: Leafhopper adults are quick and can be observed by running your hand over the top of the plants as you approach them and observing adults that fly off the plants. On the same plants, turn over each leaf to determine the number of nymphs per trifoliolate.

Economic Threshold: Unifoliolate stage – 0.25 leafhoppers per trifoliolate; second trifoliolate stage – 0.5 leafhoppers per trifoliolate; fourth trifoliolate stage – 1.0 leafhopper per trifoliolate; first bloom-2.0 leafhoppers per trifoliolate.

DEFOLIATORS

Grasshoppers

Economic Threshold: Substantial yield loss does not occur until up to 35% defoliation occurs before bloom and 15% after bloom.

Beans (Dry Edible) Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Beans (N)	Seed Treatment			
	Stress Shield 600 (N)	A co-pack of Raxil Pro and StressShield 600			
Seedcorn Maggot	Sow seeds as shallow as possible in a warm, well-prepared seedbed. If manure is used, apply and plow it under the previous fall.				
	Cruiser Maxx Beans (N)	A seed treatment combining Cruiser 5FS and Apron Maxx RTA			
	Agrox CD, Agrox B-2, DCT (OP)	See Seed Treatments section of Guide to Crop Protection. For control of Seedcorn maggots only		Seed Treatment	N/A
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Sap or Fluid Feeders					
Lygus Bugs	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
	Sevin XLR Plus (C)	2.12-2.59 L	5	A or G	649
	Cygon 480-AG (OP)	0.28-0.40 L	7	A or G	60-450

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Potato Leafhoppers	Cruiser Maxx Beans (N)	A seed treatment combining Cruiser 5FS and Apron Maxx RTA			
	Stress Shield 600 (N)	Seed Treatment			
	Gaucho 480 FL (N)	Apply at a rate of 130 ml per 100 kg seed			
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
	Sevin XLR Plus (C)	1.01 L	5	A or G	649
	Malathion 500 (OP)	0.56-1.21 L	1	A or G	4302
	Malathion 85E (OP)	0.297-0.544 L	3	G	5,500
	Cygon 480-AG (OP)	0.28-0.40 L	7	A or G	60-450
Aphids	Movento	75 - 111 ml	7	G	>2,000
	Matador (P)	34 - 94 ml	14	A or G	64-110
	Malathion 500 (OP)	0.56-1.21 L	1	A or G	4302
	Malathion 85E (OP)	0.297-0.544 L	3	G	5,500
	Dibrom (OP)	0.445-0.890 L	4	A or G	345
	Cygon 480-AG (OP)	0.28-0.40 L	7	A or G	60-450
Defoliators and Borers					
Grasshoppers	Spreadable Bran Baits				
	Eco bran (C)	0.8-1.6 kg	5	G	N/A
European Corn Borer	Coragen (D)	101-152 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Variegated Cutworm	Sevin XLR Plus (C)	30-35 mL/100 m of row	5	A or G	649
Alfalfa looper	Dibrom (OP)	0.445-0.890 L	4	A or G	345

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Scouting for insects in Canaryseed

APHIDS

When and How to Monitor: Start checking for aphids when monitoring during the early heading stage of canaryseed. The head should be bent and closely inspected for aphids hiding along the small stem inside the canaryseed head.

Also check the stems, underside of leaves, and in the canaryseed boot.

Nominal Threshold: 10 to 20 aphids on 50% of the stems prior to the soft dough stage.

Canaryseed Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Sap Feeders					
Aphids	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.20 L	21	A or G	60,450
	Malathion 85E (OP)	0.277 L	14	A or G	5,500

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Canola

Scouting Calendar

Early-season: Flea beetles, cutworms, red turnip beetle, diamondback moth

Mid-season: Diamondback moth, lygus bug, cabbage seed-pod weevil, grasshoppers

Late season: Bertha armyworm, diamondback moth, Lygus bugs, grasshoppers, alfalfa looper

Cutworms

Typical Damage: Notched, wilted, dead, or cut-off plants (weed or crop seedlings). Plants missing from rows, bare patches appearing in field.

When and How to Monitor: Look for cutworms, and evidence of cutworm feeding, when monitoring canola in late May to mid-July. Often cutworms will be close to the cut or shriveled plants that they have just damaged. Cutworms will sometimes be most abundant in patches or a specific area of a field.

Nominal Threshold: 25-30% stand reduction. Sometimes it is most economical to just treat infested patches, and not whole fields.

SAP OR FLUID FEEDERS:

Lygus Bugs

Typical Damage: Attacked buds appear shrunken and bleached white. Damaged seeds appear dark brown and shriveled.

When and How to Monitor: Monitor, using an insect sweep net, from the end of flowering until seeds within the pod have become firm. Make 10 180° sweeps with a 37 cm diameter insect net at each of at least 5 sampling site. If while doing these samples populations appear to be of concern, take additional samples; a minimum of 15 samples is needed to accurately determine whether controls are economical. Sample canola for lygus bugs on a sunny day when the temperature is above 20°C and the crop canopy is dry.

Economic Threshold: 10-18 lygus bugs/10 sweeps from the end of flowering to early pod development in the upper canopy and 15-25 lygus bugs/10 sweeps in early pod ripening stage. Controls are not recommended when seeds are ripening (yellow or brown). When precipitation is greater than 100 mm from the onset of bud formation to the end of flowering, the crop may partially compensate for plant bug damage.

A table of specific economic thresholds for various expected values of canola seed and costs of control for lygus bugs in canola can be found at: <http://www.gov.mb.ca/agriculture/crops/insects/fad12s00.html>

Aphids

Economic Threshold: Control aphids in canola if densities exceed 25 aphids/10 cm shoot tip after flowering.

DEFOLIATORS:

Flea beetles

Typical Damage: Shot-holes in leaves to complete destruction of seedling plants in late May through June. Holes chewed in pods in August (occasional).

When and How to Monitor: Look for when monitoring in May through June when crop is in seedling stage. Examine 10 plants at random at each stop. Estimate overall percentage leaf loss.

Economic Threshold: When 25 percent of leaf surface is destroyed and flea beetles are present. If damage is only along the field margins and beetles are still congregated there, then control measures should be applied to the damaged areas only.

Diamondback moth

Typical Damage: Flowers clipped or chewed, outer layers of stem and pods chewed, holes chewed in pods.

When and How to Monitor: Look for when monitoring in late May through early September. Observing for adults and larvae while taking sweep net samples can determine the presence and relative abundance of diamondback moth in the field. If levels appear to be of concern, shake plants within a 50 cm x 50 cm area and count larvae on the ground or surface (such as a sweep net) that plants were shaken over. Another alternative is to clip or pull the plants and knock over a light colored surface (such as a sweep net, jacket, hood of a car, etc.). Multiply by 4 to get the number of larvae per square metre. Do this in at least 5 areas of the field.

Nominal Threshold: 100 to 150 larvae/m² in immature to flowering plants. 200 to 300 larvae/m² in plants with flowers and pods. Note that these threshold numbers are based on stands averaging 150-200 plants/m². In areas where stands are thinner, the economic threshold should be lowered accordingly. A nominal threshold of 25-33% defoliation with larvae still present can be applied for canola at seedling stage.

Bertha Armyworm

Typical Damage: Outer layers of stems and pods chewed resulting in whitish appearance, holes chewed in pods.

When and How to Monitor: Look for larvae when monitoring fields in late July through early August. At each stop, shake plants in a 1/4 m² (50 cm x 50 cm) area and carefully check soil surface for dislodged larvae. During heat of the day larvae will often be found under leaves on soil surface.

Economic Threshold: A loss of 0.058 Bu/acre for each larva/m² can be expected. Multiplying 0.058 X average number of larvae per m² X expected seed value (dollars/acre) will determine the economic loss (in dollars/acre) due to the larvae. Only if control costs (insecticide plus application costs) can be applied for less than this economic loss will insecticide applications be economical. Yield loss may be greater for canola under moisture stress.

At an expected seed value of \$6.00/bushel, the economic threshold will be between about 20 and 34 larvae/m², depending on control costs. At an expected seed value of

\$8/bushel, the economic threshold will be between about 15 and 26 larvae/m², depending on control costs. Tables showing specific economic thresholds at various expected

seed values and control costs can be found at: <http://www.gov.mb.ca/agriculture/crops/insects/fad03s01.html>

Canola Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Cabbage root maggot	<ul style="list-style-type: none">• Increased seeding rates and increased row spacing (to about 25 to 30 cm) can reduce damage to the roots by root maggots.• Cultivating prior to seeding reduces adult emergence from overwintered pupae. Root maggot infestations are greater under zero-till systems than under conventional tillage, but yields under zero tillage usually still exceed those with conventional tillage.				
Cutworms (Army, darksided, redbacked, pale western)	Seed Treatments				
	Prosper EverGol + Lumiderm	Seed Treatment			
	Helix Vibrance + Lumiderm	Seed Treatment			
	Foliar Sprays				
	Coragen (D)	101 ml	1	A or G	>5,000
	Pounce / Perm-UP (P) Ambush (P)	73 – 158 ml 57-121 ml	Treat prior to 6-leaf stage	G	1276
	Chlorpyrifos (OP)	0.354-0.486 L	21	A or G	205-418
Sap and Fluid Feeders					
Lygus Bugs	Decis 5EC (P)	60 ml	7	A or G	395
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Chlorpyrifos (OP)	0.202 - 0.405 L	21	A or G	205-418
Turnip aphid	Lagon /Cygon 480-AG (OP)	0.34-0.36 L	21	A or G	60-450
Aster leafhopper	Lagon /Cygon 480-AG (OP)	0.34-0.36 L	21	A or G	60-450
Swede midge	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
Defoliators					
Crucifer Flea beetle and/or striped flea beetle	Seed Treatments				
	Helix Vibrance (N)	A seed treatment containing Helix Xtra and Vibrance 500FS			
	Helix Vibrance + Lumiderm	Seed Treatment			
	Helix Xtra (N)	1500 ml/100 kg of seed	Seed Treatment		>5,000
	Acceleron	A seed treatment containing Helix Xtra and Acceleron			N/A
	Prosper EverGol (N)	Seed Treatment			N/A
	Prosper EverGol + Lumiderm	Seed Treatment			
	Gaucho Canola System (N)	0.833 L/ 100 kg of seed	Seed Treatment		N/A
	Gaucho Platinum (N)	1.667L/100 kg of seed	Seed Treatment		N/A
	Sombrero (N)	0.67 – 1.33 L / 100 kg seed	Seed Treatment		N/A

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Crucifer Flea beetle and/or striped flea beetle	Foliar Sprays				
	Decis 5EC (P)	40 -60 ml	7	A or G	395
	Ripcord (P)	20 ml	30	G	242-542
	UP-Cyde (P)	56.6 ml	30	A or G	355
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Pounce (P) Ambush (P)	36-73 ml 28-57 ml		A or G	1276
	Sevin XLR Plus (C)	0.2 L	Seedling application only	A or G	649
	Malathion 500 (OP)	0.44 L	7	A or G	4302
	Malathion 85E (OP)	0.217-0.346 L	7	A or G	5,500
Cabbage seedpod weevil	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Decis 5EC (for control of adults only) (P)	80 ml	7	A or G	395
Diamondback moth	Coragen (D)	51 ml	1	A or G	>5,000
	Decis 5EC (P)	40 - 60 ml	7	A or G	395
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Malathion 500 (OP)	0.22-0.34 L	7	A or G	4302
	Malathion 85E (OP)	0.109-0.168 L	7	A or G	5,500
	Chlorpyrifos (OP)	0.405 -0.607L	21	A or G	205-418
Bertha Armyworm	Seeding as early as possible and choosing early maturing varieties of canola may help minimize damage in years when outbreaks are forecasted.				
	Coragen (D)	51 - 152 ml	1	A or G	>5,000
	Decis 5EC (P)	40 - 60 ml	7	A or G	395
	Ripcord (P)	28 ml (ground) 36 ml (air)	30	A or G	242-542
	UP-Cyde (P)	81 - 113 ml	30	A or G	355
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Lannate (C)	87.4 - 206.4 g	8	A or G	30-34
	Chlorpyrifos (OP)	0.304-0.405 L	21	A or G	205-418
Alfalfa looper	Lannate (C)	87 - 206 g	8	A or G	30-34
	Chlorpyrifos (OP)	0.304-0.405 L	21	A or G	205-418
Cabbage looper	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
Beet webworm	Decis 5EC (P)	40 - 60 ml	7	A or G	395
	Lannate (C)	87.4 - 206.4 g	8	A or G	30-34

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Clover cutworm	Decis 5EC (P)	40 – 60 ml	7	A or G	395
	Lannate (C)	87.4 – 206.4 g	8	A or G	30-34
True armyworm	Chlorpyrifos (OP)	0.304-0.405 L	21	A or G	205-418
Imported Cabbageworm	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
Variegated cutworm	Chlorpyrifos (OP)	0.354-0.486	21	A or G	205-418
Grasshoppers	Spreadable Bran Baits				
	Eco bran (C)	0.8-1.6 kg	Treat only seedlings	G	N/A
	Sprays				
	Decis 5EC (P)	40 – 60 ml (Ground); 60 ml (Aerial)	7	A or G	395
	Matador /Silencer (young grasshoppers only) (P)	25 – 34 ml (Ground) 34 ml (Aerial)	7	A or G	64-110
	Ripcord (young grasshoppers only) (P)	20 – 28 ml	30	G	242-542
	Malathion 500 (OP)	0.45-0.69 L	7	A or G	4302
	Malathion 85E (OP)	0.217-0.346 L	7	A or G	5,500
	Chlorpyrifos (OP)	0.235-0.354 L	21	A or G	205-418
	Lagon / Cygon 480-AG / Cygon 480 EC (OP)	0.34-0.36 L	21	A or G	60-450
Slugs	Sluggo Professional	10-20 kg		G	>5,000

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Chickpeas Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Pulses (N)	A seed treatment containing Cruiser-5FS and Apron Maxx RFC			
	Stress Shield 600 (N)	Seed Treatment			
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Sap Feeders					
Pea Aphids	Movento	75 - 111 ml	7	G	>2,000
	Matador /Silencer (P)	34 - 94 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Potato Leafhopper	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Defoliators					
Grasshoppers	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, P=pyrethroids, N=neonicotinoids.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Corn (Field Corn)

Cutworms

Typical Damage: Notched, wilted, dead, or cut-off plants (weed or crop seedlings). Plants missing from rows, bare patches appearing in field.

When and How to Monitor: Look for cutworms, and evidence of cutworm feeding, when monitoring corn in late – May to mid-July. Often cutworms will be close to the cut or shriveled plants that they have just damaged. Cutworms will sometimes be most abundant in patches or a specific area of a field. At each stop, examine 100 plants in a row. Calculate percentage of plants cut off or showing leaf feeding.

Economic Threshold: When 3-6% of plants are cut and small larvae less than 1 inch present. Sometimes it is most economical to just treat infested patches, and not whole fields.

European corn borer

Typical Damage: Shot-holes in leaves. Holes in stalk, tassels and ears. Damage may cause stalk breakage prior to harvest or cobs to fall to the ground. Nutrient flow in the plant may be restricted, resulting in smaller cobs.

When and How to Monitor: Begin looking for European corn borer when field scouting in early July. At 5 locations, examine 10 plants for young larvae and egg masses. Calculate percentage of plants infested. Scout every 5 to 7 days until the end of July or larvae start to tunnel into the stalks.

Economic Threshold: The level of European corn borer where control becomes economical depends on the value of the crop, and cost of control. Information on determining specific economic thresholds for European corn borer in corn can be found at <http://www.gov.mb.ca/agriculture/crops/insects/fad46s00.html>, or from your local agriculture office. These thresholds are based on a 5% yield loss per corn borer per plant on average. If the majority of larvae have bored into the stalk, do not apply insecticide, as they are ineffective once the larvae have entered the stalk.

Corn (Field Corn) Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Cutworms	Matador /Silencer (P)	34 ml	14	A or G	64-110
	Ripcord (black, white, darksided, redbacked, army, pale western) (P)	70 ml	21	G	242-542
	Pounce / Perm-UP (P) Ambush (P)	73 - 158 ml 57-121 ml	Treat prior to 6 leaf stage	G	1030
	Chlorpyrifos (darksided, black, redbacked) (OP)	0.971 L (Pre-plant treatment), 0.486-0.971 L (seedling treatment)	70	G	205-418
Wireworms	Cruiser Extreme 250 (N)	A seed treatment combining Cruiser 5FS, Maxim XL, Apron XL, and Dynasty 100FS			
	Poncho 250 (N)	0.25 mg of Poncho 600 per kernel	Seed Treatment	Seed Treatment	500-1,000
	Sombrero (N)	0.16 mg per kernel	Seed Treatment	Seed Treatment	N/A
Seedcorn maggot	Cruiser Extreme 250 (N)	A seed treatment combining Cruiser 5FS, Maxim XL, Apron XL, and Dynasty 100FS			
	Poncho 250 (N)	0.25 mg of Poncho 600 per kernel	Seed Treatment	Seed Treatment	500-1,000
	Agrox CD (OP)	50 g/25 kg of seed	Seed Treatment	Seed Treatment	N/A
	Agrox B-2 (OP)	See label	Seed Treatment	Seed Treatment	N/A
Sap Feeders					
Spider mites	Oberon	162-243 ml	Green forage - 5 Grain or stover - 30	A or G	>2,000
Defoliators and Borers					
Grasshoppers	Spreadable Bran Baits				
	Eco bran (C)	0.8-1.6 kg	1	G	N/A
	Sprays				
	Sevin XLR Plus (C)	0.50-1.01 L	1	A or G	649

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
European Corn Borer	Stalk Management: Primary tillage such as chisel plowing or moldboard plowing in the fall can reduce overwintering populations. Mowing corn stalks after harvest can reduce overwintering populations up to 85%.				
	Resistant Cultivars: Some cultivars of Bt corn are resistant to feeding by European corn borer. If planting cultivars of Bt corn, a refuge of non-Bt cultivars is required to be planted to reduce the odds of European corn borer developing resistance to Bt corn. Growers of Bt corn are also required to monitor their crop for the presence of European corn borer and any feeding damage. A table of registered Bt corn products in Canada (as of September 2011) is available at: http://www.cornpest.ca/index.cfm/bt-corn/registered-bt-hybrids/				
	Dipel 2X DF (M)	0.23-0.45 kg	0	G	>4,000
	Coragen (D)	101 – 152 ml	14	G	>5,000
	Matador/ Silencer (P)	34-76 ml	14 (silage) 21 (field corn)	A or G	64-110
	Decis 5EC (P)	0.1-0.12 L	N/A	G	395
	Ripcord (P)	70 ml	5	A or G	242-542
	UP-Cyde (P)	113 ml	5	A or G	355
	Sevin XLR Plus (C)	1.01-1.6 L	1	A or G	649
	Malathion 85E (OP)	0.445-0.544 L	5	A or G	5,500
Corn Earworm	Some cultivars of Bt corn are resistant to feeding by corn earworm.				
	Coragen (D)	101 – 152 ml	14	G	>5,000
	Matador/ Silencer (P)	34-76 ml	14 (silage) 21 (field corn)	A or G	64-110
	Ripcord (P)	70 ml	5	A or G, see product label	242-542
	Sevin XLR Plus (C)	1.01-1.6 L	1	A or G	649
	Malathion 85E (OP)	0.445-0.544 L	5	A or G	5,500
Armyworm	Coragen (D)	101 – 152 ml	14	G	>5,000
	Matador /Silencer (P)	34 ml	14 (silage) 21 (field corn)	A or G	64-110
Fall armyworm	Some cultivars of Bt corn are resistant to feeding by fall armyworm				
	Coragen (D)	101 – 152 ml	14	G	>5,000
	Matador/Silencer (P)	34 ml	14	A or G	64-110
	Pounce (P)	73 ml	1	G	1030
	Sevin XLR Plus (C)	1.01-1.6 L	1	A or G	649

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: M=microbials, D=diamides, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Fababean Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground Feeders					
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
Wireworms	Cruiser Maxx Pulses (N)	A seed treatment containing Cruiser 5FS and Apron Maxx RFC			
	Stress Shield 600 (N)	Seed Treatment			
Sap and Fluid Feeders					
Lygus Bugs	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Potato Leafhopper	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Pea Aphid	Matador /Silencer (P)	34 - 94 ml	14 (Matador) 21 (Silencer)	A or G	64-110

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, N=neonicotinoids, P=pyrethroids.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Flax

Cutworms

Typical Damage: Notched, wilted, dead, or cut-off plants (weed or crop seedlings). Plants missing from rows, bare patches appearing in field.

When and How to Monitor: Look for cutworms, and evidence of cutworm feeding, when monitoring in late May to mid-July. Often cutworms will be close to the cut or shriveled plants that they have just damaged. Cutworms will sometimes be most abundant in patches or a specific area of a field. In areas of the field where cutworm damage is noticeable, check around damaged plants in a 0.25 m² (50cm x 50cm) area. Use trowel or shovel to carefully search through top 5 cm of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m². Repeat in several locations to get an accurate assessment of what cutworm levels are.

Economic Threshold: 4-5 larvae/m². Sometimes it is most economical to just treat infested patches, and not whole fields.

Aphids

Typical Damage: Extract plant fluids from the stems, leaves and developing bolls. Can cause fewer seeds to be produced.

When and How to Monitor: The easiest way to detect aphids in flax is to sample the upper portions of the plant with an insect sweep net when the crop is in full bloom, or

tap plants over a white tray or bucket. If aphids are found, fields need to be more closely inspected by randomly sampling plants. To inspect plants, lightly tap the plants on a white surface, such as a tray or the canvas of a sweep net, to dislodge the insects. Plants can be severed at the base prior to tapping if desired. Inspect a minimum of 25 plants at full bloom and 20 plants at early green boll randomly in the field to provide an accurate estimate of aphid density. Record total number of aphids and calculate average per plant.

If control is not warranted at full bloom, aphid densities should be assessed again at the green boll stage.

Economic Threshold: Varies with crop value and control costs, but generally about 3 aphids per main stem at full bloom or 8 aphids per main stem at the green boll stage.

The yield loss of flax is 0.3346 bu/acre per aphid per plant for crops sampled at full bloom and 0.1275 bu/acre per aphid per plant for crops sampled at the green boll stage.

The potato aphid is highly susceptible to attack by fungi (especially in years of high rainfall and humidity in late June and July). Aphid populations sampled at full bloom that have many diseased insects should be sampled again at the early green boll stage to determine the effect of the disease on aphid densities.

Beet webworm

Nominal Threshold: >10 larvae/m²

Flax Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	No insecticides registered for the control of wireworms in flax				
Cutworms (Army, darksided, redbacked, pale western)	Coragen (D)	101 ml	1	A or G	>5,000
	Decis 5EC (P)	80 ml	40	A or G	395
	Pounce / Perm-UP (P) Ambush (P)	73 - 158 ml 57-121 ml	Treat prior to 6 leaf stage	G	1030
	Chlorpyrifos (OP)	0.354-0.486 L	21	A or G	205-418
Sap Feeders					
Potato Aphid	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.18 L	21	A or G	60-450
Defoliators					
Grasshoppers	Decis 5EC (P)	40 - 60 ml	40	A or G	395
	Matador /Silencer (young grasshoppers only) (P)	25 - 34 ml (Ground) 34 ml (Aerial)	7	A or G	64-110
	Malathion 500 (OP)	0.44-0.68 L	7	A or G	4302
	Malathion 85E (OP)	0.217-0.346 L	7	A or G	5,500
Bertha Armyworm	Coragen (D)	51 - 152 ml	1	A or G	>5,000
	Lannate (C)	89 - 109 g	8	A or G	30-34
	Chlorpyrifos (OP)	0.304-0.405 L	21	A or G	205-418
True Armyworm	Chlorpyrifos (OP)	0.354-0.486 L	21	A or G	205-418
Clover Cutworm	Decis 5EC (P)	40 - 60 ml	40	A or G	395
Variegated cutworm	Chlorpyrifos (OP)	0.354-0.486 L	21	A or G	205-418
Beet Webworm	Decis 5EC (P)	40 - 60 ml	40	A or G	395

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Forage Grasses (Timothy, etc.) Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Defoliators					
Grasshoppers	Spreadable Bran Baits				
	Eco bran (C)	0.8-1.6 kg	1-2	G	N/A
	Sprays				
	Matador /Silencer (P) (on timothy)	25 - 34 ml	14	G	64-110
	Sevin XLR Plus (C)	0.49-0.93 L (for nymphs or sparse vegetation) 0.93-1.42 L (for adults or application to dense vegetation)	1	A or G	649
	Malathion 500 (OP)	0.69 L	7	A or G	4302
	Lagon /Cygon 480-AG (OP)	0.17 - 0.22 L (nymphs) 0.34 - 0.40 L (adults)	2-28	A or G	60-450
European skipper (on timothy)	Dipel 2X DF (M)	57 - 111 g	N/A	A or G	>4,000
Armyworm	Coragen (D) (seed production only)	101 - 152 ml	0	G	>5,000

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: M=microbials, D=diamides, P= pyrethroids, C=carbamates, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Lentils

Grasshoppers

When and How to Monitor: Look for when monitoring fields from the early bud stage through pod development.

Economic Threshold: 2 grasshoppers/m² during the flowering and podding stages, especially if two-striped grasshopper is the dominant species.

Lygus Bugs

When and How to Monitor: Look for adult lygus bugs when monitoring lentils during blooming and podding

by using a sweep net, making 25 180° sweeps in at least 5 randomly selected places in a field.

Economic Threshold: Insecticide treatment is recommended when 10 or more adults are collected per 25 sweeps.

Pea aphid

Economic Threshold: 30-40 aphids per 180° sweep of a 38 cm (15 inch) diameter insect net, and few natural enemies are present, and when aphid numbers do not decline over a 2-day period.

Lentil Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Pulses (N)	A seed treatment containing Cruiser 5FS and Apron Maxx RFC			
	Stress Shield 600 (N)	Seed Treatment			
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
	Decis 5EC (P)	80 ml	30	A or G	395
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
	Pounce / Perm-UP (P)	73 - 158 ml	Treat prior to 6-leaf stage	G	1030
	Ambush (P)	57-121 ml			
	Chlorpyrifos (for pale western cutworm only) (OP)	0.354-0.486 L	21-60	A or G	205-418
Sap and Fluid Feeders					
Lygus Bugs	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Potato Leafhopper	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Pea Aphid	Movento	75 - 111 ml	7	G	>2,000
	Matador /Silencer (P)	34 - 94 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Defoliators					
Grasshoppers	Decis 5EC (P)	40-60 ml (ground) 60 ml (air)	30	A or G	395
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
	Malathion 500 (OP)	0.68 L	30	A or G	4302
	Malathion 85E (OP)	0.336 L	14	A or G	5,500
	Chlorpyrifos (OP)	0.235-0.486 L	21-60	A or G	205-418

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, N=neonicotinoids, P= pyrethroids, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Mustard Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Root Maggot	No insecticides registered				
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Sap Feeders					
Lygus Bugs	Decis 5EC (P)	60 ml	7	A or G	395
Defoliators					
Flea beetles	Seed Treatments				
	Helix (N)	1500 ml/100 kg of seed	N/A	Seed Treatment	>5,000
	Helix Xtra (N)	1500 ml/100 kg of seed	N/A	Seed Treatment	>5,000
	Prosper (N)		Seed Treatment	Seed Treatment	N/A
	Gaucho Canola System (N)	0.833 L/100 kg of seed	N/A	Seed Treatment	N/A
	Gaucho Platinum (N)	1.667 L/100 kg of seed	N/A	Seed Treatment	N/A
	Sombrero (N)	0.67 - 1.33 L / 100 kg seed	Seed Treatment	Seed Treatment	N/A
	Sprays				
	Decis 5EC (P)	40 - 60 ml	7	A or G	395
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	UP-Cyde (P)	57 ml	30	G	355
	Malathion 85E (OP)	0.217-0.346 L	7	A or G	5,500
Cabbage seedpod weevil	Note: Yellow mustard (<i>Sinapis alba</i>) is resistant to cabbage seedpod weevil; oriental and brown mustards (<i>Brassica juncea</i>) are susceptible to feeding by cabbage seedpod weevil.				
	Matador /Silencer (adults) (P)	34 ml	7	A or G	64-110
	Decis 5 EC (for control of adults only) (P)	80 ml	7	A or G	395
Diamondback Moth	Coragen (D)	51 ml	1	A or G	>5,000
	Decis 5 EC (P)	40 - 60 ml	7	A or G	395
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Malathion 85E (OP)	0.109-0.168 L	7	A or G	5,500
Bertha Armyworm	Coragen (D)	51 - 152 ml	1	A or G	>5,000
	Decis 5EC (P)	40 - 60 ml	7	A or G	395
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	UP-Cyde (P)	81 - 113 ml	30	G	355
Clover Cutworm	Decis 5EC (P)	40 - 60 ml	7	A or G	395
Imported cabbageworm	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
Cabbage looper	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
Beet webworm	Decis 5EC (P)	40 - 60 ml	7	A or G	395
Grasshopper	Decis 5EC (P)	40 - 60 ml	7	A or G	395
	Matador /Silencer (young grasshoppers only) (P)	25- 34 ml(Ground) 34 ml (Aerial)	7	A or G	64-110
	Malathion 85E (OP)	0.217-0.346 L	7	A or G	5,500

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, N= neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Oats - See small grain cereals

Grasshopper Management on Pastures, Rangelands, Hay, Headlands, and Roadsides

Note: Insects for biological control of weeds such as leafy spurge may be introduced and established in some areas of Manitoba and Saskatchewan. If grasshopper numbers become high, consider using control strategies and insecticides that will minimize harm to these biological control agents.

Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Reduced Agent and Area Treatments (RAATs): Grasshoppers on rangelands may be managed by applying certain insecticides in treated swaths, which alternate with untreated swaths. This can reduce the cost of control and amount of insecticide used by more than 50 percent, while resulting in effective control. For more information on managing grasshoppers by this method see: http://www.grasshoppercontrol.com/App/PDF/6.pdf				
Spreadable Bran Baits				
Nolo Bait (Pastures, Rangelands)	Minimum of 0.45 kg		A or G	
Eco bran (pastures, rangelands, field borders, headlands, right-of-way, roadsides, wastelands) (C)	0.8-1.6 kg	0-2 (see label)	G	N/A
Sprays				
Decis 5EC (Rangeland, pastures, roadside) (P)	40 – 60 ml	N/A	A or G (Rangeland, pastures) G (Roadsides)	395
Ripcord (Roadsides, headlands, and summerfallow) (young grasshoppers only) (P)	20 – 28 ml	Treated areas must not be grazed or cut or hay.	G	242-542
Matador (Unimproved pasture, summerfallow) (young grasshoppers only) (P)	25 – 34 ml (Ground) 34 ml (Aerial)	3	A or G	64-110
Silencer (Unimproved pasture) (young grasshoppers only) (P)	25 – 34 ml (Ground) 34 ml (Aerial)	3	A or G	64-110
Sevin XLR Plus (ditchbanks, field borders, headlands, pastures, rangelands, right-of-way, wastelands) (C)	0.48-1.41 L	0 (ditchbanks, field borders, headlands, right-of-way, wasteland); 1 (pastures, rangeland)	A or G	649
Malathion 500 (Hay only) (OP)	0.69 L	7	A or G	4302
Malathion 85E (pastures, rangelands) (OP)	0.336 L	Do not apply to fields occupied by dairy animals, but may be grazed or harvested on the day of application.	G	5,500

Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Dibrom (Rangeland, pastures, dairy and horse paddocks) (OP)	0.22-0.34 L (young grasshoppers) 0.28-0.4 L (adult grasshoppers)	4	A or G	345
Lagon /Cygon 480 EC /Cygon 480-AC (pasture, wasteland) (OP)	0.22 L (nymphs) 0.34-0.41 L (Adults)	2 days - 0.22L rate 7-28 days - 0.34-0.41L rates (see labels)	A or G	60-450
Lagon (Hay) (OP)	0.17-0.22 L	2	A or G	60-450
Chlorpyrifos (OP)	Ungrazed and unoccupied areas such as roadsides, right of way, and fence lines adjacent to barley, wheat, oats, or canola, and lentils.		A or G	205- 418

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: P=pyrethroids, C=carbamates, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Peas (Field Peas)

Cutworms

Nominal Threshold: 2 to 3 cutworms per square metre.

SAP FEEDERS

Aphids

When and How to Monitor: Look for when monitoring field peas at the beginning of flowering. Take 180° sweeps or check 10 8-inch (20 cm) plant tips at each stop. Record

total number of aphids and calculate average per sweep or plant tip.

Economic Threshold: If, at the beginning of flowering, there are 9 to 12 aphids per sweep or 2-3 aphids per 8-inch (20 cm) plant tip, an insecticide application when 50 percent of plants have produced some young pods will be cost-effective.

Peas (Field Peas) Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Pulses (N)	A seed treatment containing Cruiser 5FS and Apron Maxx RFC			
	Stress Shield 600 (N)	Seed Treatment			
Seedcorn Maggot	Agrox CD (OP)		Seed Treatment	Seed Treatment	N/A
	Agrox B-2 (OP)		Seed Treatment	Seed Treatment	N/A
Cutworms (Army, black, darksided, pale western, red-backed, white)	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
	Pounce / Perm-UP (P)	73 – 158 ml	Treat prior to 6 leaf stage	G	1030
	Ambush (P)	57-121 ml			
Sap and Fluid Feeders					
Leafhoppers	Malathion 85E (OP)	0.445 L	3	A or G	5,500
Pea Aphid	Movento	75 – 111 ml	7	G	>2,000
	Matador /Silencer (P)	34 - 94 ml	14 (Matador) 21 (Silencer)	A or G	64-110
	Lannate (C)	0.206 kg	1	G	30-34
	Malathion 85E (OP)	0.445 L	3	A or G	5,500
	Lagon/Cygon 480 EC (OP)	0.11-0.17 L	3-21 (see labels)	A or G	60-450
Defoliator					
Grasshoppers	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110
Alfalfa Looper	Sevin XLR Plus (C)	1.90 L	3	G	649
	Lannate (C)	0.206 kg	1	G	30-34
Pea leaf weevil	Cruiser Maxx Pulses (N)	A seed treatment combining Cruiser 5FS and Apron Maxx RTA			
	Stress Shield 600 (N)	Seed Treatment			
	Matador /Silencer (P)	34 ml	14 (Matador) 21 (Silencer)	A or G	64-110

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Scouting and Thresholds for Insects in Potatoes

Aphids

Typical damage: Several species of aphids are sap feeders on potato leaves. At very high numbers this sap feeding may cause plants to wilt in small localized areas of the field as "aphid holes". The greatest injury is due to transmission of viruses. Identification and control of aphids is critical in potato seed production to prevent virus spread. In commercial production tuber quality may be reduced by net necrosis of tubers.

When and How to Monitor: Aphid identification and scouting should start in early July when aphids begin to be observed in fields. Sample 25 lower canopy leaves from each of 4 areas in the field (100 leaves in total). Count potato aphids and green peach aphids on each compound leaf, using a magnifying device to identify the species.

Economic threshold: For seed potatoes = 3-10 green peach aphids / 100 leaves. For processing potatoes = 30-100 green peach aphids / 100 leaves. There are no economic thresholds for buckthorn and potato aphids. These thresholds relate to transmission of potato leafroll virus and are not useful in determining infectivity relative to potato virus Y. No economic thresholds have been established for aphids that relate to potato virus Y transmission.

Leafhoppers

Typical damage: the potato leafhopper injects a toxin into the plant which results in hopper burn, a yellowing and curling of the tips and margins of the leaflets, which ultimately turn brown and brittle. Damaged plants die prematurely and yield may be reduced.

When and How to Monitor: Nymphs are scouted by visual inspection; sample 100 plants from 3-5 areas of the field.

Count the wingless nymphs on compound leaves taken from mid canopy. Adults are sampled with a sweep net (20 sweeps per location at 5 locations for a total of 100 sweeps).

Economic threshold: Nymphs-1 nymph per 10 leaves. Adults-1 leafhopper per sweep.

Colorado potato beetle

Typical damage: Larvae feeding may cause extensive defoliation of leaves and is capable of transmitting spindle tuber virus and bacterial ring rot.

When and How to Monitor: Start scouting for larvae 2 weeks after crop emergence. On field edges, count number of beetles on 20 separate plants. Record % defoliation of leaves. Repeated scouting is required since beetles have developed resistance to many insecticides and 2 generations may occur during the year.

Economic threshold: Economic threshold based on beetle numbers may vary by cost of treatment, expected returns and variety. Typical thresholds are 18 larvae/ 20 plants for Russet Burbank vs 6 larvae/20 plants for Norland. Treat when defoliation exceeds 10%.

Potato flea beetle

Typical damage: Beetle feeding causes "shot holes" in the leaves. Two generations may attack the foliage.

When and How to Monitor: Estimate feeding damage on the leaf or numbers of beetles on plants.

Economic threshold: Early in the season treat if greater than 10% defoliation. Later in the season (August) treat if greater than 25% defoliation or with greater than 65 beetles per plant for Norland or 300 beetles per plant for Russet Burbank.

Potatoes* Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate / Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Titan (N)	20.8 ml per 100 kg potato seed pieces			2,000
	Pyrifos 15G (OP)	0.1 kg per 100 metres of row	70	G	
	Pyrinex 480 EC (OP)	0.97 L (based on 90 cm row spacing)	70	G	409
	Thimet (OP)	0.14 kg in sandy soils 0.215 kg in heavy soils	At planting application. Do not harvest before 90 days after planting time.	G	27-31

Insect	Insecticide (and insecticide group ¹)	Rate / Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Cutworms (Army, darksided, pale western, redbacked, black, white)	Pounce / Perm-UP (I ²) Ambush (I ²)	73 – 158 ml 57-121 ml	Treat prior to 6-leaf stage	G	1276
	Ripcord (I ²)	70 ml	21	G	242-542
	Chlorpyrifos (Redbacked, black, and darksided cutworms only) (OP)	0.971 L (pre-plant) 0.486-0.971 L (seedling)	7	G	205 – 418
Sap or Fluid Feeders					
Aphids	Seed Piece Treatments				
	Actara 240SC (N)	See chart on label	N/A	Seed Treatment	>5,000
	Admire SPT / Alias 240 SC / Grapple / Grapple ₂ (N)	11.79 – 17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	N/A	Seed Treatment	>4,870
	Cruiser Maxx D Potatoes (N)	See chart on label	NA	Seed Treatment	
	Titan (N)	10.4-20.8 ml per 100 kg potato seed pieces	N/A	Seed Treatment	2,000
	In-Furrow Application				
	Actara 240SC (N)	0.15-0.20 L (based on 90 cm row spacing).		G	>5,000
	Admire 240 F / Alias 240 SC / Grapple / Grapple ₂ (N)	0.344-0.526 L (based on 90 cm row spacing).		G	4143-4870
	Foliar Sprays				
	Fulfill (HFB)	78.1 g	14	A or G	>5,000
	Beleaf (HFB)	49-65 g	7	G	
	Movento	89 – 148 ml	7	A or G	>2,000
	Actara 240SC (N)	44.1 ml	7	A or G	
	Actara 25WG (N)	42.5 g	7	A or G	>5,000
	Admire 240 F / Alias 240 SC / Grapple / Grapple ₂ (N)	81 ml	7	G	4143-4870
	Assail (N)	22.7-34.8 g	7	G	1,064
	Clutch (N)	28-43 g	14	A or G	4,300
	Concept (N + P)	263 ml	7	G	
	Lannate (C)	0.2185 kg	3	G	30-34
	Vydate (C)	0.93-1.21 L	7	G	9-10
	Malathion 500 (OP)	0.56-0.80 L	3	A or G	4302
	Malathion 85E (OP)	0.297-0.445 L	3	G	5,500
	Lagon / Cygon 480 EC / Cygon 480-AG (OP)	0.22-0.44 L	7	G	60-450
	Imidan (OP)	0.65 kg	7	G	285
	Orthene (OP)	0.30-0.44 kg	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	1	G	110
	Thionex EC (OC)	0.6-0.8 L	2	G	110
	Thiodan (OC)	0.6-0.8 L	1	G	107.2
Potato psyllid	Movento	89 – 148 ml	7	A or G	>2,000

Insect	Insecticide (and insecticide group ¹)	Rate / Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Leafhoppers	Seed Piece Treatments				
	Actara 240SC (N)	See chart on label	N/A	Seed Treatment	>5,000
	Admire SP/T / Alias 240 SC / Grapple / Grapple ₂ (N)	11.79 – 17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	N/A	Seed Treatment	>4,870
	Cruiser Maxx D Potatoes (N)	See chart on label	NA	Seed Treatment	
	Titan (N)	10.4-20.8 ml per 100 kg potato seed pieces	N/A	Seed Treatment	2,000
	In-Furrow Application				
	Actara 240SC (N)	0.15-0.20 L (based on 90 cm row spacing).		G	>5,000
	Admire 240F / Alias 240 SC / Grapple / Grapple ₂ (N)	0.344-0.526 L (based on 90 cm row spacing).	N/A	G	4143-4870
	Foliar Sprays				
	Actara 240SC (N)	44.1 ml	7	A or G	
	Actara 25WG (N)	42.5 g	7	A or G	
	Clutch (N)	28-43 g	14	A or G	4,300
	Pounce / Perm-UP (P)	73 – 105 ml	1	A or G	1276
	Decis SEC (P)	40 – 60 ml	3	A or G	395
	Matador / Silencer (P)	34 ml	7	A or G	64-110
	Ripcord (P)	25 – 50 ml	7	A or G	242-542
	UP-Cyde (P)	57 ml	7	A or G	355
	Concept (N + P)	263 ml	7	G	
	Sevin XLR Plus (C)	0.50 L	7	A or G	649
	Lannate (C)	0.2185 kg	3	G	30-34
	Vydate (C)	0.93-1.21 L	7	G	9-10
	Malathion 500 (OP)	0.56-0.80 L	3	A or G	1375-2800
	Malathion 85E (OP)	0.297-0.445 L	3	G	5,500
	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.22-0.44 L	7	G	60-450
	Dibrom (OP)	0.44 L	4	A or G	345
	Imidan (OP)	0.65 kg	7	G	285
	Orthene (OP)	0.30-0.44 kg	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	1	G	110
	Thionex EC (OC)	0.6-0.8 L	2	G	110
	Thiodan (OC)	0.6-0.8 L	1	G	107.2

Insect	Insecticide (and insecticide group ¹)	Rate / Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Lygus bugs	Pounce / Perm-UP (P) Ambush (P)	73 – 105 ml 57-81 ml	1	A or G	1276
	Decis 5EC (P)	40 – 60 ml	3	A or G	395
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Ripcord (P)	50 ml	7	A or G	242-542
	UP-Cyde (P)	81 ml	7	A or G	355
	Concept (N + P)	263 ml	7	G	
	Sevin XLR Plus (C)	2.12 – 2.59 L	7	A or G	649
	Vydate (C)	0.93-1.21 L	7	G	9-10
	Lagon /Cygon 480 EC (OP)	0.22-0.44 L	7	G	60-450
	Chlorpyrifos (nymphs only) (OP)	0.405 L	7	G	205 – 418
	Orthene (OP)	0.30-0.44 kg	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	1	G	110
	Thionex EC (OC)	0.6-0.8 L	1	G	110
	Thiodan (OC)	0.6-0.8 L	1	G	107.2
Defoliators					
Colorado potato beetle	Note: Colorado potato beetles have been found to be resistant to several families of insecticides in localized areas of Manitoba. Rotation between different families of insecticides is essential.				
	Seed Piece Treatments				
	Actara 240SC (N)	See chart on label	N/A	Seed Treatment	>5,000
	Admire SPT / Alias 240 SC / Grapple /Grapple ₂ (N)	11.79 – 17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	N/A	Seed Treatment	>4,870
	Cruiser Maxx D Potatoes (N)	See chart on label	NA	Seed Treatment	
	Titan (N)	10.4-20.8 ml per 100 kg potato seed pieces	N/A	Seed Treatment	2,000
	In-Furrow Application				
	Actara 240SC (N)	0.15-0.20 L (based on 90 cm row spacing).		G	>5,000
	Admire / Alias 240 SC / Grapple /Grapple ₂ (N)	0.345 -0.525 L	7	G	4143-4870
	Clutch (N)	108-181 g (based on 90 cm row spacing)	14	G	4,300
	Foliar Sprays				
	Rimon (SB)	0.17-0.33 L	14	G	>5,000
	Entrust (S)	20-40 g	7	G	>5,000
	Success (S)	34 – 67 ml	7	G	>5,000
	Delegate (S)	65-97 g	7	G	>5,000
	Coragen (D)	101 – 152 ml	14	A or G	>5,000
	Actara 240SC (N)	44.1 ml	7	A or G	
	Actara 25WG (N)	42.5 g	7	A or G	>5,000

Insect	Insecticide (and insecticide group ¹)	Rate / Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Colorado potato beetle <i>continued</i>	Admire / Alias 240 SC / Grapple / Grapple ₂ (N)	81 ml	7	G	4143-4870
	Assail (N)	16.2-32.4 g	7	G	1,064
	Clutch (N)	28-43 g	14	A or G	
	Pounce / Perm-UP (P) Ambush (P)	73 - 105 ml 57-81 ml	1	A or G	1276
	Decis 5EC (P)	40 - 60 ml	3	A or G	395
	Matador / Silencer (P)	34 - 50 ml	7	A or G	64-110
	Ripcord (P)	25 - 50 ml	7	A or G	242-542
	Up-Cyde (P)	57 ml	7	A or G	355
	Concept (N + P)	263 ml	7	G	
	Sevin XLR Plus (C)	0.50 L	7	A or G	649
	Vydate (C)	0.93-1.21 L	7	G	9-10
	Malathion 500 (OP)	0.56-0.80 L	3	A or G	4302
	Malathion 85E (OP)	0.297-0.445 L	3	G	5,500
	Dibrom (OP)	0.44 L	4	A or G	345
	Imidan (OP)	0.65 kg	7	G	285
	Chlorpyrifos (larvae only) (OP)	0.405 L	7	G	205 - 418
	Thionex 50W (OC)	0.45-0.61 kg	1	G	110
	Thionex EC (OC)	0.6-0.8 L	2	G	110
	Thiodan (OC)	0.6-0.8 L	1	G	107.2
Potato Flea Beetle	Admire SPT / Alias 240 SC / Grapple / Grapple ₂ (N)	11.79 - 17.69 ml per 100 pounds (45.36 kg) of potato seed tubers	N/A	Seed Treatment	>4,870
	Titan (N)	10.4-20.8 ml per 100 kg potato seed pieces			2,000
	Admire 240 F / Alias 240 SC / Grapple / Grapple ₂ (N)	Soil Application: 0.344-0.526 L (based on 90 cm row spacing).	N/A	G	4143-4870
	Pounce / Perm-UP (P) Ambush (P)	73 - 105 ml 57-81 ml	1	A or G	1276
	Decis 5EC (P)	40 - 60 ml	3	A or G	395
	Matador / Silencer (P)	34 ml	7	A or G	64-110
	Ripcord (P)	25 - 50 ml	7	A or G	242-542
	UP-Cyde (P)	57 ml	7	A or G	355
	Concept (N + P)	263 ml	7	G	
	Sevin XLR Plus (C)	0.50 L	7	A or G	649

Insect	Insecticide (and insecticide group ¹)	Rate / Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Potato Flea Beetle <i>continued</i>	Lannate (C)	0.2185 kg	3	G	30-34
	Vydate (C)	0.93-1.21 L	7	G	9-10
	Dibrom (OP)	0.44 L	4	A or G	345
	Imidan (OP)	0.65 kg	7	G	285
	Chlorpyrifos (OP)	0.405 L	7	G	205 - 418
	Orthene (OP)	0.30-0.44 kg	21	G	1030-1447
	Thionex 50W (OC)	0.45-0.61 kg	1	G	110
	Thionex EC (OC)	0.6-0.8 L	1	G	110
	Thiodan (OC)	0.6-0.8 L	1	G	107.2
Variegated cutworm	Coragen (D)	101-152 ml	1	A or G	>5,000
	Pounce / Perm-UP (P) Ambush (P)	73 ml 57 ml	1	G	1276
	Ripcord (P)	70 ml	7	G	242-542
	Sevin XLR Plus (C)	0.1-0.125 L / 300 m of row	7	A or G	649
	Lannate (C)	0.11 -0.22 kg	3	G	30-34
Armyworm	Coragen (D)	101-152 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
Stem Borers					
European Corn Borer	Rimon (SB)	0.17-0.33 L	14	G	>5,000
	Entrust (S)	35.4 g / acre	7	G	>5,000
	Success (S)	59 ml	7	G	>5,000
	Delegate (S)	65 g	7	G	>5,000
	Coragen (D)	101 - 152 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	7	A or G	64-110
	Pounce /Perm-UP (P) Ambush (P)	73 ml 57 ml	1	A or G	1276
	Concept (N + P)	263 ml	7	G	
	Sevin XLR Plus (C)	1.01 - 2.59 L	7	A or G	649

*Before using any pesticide on potatoes, consult the list of Agricultural Pesticides Approved for Use, available from Simplot Canada and McCain Foods (Canada).

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: SB=substituted benzoylurea, S=spinosyns, HFB=Homopteran feeding blockers, D=diamides, N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates, OC=organochlorines.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Rye

Information on typical damage, when and how to monitor, and economic thresholds for cutworms, aphids and armyworms in rye can be found in the section on field scouting in small grain cereals (wheat, barley, oats).

Rye Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Cereals Seed Treatment (N)	325 ml per 100 kg of seed		Seed treatment	
	Cruiser Maxx Vibrance Cereals Seed Treatment	325 – 650 ml per 100 kg of seed		Seed treatment	
Cutworms (Army, black, darksided, pale western, red-backed, white)	Pounce / Perm-UP (P) Ambush (P)	73 – 158 ml 57-121 ml	Treat prior to 6 leaf stage	G	1030
Sap Feeders					
Aphids	Malathion 500 (OP)	0.60-0.80 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
Defoliators					
Grasshoppers	Spreadable Bran Baits				
	Nolo Bait	Minimum of 0.45 kg		A or G	
	Eco bran (C)	0.8-1.6 kg	14	G	N/A
	Sprays				
	Sevin XLR Plus (C)	0.50-1.01 L	14	A or G	649
	Malathion 500 (OP)	0.69 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Lagon /Cygon 480EC (OP)	0.22 L (nymphs) 0.34-0.41 L (adults)	2-28 (see labels)	A or G	60-450
True armyworm	Sevin XLR Plus (C)	1.01-2.12 L	14	A or G	649
	Malathion 500 (OP)	0.60-0.80 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: N=neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Safflower Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
Defoliators					
Grasshoppers	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.22-0.40 L	21	A or G	60-450

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: D=diamides, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Small Grain Cereals (wheat, barley, oats)

BELOWGROUND AND SURFACE FEEDERS

Cutworms

Typical Damage: Notched, wilted, dead, or cut-off plants. Plants missing from rows, bare patches appearing in field.

When and How to Monitor: Look for cutworms, and evidence of cutworm feeding, when monitoring in late May to mid-July. Often cutworms will be close to the cut or shriveled plants they have just damaged. Cutworms will sometimes be most abundant in patches or a specific area of a field. In areas of the field where cutworm damage is noticeable, check around damaged plants in a 0.25 m² (50cm x 50cm) area. Use trowel or shovel to carefully search through top half to 1 inch of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m². Repeat in several locations to get an accurate assessment of what the cutworm levels are.

Economic Threshold: Pale western cutworm – 3-4/m²; Redbacked and army cutworm – 5-6/m². Well established fall-seeded crops or spring seeded crops with good moisture conditions can tolerate higher numbers. Sometimes it is most economical to just treat infested patches, and not whole fields.

SAP FEEDERS

Aphids

Typical Damage: Visible wilting of plants, yellow patches in fields, plants are sticky.

When and How to Monitor: Look for aphids when monitoring prior to the soft dough stage. While monitoring the field, using a sweep net or tapping plants over a white tray or bucket can alert you to the presence and relative abundance of aphids. If aphid levels appear concerning, a more thorough examination is needed. Count aphids on 20

randomly selected stems in each of 5 areas. Counts should be at least 50 paces apart, and observations should be made well into the center of the field. Too frequently farmers become alarmed after checking a few plants along the margins, especially near shelterbelts, where populations are high. Record the total number of aphids and calculate the average per plant.

Economic Threshold: 12-15 aphids/stem prior to the soft dough stage.

Barley Thrips

When and How to Monitor: Sampling should begin when the flag leaf is first visible and continue until the head is completely emerged from the boot. Barley thrips exhibit an edge effect; there are usually more thrips near protected field margins than other areas of the field. Most thrips can be found under the top 2 leaf sheaths. Unroll the leaf sheaths away from the stem to find the thrips.

Economic Threshold: Insecticide treatments are only effective when applied before heading is complete.

Treat when thrips are equal to or greater than the number calculated by:

Threshold (Thrips/stem) = (Cost of Control ÷ expected \$ value per bushel)/0.4

DEFOLIATORS

Grasshoppers

Typical Damage: Black strips along margins of newly emerging crops, head clipping later in season.

When and How to Monitor: Look for grasshoppers when monitoring fields from late – may through to harvest. Check along edges of crop, particularly areas adjacent to hayland, pastures and roadsides. Estimate number of hoppers/yard² (m²).

Economic Threshold: 8-13 grasshoppers/m². Early in the season, when grasshoppers are small, 18 grasshoppers/m² and visible crop damage may be a more appropriate threshold.

A rough estimate for an economic threshold for grasshoppers in crops to be used as greenfeed has been suggested at 20 grasshoppers/m² or higher.

Armyworms

Typical Damage: Leaves stripped from plants, awns chewed from heads, heads clipped.

When and How to Monitor: Check the soil surface for armyworms, and the plants for feeding, when monitoring in mid-June through early-August. At each stop shake plants and carefully check soil surface for dislodged larvae. During the day larvae may be under plant trash, soil clods or in soil cracks. Check the backs of armyworms for parasite eggs.

Economic Threshold: Four unparasitized larvae, small-

er than 2.5 cm (1 inch) per square foot. For migrating Armyworms: Treat a couple of swaths ahead of the infestation in the direction of movement to form a barrier strip.

SEED FEEDERS ONLY

Wheat Midge (wheat only)

When and How to Monitor: Monitor wheat in July when crop emerges from boot stage until flowering. Check crop canopy at dusk for signs of wheat midge adult activity. At each stop, examine 10 heads. Record the number of midge adults observed on or near heads. Calculate average number of midge per head.

Sticky traps may be used to capture adult midge activity in wheat fields.

Economic Threshold: For yield only: 1 adult midge per 4 to 5 heads. At this level of infestation, wheat yields will be reduced by approximately 15% if the midge is not controlled. To maintain optimum grade: 1 adult midge per 8 to 10 wheat heads during the susceptible stage.

Small Grain Cereals (wheat, barley, oats) Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ³
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Cereals Seed Treatment (N)	325 ml per 100 kg of seed		Seed treatment	
	Cruiser Maxx Vibrance Cereals Seed Treatment	325 - 650 ml per 100 kg of seed		Seed treatment	
	Raxil ProShield (N)	A co-pack of Raxil Pro and StressShield 600			
	Raxil WW (N)	A combination of Raxil MD and Stress Shield (imidacloprid).			
Cutworms (Army, dark sided, pale western, redbacked)	Decis 5EC (P)	80 ml	31 (oats) 40 (barley, wheat)	A or G	395
	Ricord (barley and wheat only) (P)	70 ml	21	G	242-542
	Pounce / Perm-UP (P)	73 - 158 ml	Treat prior to 6-leaf stage	G	1030
	Ambush (P)	57-121 ml			
	Chlorpyrifos (OP)	0.354-0.486 L	60	A or G	205-418
Sap and Fluid Feeders					
Aphids	Malathion 500 (OP)	0.60-0.8 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Cygon 480 EC /Cygon 480-AG (OP)	0.17 L	2	A or G	60-450
Thrips	Lannate (C)	0.1214kg	20	A or G	30-34
	Lagon /Cygon 480 EC /Cygon 480-AG (OP)	0.40 L	7-21 (see labels)	A or G	60-450
Brown Wheat Mite	Chlorpyrifos (OP)	0.253 L	60	A or G	205 - 418

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ³
Defoliators					
Grasshoppers	Spreadable Bran Baits				
	Nolo Bait	Minimum of 0.45 kg		A or G	
	Eco bran (C)	0.8-1.6kg	14 (oats, wheat) 28 (barley)	G	
	Sprays				
	Decis 5EC (P)	40 – 60 ml (ground) 60 ml (air)	31 (oats) 40 (wheat, barley)	A or G	395
	Ripcord (young grasshoppers only) (wheat and barley only) (P)	20 – 28 ml	30 (wheat) 45 (barley)	G	242-542
	Matador /Silencer (young grasshoppers only) (P)	25 – 34 ml (Ground) 34 ml (Aerial)	Do not apply within 28 days of harvest or 14 days of livestock foraging	A or G	64-110
	Sevin XLR Plus (C)	0.50-1.01 L	14 (wheat, oats) 28 (barley)	A or G	649
	Malathion 500 (OP)	0.68 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Chlorpyrifos (OP)	0.235-0.354 L	60	A or G	205-418
	Lagon /Cygon 480EC (OP)	nymphs-0.22 L adults-0.34-0.40 L	2-28 (see labels)	A or G	60-450
Cereal Leaf Beetle	Malathion 500 (OP)	0.22 – 0.45 L	7	A or G	4302
	Malathion 85E (OP)	0.435 L	7	A or G	5,500
True Armyworm	Delegate (S)	40-81 g	21	G	>5,000
	Matador /Silencer (P)	34 ml	Do not apply within 28 days of harvest or 14 days of livestock foraging	A or G	64-110
	Sevin XLR Plus (C)	1.01-2.12 L	14 (wheat, oats) 28 (barley)	A or G	649
	Lannate (C)	0.1093-0.2185kg	20	A or G	30-34
	Malathion 500 (OP)	0.60-0.80 L	7	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Chlorpyrifos (OP)	0.354 – 0.486 L	60	A or G	205-418
Slugs	Sluggo Professional	10-20 kg		G	>5,000

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Pests of Seed Only					
Wheat Midge (a pest of wheat only)	Rotate Crops – Continuous wheat cropping encourages higher wheat midge populations.				
	Resistant Varieties – the following varieties are resistant to feeding by wheat midge: Red spring wheat varieties: AC Goodeve, AC Unity, AC Fieldstar, AC Shaw, CDC Utmost, and AC Vesper. Extra-strong wheat variety: AC Glencross.				
	Biological Control - A parasitoid, <i>Macroglanes penitrens</i> , was found to control an average of 32% of the wheat midge in Saskatchewan.				
	Chlorpyrifos (OP)	0.336-0.405 L	60	A or G	205-418
	Lagon / Cygon 480 EC / Cygon 480-AG (OP)	0.40 L	21	A or G	60-450
Stem-Borers					
Hessian Fly	<ul style="list-style-type: none"> • Never plant wheat in the same field 2 years in a row in areas where Hessian flies are a problem. • The spring wheat cultivar Superb is partially resistant to the Hessian fly. • Early seeded spring wheat is less susceptible to stem breakage caused by Hessian fly than later seeded wheat. • Winter wheat planted in September will likely be free of Hessian flies. 				
Wheat Stem Maggot	Crop rotation and stubble cultivation may reduce populations				
Wheat stem sawfly	Solid-stem wheat varieties (such as the hard red spring wheat varieties AC Lillian, AC Abbey and AC Eatonia) can reduce damage by wheat stem sawfly larvae compared to susceptible varieties, however the level of control can vary depending on environmental conditions.				

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: S=spinosyns, N= neonicotinoids, P=pyrethroids, C=carbamates, OP=organophosphates

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Scouting for insects in Soybeans

Soybean Aphid

Typical Damage: Soybean aphids suck sap from soybean plants. Infested leaves may wilt or curl when infestations are large. Other symptoms may include plant stunting, reduced pod and seed count, and yellowing of leaves.

When and How to Monitor: Check 30 plants (6 plants in 5 areas) per field. Examine the entire plant and estimate populations of soybean aphids (counting exact numbers will not be possible or practical with higher populations). Once soybean aphid numbers reach 250 aphids per plant, scout the field frequently to determine if soybean aphid numbers are increasing. A population can stay at 250-300 aphids per plant and not cause economical yield loss. If the aphid levels are not rising above 250-300 per plant, there is a good indication that field conditions are favouring natural enemies (such as beneficial insects and fungi) that are helping control the aphids.

Economic Threshold: When there are on average at least 250 aphids per plant and the population is increasing, and the plants are in the R1 (beginning bloom) to R5 (beginning seed) growth stages, treatment would be economical. This threshold gives an approximate 7-day lead time before aphid populations are expected to exceed the economic injury level (670 aphids per plant), where cost of control is equal to yield loss. When soybean aphid populations are not actively increasing above 250 aphids per plant, natural enemies are keeping up with the aphid population. Do not use an insecticide in this case, as it will kill the natural enemies which may enable the aphid population to increase above the economic injury level.

Soybean Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Cruiser Maxx Vibrance Beans (N)	A seed treatment combining Cruiser Maxx Beans and Vibrance 500FS			
	Stress Shield 600 (N)	Seed Treatment			
Seedcorn Maggot	Cruiser Maxx Vibrance Beans (N)	A seed treatment combining Cruiser Maxx Beans and Vibrance 500FS			
	Stress Shield 600 (N)	Seed Treatment			
	Agrox CD, Agrox B-2 (OP)	See "Disease Control-Seed Treatments" section of this guide			N/A
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
	Matador /Silencer (P)	34 ml	21	A or G	64-110
Sap or Fluid Feeders					
Soybean Aphid	Movento	75-111 ml	21	A or G	>2,000
	Matador /Silencer (P)	34-94 ml (Matador) 34 ml (Silencer)	21	A or G	64-110
	Concept (N + P)	132 - 263 ml	20	G	
	Lagon /Cygon 480 EC /Cygon 480-AG (OP)	0.28-0.40 L	30	A or G (lagon, cygon 480 EC) G only (Cygon 480-AG)	60-450
Leafhoppers	Lagon /Cygon 480 EC /Cygon 480-AG (OP)	0.28-0.40 L	30	A or G (lagon, cygon 480 EC) G only (Cygon 480-AG)	60-450
Lygus bugs	Matador /Silencer (P)	34 ml	21	A or G	64-110
	Lagon /Cygon 480 EC /Cygon 480-AG (OP)	0.28-0.40 L	30	A or G (lagon, cygon 480 EC) G only (Cygon 480-AG)	60-450
Spider mites	Lagon /Cygon 480 EC /Cygon 480-AG (OP)	0.40 L	30	A or G (lagon, cygon 480 EC) G only (Cygon 480-AG)	60-450
Defoliators					
Corn Earworm	Coragen (D)	101-152 ml	1	A or G	>5,000
Grasshoppers	Matador /Silencer (P)	34 ml	21	A or G	64-110

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: N= neonicotinoids, P=pyrethroids, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Stored Grain Insect Control-

See Insect Control in Stored Grain; after Insect Management Charts (pages 494-497).

Summerfallow-

See grasshopper management on Pastures, etc.

Scouting for insects in Sunflowers

BELOWGROUND AND SURFACE FEEDERS

Cutworm

Typical Damage: Notched, wilted, dead, and cut-off plants (weed or crop seedlings). Plants missing from rows, bare patches appearing in field.

When and How to Monitor: Look for cutworms, and evidence of cutworm feeding, when monitoring sunflowers in late May to mid-July. Often cutworms will be close to the cut or shriveled plants that they have just damaged. Cutworms will sometimes be most abundant in patches or a specific area of a field. In areas of a field where cutworm damage is noticeable, check around damaged plants in a 0.25 m² (50cm x 50cm) area. Use trowel or shovel to carefully search through top half to 1 inch of soil for cutworm larvae. Multiply the number of cutworms found by 4 to get the number per m². Repeat in several locations to get an accurate assessment of what the cutworm levels are.

Nominal Threshold: 1 cutworm or more per square foot (30 by 30 cm) or if there is a 25 to 30% stand reduction. Sometimes it is most economical to just treat infested patches, and not whole fields.

DEFOLIATORS

Sunflower Beetle

Typical Damage: Adults: Leaves of seedling plants chewed or completely destroyed late May through June, shot-holes or large areas of leaves chewed July through August. Larvae: Leaves of plants chewed or completely destroyed.

When and How to Monitor: Adults: Look for when monitoring sunflower seedlings in May through June. Examine 10 plants at random at each stop. Larvae: Look for when monitoring sunflowers in July through mid-August. Examine 10 plants at random at each sampling site. Peel back leaves around growing tip and record total number of larvae found. Calculate average number per plant.

Economic Threshold: Adults: 1-2/seedling; Larvae: 10 to 15/plant or 25-30% defoliation.

Insects affecting the seeds

PESTS OF SEED ONLY

Red Sunflower Seed Weevil

Typical Damage: Seeds partly or completely destroyed, exit hole in hull. Shriveled kernels, kernels completely destroyed.

When and How to Monitor: Monitor fields when ray petals being to form and continue every 2 to 3 days until pollination is complete. When scouting, use the X pattern and begin counts at least 70 to 100 feet into the field to avoid margin effects. Examine 5 plants at each site for a total of 25 plants. For checking individual sunflower heads, brush the face of the head vigorously to bring the weevils to the surface, or use a commercial preparation of mosquito repellent containing diethyl toluamide (DEET) to spray the heads. This will cause the weevils to move out of hiding spots. Record total number of weevils and calculate average per head.

Economic Threshold:

Confection Sunflowers: 1-2 weevils/plant. Control is based on a need to keep seed damage below 3 or 4% because of industry standards.

Oilseed sunflowers: 12-14 weevils/head.

The ideal plant stage to treat is when most plants in the field are at 40% pollen shed (R5.4).

Banded Sunflower Moth

When and How to Monitor: Look for banded sunflower moth adults when monitoring fields in the late bud (R-4) to early bloom (R5.1) plant growth stage. Count moths on 20 plants from 5 different sites for a total of 100 plants. Sampling in early evening or early morning when the moths are most active gives the most accurate counts.

Sampling strategies based on scouting for adult moths during daylight hours, and counting eggs, have also been developed.

Economic Threshold: 1 moth per 2 plants when monitoring in the early evening or early morning.

If monitoring for eggs or adult moths during daylight hours, tables for determining economic thresholds can be found at: <http://www.ag.ndsu.edu/pubs/plantsci/pest/e823.pdf> If treatment is warranted, it should be applied at the R5.1 sunflower plant growth stage.

Lygus bugs

Economic Threshold: Confection - One adult lygus bug per 9 heads can result in economic losses through the reduction in seed quality. Lygus bug management should be initiated between the R4 to R5.1 stage if adult densities reach economic levels. Oilseed sunflowers are not believed to be at risk of damage from lygus bugs.

Sunflowers Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Belowground and Surface Feeders					
Wireworms	Wireworms may sometimes damage sunflowers. Seeding sunflowers when the soil temperature is at least 8 to 10°C at 1 to 1.5 inches depth may minimize damage by wireworms.				
	Cruiser Maxx Sunflowers (N)	A seed treatment combining Cruiser 5FS with Maxim 480FS and Apron XL. Sunflower Seeds can not be treated with Cruiser Maxx Sunflowers in Canada.			
Cutworms	Coragen (D)	101 ml	1	A or G	>5,000
	Pounce / Perm-UP (P)	73 – 158 ml	Treat prior to 6-leaf stage	G	1030
	Ambush (P)	57-121 ml			
	Chlorpyrifos (OP)	0.486 L	42	G	205-418
Defoliators					
Sunflower Beetle	Cruiser Maxx Sunflowers (N)	A seed treatment combining Cruiser 5FS with Maxim 480FS and Apron XL. Sunflower Seeds can not be treated with Cruiser Maxx Sunflowers in Canada			
	Decis 5EC (P)	40 ml	70	A or G	395
	Matador / Silencer (P)	17– 25 ml(Ground) 34 ml (Aerial)	7	A or G	64-110
	Ripcord (P)	28 ml	70	A or G	242-542
	UP-Cyde (P)	40 ml	70	A or G	355
Pests of Head and Seeds					
Lygus bugs	Matador	34 ml	7	A or G	64-110
Sunflower Seed Weevil	Early planting helps to reduce seed damage by sunflower seed weevils				
	Ripcord (P)	28 ml	70	A or G	242-542
	UP-Cyde (P)	40 ml	70	A or G	355
	Chlorpyrifos (OP)	0.486 L	42	A or G	272/205
Banded sunflower moth	Late planting may provide some control				
	Coragen (D)	101 - 152 ml	1	A or G	>5,000
Sunflower Moth	Dipel 2X DF (young larvae) (M)	127 – 253 g	N/A	A or G	> 4,000
Sunflower Midge	Some cultivars show some resistance to feeding by sunflower midge				

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: M=microbial, D=diamides, P=pyrethroids, C=carbamates, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Field Scouting in Sweet Clover

DEFOLIATORS

Sweetclover Weevil

Typical Damage: Adults chew crescent-shaped and jagged notches in leaves and can completely defoliate plants.

When and How to Monitor: Inspect clover seedlings for weevil damage in spring as the seedlings emerge. In mid-summer and throughout August, inspect first-year clover

stands for damage along crop margins. Invading weevils move into these stands only as far as necessary to satisfy their food requirements, so an insecticide application to affected field margins is usually all that is required. Visually estimating the number of weevils per plant must be done carefully because weevils fall from plants easily and are difficult to see on the ground.

Economic Threshold: 1st year stands: 1 weevil adult/3 seedlings (1/5 seedlings under dry conditions). 2nd year stands: 9-12 weevil adults/plant.

Sweet Clover Insect Management Chart

Insect	Insecticide (and insecticide group ¹)	Rate/Acre	Preharvest interval (days)	Application (A=aerial; G=ground)	LD ₅₀ (Mammalian Toxicity) ²
Defoliators					
Grasshoppers	Spreadable Bran Baits				
	Eco bran (C)	0.8-1.6kg	2	G	N/A
	Sprays				
	Sevin XLR Plus (C)	0.50-1.01 L	2	A or G	649
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
Sweetclover Weevil	Locate new seedlings as far as possible from 2nd-year clover. Cultivating second-year stands of sweet clover silage and hay as soon as possible after the crop is taken kills the new-generation weevil larvae in the soil.				
	Malathion 500 (OP)	0.56-1.01 L	7-Cattle may be returned immediately after spraying.	A or G	4302
	Malathion 85E (OP)	0.445-0.544 L			5,500
	Lagon /Cygon 480 EC / Cygon 480-AG (OP)	0.34-0.45 L	28	A or G	60-450
Alfalfa weevil	Coragen (D) (suppression only)	152-202 ml	0	G	>5,000
Blister Beetle	Sevin XLR Plus (C)	1.01-1.6 L	2	A or G	649
Beet Webworm	Sevin XLR Plus (C)	1.01-2.12 L	2	A or G	649
Sap or Fluid Feeders					
Lygus Bugs	Dibrom (OP)	0.44-0.88 L	4	A or G	345
Leafhoppers	Sevin XLR Plus (C)	1.01-1.6 L	2	A or G	649
	Malathion 85E (OP)	0.445-0.544 L	7	A or G	5,500
	Dibrom (OP)	0.44-0.88 L	4	A or G	345
Aphids	Dibrom (OP)	0.44-0.88 L	4	A or G	345

ALWAYS CONSULT THE INSECTICIDE LABEL BEFORE APPLYING ANY INSECTICIDE.

¹ Insecticide Group: C=carbamates, OP=organophosphates.

² LD₅₀ values represent the relative toxicity of a pesticide. They represent the dose (in mg/kg body weight) that will kill 50% of the test animals. Thus the lower the number the greater the toxicity. Values given are for oral LD₅₀.

Timothy-

See forage grasses

Wheat-

See small grain cereals

Insect Control in Stored Grain

Prevention

Clean in and around storage facilities. Grain storage facilities, and the area around storage facilities, should be cleaned thoroughly prior to storing grain.

Clean equipment used to move grain. Grain left in equipment throughout the summer months can result in new grain that is being placed into storage becoming infested. Combines, truck beds, grain wagons, augers and other equipment used to move grain should be cleaned of grain residue. Other potential sources of grain infesting insects include livestock feeds, old seed bags, spilled grain, etc.

Inspect grain storage facilities for signs of deterioration, especially for leaks or holes through which insects or rodents can gain access to the stored grain. Moving and storing the grain in clean facilities will eliminate one source of infestation. However, grain stored for long periods of time still has the potential for renewed infestations.

Treating storage facilities. Depending on the commodity to be stored, storage facilities may additionally be sprayed or dusted, if needed, with a recommended insecticide before storing grain in the bin (e.g. malathion, diatomaceous earth or cyfluthrin – refer to product labels for details). *Note* – some commodities, such as canola, flax and sunflowers, should not be stored in facilities recently treated with malathion or cyfluthrin (Tempo).

Dry and Cool Grain. Ideally, the grain should be dry before being put into storage, and cooled as quickly as possible. For long-term storage, producers are urged to lower the grain temperature below 15°C as soon as possible after the grain is placed in storage. At 15°C the stored product insects stop laying eggs and development stops. Aeration systems used during the night immediately after harvest should have the grain below 15°C in about 2 weeks. Grain that is not moved or aerated after harvest can remain warm enough to allow insects to survive the winter. Convection currents arising from this warm air can also promote condensation, sprouting (heating) and mould growth in unmanaged grain. These conditions are very attractive to stored product pests and support their development.

Once the grain mass is cooled to the desired temperature, fans should be sealed to prevent unwanted air migration through the mass that could result in early grain mass warm-up. Cold grain has a longer storage life than warm grain.

Note however that under cool grain temperatures, insect movement is reduced to the point that some insecticides may not be effective.

Monitoring for Insects

Regular monitoring of the stored grain is the next step in determining the presence and potential for serious infestations. Either the presence of insects or damaged kernels will give an indication of a problem. One means of detecting insects in stored grain is through the use of probe traps, available from numerous crop protection agents. Often the

first indication of an infestation will be found near the top centre of a storage bin, and therefore, this is where traps should be placed. Monitoring should take place once every 7-10 days during the onset of storage (first 60 days) and then the frequency of monitoring may be adjusted.

Identifying insects in stored grain

Correct identification of insects found in stored grain is important in determining the most appropriate control methods. Some of the insects found in stored grain feed on the grains, while others feed on fungus that may be developing in the stored grain.

Grain feeders:

Insects that feed on the grain include rusty grain beetles, red flour beetles, and sawtoothed grain beetles.

The **rusty grain beetle** is the most common stored product insect. Heavy infestations of this insect cause grain to heat and spoil.

The **red flour beetle** is the another common insect pest of stored grain in the prairies. Red flour beetles cannot feed on undamaged, dry seed with less than 12% moisture content. They prefer grain dust, broken grain and milled stocks. Red flour beetles can be controlled by moving grain, whether or not it is cold.

Sawtoothed grain beetles are more common in stored oats than in stored wheat and barley.

Fungus feeders:

Insects that feed on fungus in the grain bin include the foreign grain beetle, hairy fungus beetle, psocids, and grain mites.

Foreign grain beetles resemble the rusty grain beetle, but can be distinguished from it by club-shaped antennae. Also, when placed in a glass jar, foreign grain beetles will climb up the sides, while rusty grain beetles cannot. While foreign grain beetle is considered a fungus feeder, they will feed on grain if the moisture content is in the high end of the acceptable range (eg. 14.5% mc wheat).

Grain mites are whitish, about 0.2 to 0.5 mm long, and can be hard to see with the naked eye. About eight kinds of mites are common in farm granaries and elevators.

Psocids are soft-bodied insects, about 1 mm long, with long antennae relative to the body size.

Fungus feeding insects and mites cannot survive in dry grain. Chemical control is not necessary for fungus feeding pests in stored grain. Practices that result in the grain drying may be all that is needed to control such pests.

Information to help identify insect pests of stored grain can be found at: <http://www.grainscanada.gc.ca/storage-entrepose/keys-cles/sgp-irg/sgp-irg-eng.htm>

Control Techniques:

The Canada Grain Act states there is zero tolerance for any primary insects (those that feed on whole, sound grain) in grain delivered to elevators. Outlined below are some control techniques and when and how these techniques can be best used.

Cold Temperatures

Rusty grain beetles are cold hardy and can survive subzero temperatures. Rusty grain beetles and other stored grain insects can be killed by reducing core grain temperatures as follows:

Time Required to Kill Insects at Various Grain Temperatures	
Grain Temperature	Time required to kill insects
-5°C	12 weeks
-10°C	8 weeks
-15°C	4 weeks
-20°C	1 week

Cooling the grain, through aeration or moving the grain several times during mid-winter, should provide effective control of rusty grain beetles.

Phostoxin

Company: Degesch America Inc. (Phostoxin); PCP#15736 (Round tablets).

Formulation: 55% aluminum phosphide.

Formulation	Primary Use	Container Size
Phostoxin tablets (3g each)	On the farm or country elevator	500 tablets
Phostoxin pellets (0.6g each)	In terminals	2500 pellets
Phostoxin tablets prepac	Containers	4 strips of 33 tablets to a pouch

Insects and other pests controlled: Rusty grain beetle, red flour beetle, sawtoothed grain beetle, granary weevil, Indian meal moth, yellow mealworm, lesser grain borer, nematodes, mice and rodents.

Approved for use on the following stored grains: Barley, corn, dried peas, lentils, millet, oats, rye, soybeans, sunflower seeds, triticale and wheat.

Restricted Product: The use and sale of Aluminum Phosphide (Phostoxin) is restricted to licensed pesticide applicators possessing a valid fumigation licence (Saskatchewan) or stored agricultural products license (Manitoba).

Phostoxin can only be used in conjunction with a detailed fumigation management plan.

Rate and Minimum Exposure Period: Refer to labels to determine rate. For grain bins a dosage of 250-500 tablets (or 880-2560 pellets) per 100m³ of bin space being treated (not volume of grain) is recommended. It is important to ensure that bins are relatively secure. It is not advisable to

High Temperatures

All four stages of insects (eggs, larvae, pupae and adults) will die if subjected to high temperatures for a sufficient period of time. The most realistic use of high temperatures for controlling insects would be when the grain was tough and in need of drying. In this case, the insects would be killed at the same time the grain is dried. The insects need to be exposed to temperatures in the range of 50 to 55°C for approximately 15 minutes. Note – the baking quality of wheat is damaged if the temperature of the grain reaches 60°C for any significant length of time.

Moving Grain

Moving grain using cyclone-based pneumatic conveyors (grain vacs) at about 200 bushels per hour has been shown to be an effective means of controlling insects in stored grain. Loading the grain using a pneumatic grain conveyor removes insects from grain being delivered to elevators.

use phosphine products in bins that are leaky or not well sealed.

The following table may be used as a guide to determine the minimum length of exposure period to phostoxin at the indicated temperatures:

Temperature	Exposure Period
Below 5°C (40°F)	Do not fumigate
5°C-12°C (40°-53°F)	10 days
13°-15°C (54°-59°F)	5 days
16°-20°C (60°-68°F)	4 days
above 20°C (68°F)	3 days

If the grain is less than 5°C then the tablets will not release the gas until the grain temperature warms up. This may result in poor control and accidental exposure to phostoxin at a later date in grain handling facilities. Very dry grain will also slow the release of the gas from the pellets. A shortened exposure period cannot be compensated for by increased dosage. Also ensure that storage is well ventilated for at least 24 hours after the required time for fumigation.

Protect-It, Insecto

Company: Hedley Technologies Ltd. (Protect-It)
PCP#24259; Natural Insecto Products Inc. (Insecto)
PCP#22489

Formulation: 90 percent Diatomaceous Earth (DE)

Insects controlled: Rusty grain beetle (Protect-It only), rice weevil, granary weevil, Angoumois grain moth, Mediterranean flour moth, Indian meal moth, red flour beetle and Tribolium.

Approved for use on the following stored products: Feed grains, seed, stored grains, wheat, barley, buckwheat, corn, oats, rye, flax, peas, soybeans and sorghum. Also registered for structural treatment of empty grain storage and transportation containers.

How it works: Diatomaceous earth damages the cuticle of the insect, reducing the insect's ability to retain moisture. The insect eventually dies from dehydration.

Rate, for empty storage structures: Use a dust blower or aeration fan to get diatomaceous earth into the cracks, crevices and void spaces of the structure being treated. Dust areas at a rate of 1 kg per 200 square metres (5 g/m²).

Rate, while grain is being placed into storage: Protect-It; The application rate for Protect-It varies by crop and insect species, ranging from 100 g/tonne for control of rusty grain beetle in wheat to 1000 g/tonne for red flour beetle in corn. Refer to the label for details.

Insecto; Apply to grain at the time of storage at a rate of 0.5 to 1 Kg per metric ton of grain (500-1000 ppm).

Precautions: The application of DE will lower the test weight measurement of the grain, but usually not to the point of downgrading. If test weight loss is excessive, the grain can be diluted with untreated grain. DE is non-toxic to humans and animals.

Malathion Grain Dust

Company: Interprovincial Co-operative (Malathion Grain Protectant Dust) PCP#17222; United Agri Products (Malathion Grain Protector Dust) PCP# 15896

Formulation: 2% malathion

Insects controlled: confused flour beetles, flat grain beetles, granary weevil, Indian meal moth, lesser grain borer, rusty grain beetle and sawtoothed grain beetle.

Approved for use on the following stored grains: Wheat, rye, barley and oats as stored grains.

Malathion Grain Dust can be applied to grain as it is being loaded into a bin or being turned by adding gradually at the grain auger. It can also be used to control surface infestations by applying to the grain surface and raking in to 15 cm depth of the grain. Malathion controls insects by ingestion and contact and insects must be active for it to be effective.

Rate:

Crop	Rate-g/1000 kg (tonne) grain
Wheat	415
Rye	450
Barley	520
Oats	735

Do not apply to grain within 7 days of sale.

Be aware that the Canadian Grain Commission allows only 8 ppm of malathion residues in stored grains.

Malathion 500, Malathion 85E

Refer to labels for these products for insect and mite control in empty grain bins, grain elevators, grain box cars and flour mills.

Note – Some commodities, such as canola, should not be stored in facilities recently treated with malathion.

Malathion 500 (IPCO)

Insect	Rate	Note
Rusty grain beetle, sawtoothed grain beetle, confused flour beetle, grain mite, granary weevil, Indian meal moth, lesser grain borer (empty grain bins)	250-300 ml/5 L of water on 100 m ²	May be used within 1 day of grain storage

Malathion 85E (United Agri Products),

Insect	Rate	Note
Rusty grain beetle, red flour beetle, sawtoothed grain beetle, confused flour beetle, grain mite, granary weevil, Indian meal moth, lesser grain borer, flat grain beetles, rice weevils (empty grain storage facilities)	Mix 490 ml in 15 L of water. Apply 5 L of mixture on 100 m ²	Wait until spray has thoroughly dried before storing grain in treated areas.

Tempo 20 WP

Company: Bayer CropScience PCP#25673

Formulation: 20% cyfluthrin. Tempo is a group 3 (pyrethroid) insecticide.

Application: Tempo can be used to control insects in grain storage facilities, truck beds and other areas where grain

is stored *before filling these areas with grain*. Cleaning of all areas prior to use of Tempo 20 WP insecticide will increase levels of control. See the insecticide label for specific mixing instructions.

Insecticide Product Pages

For rates and preharvest intervals for insecticides, see the insect management charts on pp. 457-492.

Actara

Insecticide Group – 4A

(Refer to page 456)

Company:

Syngenta

Formulation:

Actara 240SC (PCP#28407): 240 g/L thiamethoxam formulated as a soluble concentrate.

Container size - 2 X 2.04 L

Actara 25WG (PCP#28408): 25% thiamethoxam formulated as a water dispersible granule.

Container size - 4 X 850 g

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, aphids, potato leafhopper

Application:

***Actara* 240SC - Soil application:** Apply as an in-furrow spray during planting to allow the insecticide to be absorbed by plant roots. For 90 cm row spacing, apply 151 to 196 mL/acre. Use the higher rate for extended control. DO NOT follow a soil application with a foliar application.

Potato seed piece treatment: Choose the appropriate rate from the chart on label, based on seeding rate. Apply only in areas with adequate ventilation or in areas that are equipped to remove mist or dust. Best results are obtained if potatoes are planted immediately after *Actara* 240SC is applied to seed. When transporting cut and treated seed ensure the seed is covered. DO NOT apply a subsequent treatment of in-furrow or foliar application of thiamethoxam or other Group 4 insecticide following seed piece treatment with *Actara* 240SC.

Foliar application: *Actara* may be applied by ground or air. For ground application use a minimum of 40 L / ac unless otherwise indicated on label. A maximum of 2 foliar applications of *Actara* may be made per season. DO NOT exceed a total of 88 g/acre. Allow at least 7 days between applications. DO NOT use a foliar application of *Actara* following in-furrow or soil application of *Actara*.

How it Works:

Actara is a systemic (taken up into the plant foliage after application), chloronicotinyl insecticide.

Restrictions:

Rainfastness:

Avoid application of this product when heavy rain is forecast. *Actara* is rainfast once spray has dried on treated plants.

Preharvest Interval:

DO NOT harvest within 7 days of application.

Re-Entry:

DO NOT re-enter treated areas for 12 hours after foliar application.

Re-cropping: No restrictions following the harvest of sorghum, wheat, barley, canola, potatoes or cover crops. For all other crops 120 day plant-back interval is required.

Tank mix: Potatoes - *Actara* 240SC can be mixed with *Quadris*® Flowable fungicide and *Ridomil*® Gold 480SL fungicide (or *Ridomil* Gold 480EC fungicide).

Buffer Zones: Buffer zones are required for the protection of terrestrial and freshwater habitats. Refer to specific label for buffer zones required.

Precautions:

Actara is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Wait a minimum of 5 days after treatment before placing beehives in a treated field.

Hazard Rating:



Actara 240SC: Caution – Poison

Actara 25WG: Caution – Eye and skin irritant

For an explanation of the symbols used here see page 11.

Assail

Insecticide Group – 4A

(Refer to page 456)

Company:

Distributed by Engage Agro Corp. (PCP#27128)

Formulation:

Acetamiprid formulated as a wettable powder-70% by weight

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa (seed production only)	Alfalfa plant bug, Lygus bugs (suppression only)
Potato	Colorado potato beetle, aphids

Application:

Ground application only. DO NOT apply by air

Apply with a minimum finished spray volume of 80 litres per acre. For best results uniform spray coverage of the host plants is important.

Begin application when insect levels reach economic thresholds. Use higher rates for heavy infestations, dense foliage or for adult stages of the Colorado potato beetle. Residual control will depend on environmental factors, plant growth, application rate and level of insect infestation.

How it Works:

Assail is a chloronicotinyl insecticide that works by contact or ingestion. It has an anti-feedant effect that can prevent pest damage to host plants prior to the death of the insect. This product rapidly degrades in the soil with no carryover effects.

Restrictions:

DO NOT make more than 2 applications per year per crop.

DO NOT apply more than once every 7 days.

DO NOT exceed a total of 48 g active ingredient (68.8 g product) per acre per season.

DO NOT apply less than 7 days prior to harvest (Preharvest interval).

Buffer Zones: An untreated buffer zone between the last spray swath and the edge of aquatic systems (such as rivers, streams, lakes, and other water bodies) must be established. For groundboom sprayers - 20 metres. DO NOT apply acetamiprid directly to water or to areas where surface water is present. Buffer zone required for sensitive terrestrial areas (grasslands, forested areas, shelterbelts, woodlots, hedgerow, rangelands) – 2 metres.

Re-entry Interval: DO NOT re-enter treated areas for a period of 12 hours after application.

Precautions:

DO NOT apply when honey bees are present in the area to be treated as acetamiprid is toxic to honey bees if exposed to direct treatment.

If this product is to be applied to a product destined for export to the United States, information on acceptable residue levels are available at www.croplife.ca.

Storage: DO NOT store in or around the home. Store unused product in a cool, ventilated, dry, locked area. DO NOT allow prolonged storage in areas where temperatures frequently exceed 46 degrees C.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Beleaf

Insecticide Group – 9C

(Refer to page 456)

Company:

FMC of Canada (PCP#29796)

(Distributed by United Agri Products)

Formulation:

50% flonicamid formulated as a water soluble granule

Container size - 1 to 100 kg

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa (seed production only)	Aphids
Potato	Aphids

Application:

DO NOT apply by air.

Ensure the spray system is clean and free of residues from previous applications. Fill the tank half full with clean water. Ensure agitation system is operating and sufficient to provide uniform spray mixing during application and until the spray tank has been emptied. Complete filling to the desired solution volume.

Thorough spray coverage of plant foliage is essential for optimum control. Apply in sufficient water volumes to ensure good coverage - Use a minimum of 38 litres per acre of water. Rates and finished spray volumes should be increased under extreme pest populations or dense plant foliage.

Scout fields and reapply if necessary.

How it Works:

Flonicamid insecticide is a member of the pyridinecarboxamide class of chemistry and controls target pests by contact and ingestion provoking rapid and irreversible feeding cessation.

Restrictions:

Allow a minimum of 7 days between applications.

DO NOT make more than 3 applications per year.

DO NOT apply more than apply more than 64 grams per acre of *Beleaf* per application.

DO NOT apply more than 192 grams per acre of *Beleaf* per season.

DO NOT use *Beleaf* in greenhouses or home gardens.

Re-cropping: There are no plant-back restrictions for potatoes. All other crops may be planted 30 days after the last application of *Beleaf*.

Precautions:

Avoid overnight storage of spray mixture. Prepare only enough spray mixture required for immediate application.

DO NOT use liquid fertilizer as a carrier for *Beleaf* insecticide.

Beleaf insecticide should not be used with spray adjuvants.

Avoid application when heavy rain is forecast.

DO NOT enter or allow entry into treated areas for 12 hours after application.

Storage: Store product in original container, in a secured, dry place separate from other pesticides, fertilizer, food or feed.

Flonicamid is toxic to beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site. Flonicamid is toxic to non-target terrestrial plants. Observe buffer zones for sensitive areas (e.g. aquatic habitats, forested areas) as specified on label directions.

If this product is to be applied to a commodity destined for export to the United States, visit Crop Life Canada's website www.croplife.ca for information on acceptable residue limits.

Hazard Rating:



Caution – Eye irritant

For an explanation of the symbols used here see page 11.

Chlorpyrifos

Insecticide Group – 1B

(Refer to page 456)

Company:

Dow AgroSciences (*Lorsban 4E* – PCP#14879)

MANA Canada (*Pyrinex 480EC* – PCP#23705;

United Agri Products (*Pyrifos 15G* – PCP#24648)

Cheminova Inc. (*Nufos 4E* – PCP#25831)

IPCO (*Citadel 480EC* – PCP#27479)

Loveland Products Inc. (*Warhawk 480EC* – PCP#29984)

NewAgco Inc. (*MPOWER Krypton* – PCP#30985)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Formulations:

Citadel, *Nufos*, *Lorsban* and *Pyrinex* - 480 g/L chlorpyrifos formulated as an emulsifiable concentrate. Container sizes (Note that container sizes may vary between products) - 10 L jug, 115 L returnable container, 208 L drum.

Pyrifos - 15% chlorpyrifos formulated as a granule

Insects Controlled and Registered Crops:

CROP	INSECT
Barley, Oats, Wheat	Army, darksided, pale western and redbacked cutworms, armyworms, grasshoppers
Wheat only	Russian wheat aphid, brown wheat mite, wheat midge
Canola	Darksided, redbacked, variegated, pale western, and army cutworms; bertha army worm, alfalfa looper, armyworm, diamond-back moth larvae, grasshoppers, lygus bug
Flax	Darksided, redbacked, variegated, pale western, and army cutworms, armyworm, bertha armyworm
Lentils	Pale western cutworm, grasshoppers
Sunflowers	Redbacked, pale western and army cutworms, sunflower seed weevil (except for <i>Pyrinex</i> and <i>Citadel</i>)
Corn	Darksided, black and redbacked cutworms
Potato	Wireworms (in-furrow at planting - <i>Pyrinex</i> and <i>Pyrifos</i> only), Colorado potato beetle (larvae), potato flea beetle, tarnished plant bug, redbacked cutworm, black cutworm, darksided cutworm

Application:

Chlorpyrifos may be applied by air or ground equipment except for the following. Ground application only for redbacked cutworm control in corn and sunflower. Ground application only for potatoes.

Pyrifos 15G may be applied by ground only and is to be applied in furrow at planting. Refer to label for specific rates with respect to row spacing.

Uniform coverage of the crop is essential in aerial applications. Apply when insects exceed economic threshold levels and use sufficient water for good coverage. Use higher rates for heavy infestations, mature insects, heavy crop canopy, or under dry soil conditions.

How it Works:

Chlorpyrifos is a broad spectrum, non-systemic insecticide and works by contact, ingestion and vapour action (inhalation).

Effects of Weather:

Avoid application under hot temperatures. Best results will be obtained for wheat midge and cutworms when application is made in evening (after 7 p.m.) or morning (before 8 a.m.). DO NOT apply to plants under extreme drought stress or crop injury may occur.

Tank mixes:

Various chlorpyrifos labels differ. Contact the specific company for supported tank mixes.

Restrictions:

Grazing: Treated cereals grown for cover crop should not be used for human or animal consumption if treated within 60 days of harvest.

Storage: Combustible. DO NOT store near heat or flame. DO NOT store with food, feed, drugs or clothing.

Wheat, barley, oats, canola, corn, flax, lentil, sunflower, potatoes - DO NOT make more than 1 application per season.

Buffer zones around sensitive areas: For all aerial applications, a buffer zone of 100 metres is required for the protection of aquatic habitats.

Precautions:

Chlorpyrifos has a high acute mammalian toxicity. Very toxic to bees, fish, birds, aquatic organisms and other wildlife. May be fatal if swallowed. Causes substantial but temporary eye injury. Harmful if absorbed through skin. May cause skin or eye irritation. Wear protective clothing, impervious gloves and goggles. Wash thoroughly with soap and water after handling and applying. Immediately remove contaminated clothing and wash before re-use. DO NOT apply or allow to drift on to workers or other persons. DO NOT apply directly to water or where runoff could

occur to adjacent aquatic sites. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. DO NOT use on flowering crops or weeds. Avoid use when bees are actively foraging.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Clutch

Company:

Valent Canada Inc. – PCP#29382

Formulation:

50% clothianidin formulated as a water dispersible granule

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, aphids, leafhoppers

Application:

In furrow application: Apply as a narrow band in-furrow at planting. For best results, direct spray on the seed pieces or seed potatoes. Use sufficient water volume to ensure uniform coverage and optimal uptake. Use higher rate when extended control is needed. DO NOT apply *Clutch* more than once per season as an in furrow treatment.

Foliar application: May be applied by air or ground. Maximum of 3 foliar applications may be made per crop per season. Application intervals must be at least 10 days apart and must be rotated with an insecticide from a different chemical family. Use sufficient water volume to ensure uniform coverage. Use higher rate when insect populations are high.

Insecticide Group – 4A

(Refer to page 456)

How it Works:

Clothianidin is in the neonicotinoid class of insecticides and works by contact or ingestion, with systemic properties that provide residual control. Residual control will depend on environmental factors, plant growth, dosage rate and level of insect infestation.

Restrictions:

DO NOT follow a soil or in furrow application of *Clutch* with a foliar application of *Clutch* or any Group 4 or 4A insecticide.

DO NOT make a foliar application of *Clutch* following a seed piece treatment or in furrow application of *Clutch*, any product containing clothianidin or other neonicotinoid class (Group 4 or 4A) insecticides.

Re-cropping: Acceptable plant-back intervals for:

canola, corn, potato - no restrictions;
soybeans - 30 days.

Precautions:

Clothianidin is persistent and may carry over. It is recommended that any products containing clothianidin not be used in areas treated with this product during the previous season.

Avoid application when heavy rain is forecast.

DO NOT enter or allow entry into treated areas for 12 hours after application.

DO NOT graze treated fields or feed treated forage or hay to livestock.

Storage: DO NOT store in or around the home. Store unused product in a cool, ventilated, dry, secure area, away from food and feed.

DO NOT use treated seed pieces for food, feed or fodder.

Clothianidin is highly toxic to bees. DO NOT apply this product to flowering crops or weeds if bees are visiting the treatment area.

Clothianidin is toxic to beneficial insects, aquatic organisms, birds, small wild mammals and non-target terrestrial plants. Observe buffer zones for sensitive areas (e.g. aquatic habitats, forested areas) as specified on label directions.

DO NOT use this chemical where groundwater contamination can occur, especially in areas where soils are permeable (e.g. sandy) and / or the water table is shallow.

If this product is to be applied to a commodity destined for export to the United States, visit Crop Life Canada's website www.croplife.ca for information on acceptable residue limits.

Hazard Rating:



Caution – Poison

– Eye irritant

For an explanation of the symbols used here see page 11.

Concept

Company:

Bayer CropScience Inc. (PCP# 29611)

Formulation:

75 g/L imidacloprid and 10 g/L deltamethrin formulated as a suspension concentrate

Container size - 5.26 L jug

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, aphids, leafhoppers, potato flea beetle, tarnished plant bugs, European corn borer (suppression only)
Soybean	Soybean aphid

Application:

Ground application only.

Apply when target pest has reached economic threshold levels. Repeat application if pest populations reach economic thresholds.

Use sufficient water volumes for thorough coverage (i.e. minimum of 40 – 80 litres of water per acre)

For control of tarnished plant bug it is recommended to

Insecticide Group – 3A, 4A

(Refer to page 456)

use *Concept* insecticide only when timing of application coincides with the timing for another pest on the label for potatoes.

How it Works:

Concept insecticide works through contact and systemic activity. Insecticide components: Imidacloprid is a chloro-nicotinyl, systemic (within the plant) insecticide that works by contact or ingestion. Deltamethrin is a non-systemic pyrethroid insecticide that works through contact and ingestion.

Restrictions:

Allow a minimum of 5 days between applications.

DO NOT make more than 3 applications of *Concept* in a year.

DO NOT apply *Concept* through any type of irrigation equipment.

DO NOT apply *Concept* following a seed treatment or soil application of any Group 4 (neonicotinoid class) insecticide.

A buffer zone of 8 metres is required between the down-wind point of application and the closest edge of aquatic habitats.

Re-cropping: Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application.

Acceptable plant-back intervals for:

Cereal grains (wheat, barley, oats) – 30 days

Pea and bean (including fababean, soybean and dry common bean) – 9 months

All other food and feed crops – 12 months

Green manure and other cover crops not intended for human or animal consumption - no plant-back interval required following treatment.

DO NOT graze or harvest cover crops for food or feed.

Precautions:

DO NOT enter or allow entry into treated areas for a period of 24 hours after application of *Concept*.

Concept is highly toxic to fish and other aquatic organisms. DO NOT allow spray drift to come in contact with lakes, streams, rivers, ponds or other aquatic areas including marshes, ponds, ditches, streams, lakes, etc. DO NOT apply *Concept* where runoff is likely to occur and be hazardous to aquatic organisms in neighboring areas.

DO NOT apply *Concept* within 15 metres of well-heads or aquatic systems. DO NOT mix, load or clean equipment within 30 metres of well-heads or aquatic systems.

This product is highly toxic to bees exposed to direct treat-

ment or residues on flowering crops or weeds. DO NOT apply this product or allow it to drift to flowering crops or weeds if bees are visiting the treatment area.

If this product is to be applied to a commodity destined for export to the United States, visit Crop Life Canada's website www.croplife.ca for information on acceptable residue limits.

Storage: DO NOT use or store in or around the home. Store unused product away from feeds, seeds, fertilizer, plants and foodstuffs.

Concept cannot be stored below freezing.

If stored for one year or longer, shake well before using.

Hazard Rating:

Warning – Eye Irritant



deltamethrin: Danger – Poison



imidacloprid: Caution – Poison

For an explanation of the symbols used here see page 11.

Coragen

Insecticide Group – 28

(Refer to page 456)

Company:

E.I. duPont Canada (PCP#28982)

Formulation:

200 g/L chlorantraniliprole formulated as a suspension.

Container sizes - 3.79 L, 6 L

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa, sweet clover	Alfalfa weevil (suppression only)
Bean	Armyworms, cutworms, European corn borer
Canola, mustard	Bertha armyworm, diamondback moth, Cutworms, swede midge, imported cabbageworm
Chickpea, lentil, pea	Armyworms, cutworms
Corn	Armyworm, fall armyworm, black cutworm, variegated cutworm, corn earworm, European corn borer
Flax	Bertha armyworm, cutworms
Forage (grass) seed	Armyworms
Potato	Armyworm, Colorado potato beetle, European corn borer, variegated cutworm
Sunflowers	Cutworms, banded sunflower moth
Safflower	Cutworms

Application:

May be applied by air or ground equipment on potatoes.

Begin application when treatment thresholds have been reached. Thorough coverage is essential for optimal control.

Use the high rate under heavy pest pressure and/or when larger larvae are present.

Spray Volume for potatoes, corn, canola, mustard and flax: Apply in a minimum finished spray volume of 40 L/acre by ground. Apply in a minimum finished spray volume of 20 L/acre by air.

How it Works:

Coragen disrupts muscle activity in the insects, resulting in paralysis. Treated pests stop feeding quickly after ingestion, become lethargic and lose mobility.

Restrictions:

DO NOT make more than 4 applications per season on alfalfa (seed production), bean, chickpea, lentil, pea, potatoes, corn, and forage seed.

Canola, flax, mustard - DO NOT make more than 3 applications per season.

DO NOT apply more than once every 5 days.

DO NOT make more than 1 application per cutting on alfalfa and sweet clover. DO NOT exceed a total of 455 ml of Coragen per acre per season.

Potatoes, bean, chickpea, lentil, pea - DO NOT apply more than once every 3 days.

Corn - DO NOT apply more than once every 7 days.

Canola, rape seed, mustard, flax - DO NOT make more than 3 applications per season. DO NOT apply more than once every 5 days.

Forage (grass) seed - Seed production only. DO NOT feed forage, fodder or hay to livestock.

Re-Entry Interval: 12 hours.

Storage: Store product in original container only, away from other pesticides, fertilizer, food or feed. Not for use or storage in or around the home. Keep container closed.

Precautions:

Toxic to aquatic organisms. Observe buffer zones specified on the label.

The use of this chemical may result in contamination of groundwater, particularly in areas where soil is permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:

Very low toxicity to mammals.

Keep out of reach of children.

Cypermethrin

Insecticide Group – 3A

(Refer to page 456)

Company:

Engage Agro corporation (*Ripcord* Insecticide – PCP#30316)
United Phosphorous, Inc. (*UP-Cycle* 2.5 EC – PCP#28795)

Formulation:

cypermethrin formulated as an emulsifiable concentrate.
(*Ripcord* Insecticide – 407 g/L; *UP-Cycle* 2.5 EC – 250 g/L)
Container size – 1L

Insects Controlled and Registered Crops:

CROP	INSECT
Wheat, barley (<i>Ripcord</i> only)	Grasshoppers, cutworms
Canola, rapeseed, mustard	Grasshopper (<i>Ripcord</i> only), flea beetle, bertha armyworm
Roadsides, headlands, summerfallow	Grasshopper (<i>Ripcord</i> only)
Sunflower	Cutworm (<i>Ripcord</i> only), sun- flower beetle, sunflower seed weevil
Corn	European corn borer, cutworms (<i>Ripcord</i> only), corn earworm (<i>Ripcord</i> only)
Potato	Colorado potato beetle, flea bee- tle, leafhopper, tarnished plant bug, cutworms (<i>Ripcord</i> only)

Application:

Ripcord may be applied by ground application only for control of grasshoppers on wheat, barley, roadsides, headlands and canola; for flea beetle control on canola; and for control of cutworms.

Ripcord may be applied by ground or air for bertha armyworm, sunflower beetle, sunflower seed weevil, corn earworm, European corn borer and Colorado potato beetle, flea beetle, leafhoppers and tarnished plant bug on potatoes. Only 1 aerial application is permitted per season for bertha armyworm on canola and sunflower beetle and seed weevil on sunflower, and 2 applications per season for corn and potatoes.

UP-Cycle may be applied by ground or air in corn, potatoes, canola, rapeseed, sunflower

Observe a 16 yard (15 m) buffer zone from water bodies and other sensitive areas when applying cypermethrin with ground application. Observe a 110 yard (100 m) buffer zone from water bodies and sensitive areas when applying by air.

Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage. Use higher rates for mature insect stages (grasshoppers) or severe infestations.

How it Works:

Ripcord and *UP-Cycle* are pyrethroid insecticides that work as a contact and stomach poison.

Effects of Weather:

Activity of *Ripcord* on grasshoppers is reduced as soil temperature increases. Application for grasshopper control should be made at temperatures below 25°C. Spraying for grasshoppers should be delayed until evening if daytime temperatures are above 25°C.

Avoid application of *UP-Cycle* when temperatures exceed 27°C

Restrictions:

Grazing: Treated crops must not be grazed or cut for hay.

Storage: Keep in original container during storage. DO NOT contaminate or store near foodstuffs.

Precautions:

Cypermethrin is of low to moderate acute mammalian toxicity and very toxic to bees, fish, and aquatic organisms. Harmful or fatal if swallowed. May be harmful if absorbed through skin.

Severely irritating to eyes. Causes skin irritation and sensitization. Wear longsleeved protective clothing and gloves when handling or applying. Wear face shield or goggles when mixing. DO NOT apply where streams, lakes and ponds may be contaminated. Avoid spraying when bees are foraging.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Decis

Insecticide Group – 3A

(Refer to page 456)

Company:

Bayer CropScience (PCP#17734)

Formulations:

50 g/L deltamethrin formulated as an emulsifiable concentrate.

Container sizes: *Decis* 5FC - 2.4 L and 9.6 L jugs

Insects Controlled and Registered Crops:

CROPS	INSECT
Alfalfa (seed crops only)	Lygus bug, alfalfa weevil
Field corn	European corn borer
Potato	Colorado potato beetle, potato flea beetle, Lygus bugs, leafhopper
Canola, rapeseed, mustard (condiment and oilseed quality <i>Brassica juncea</i> varieties)	Flea beetles, clover cutworm, bertha armyworm, diamondback moth, beet webworm, cabbage seed-pod weevil (adults only), Lygus bugs, grasshoppers (canola only)
Sunflower	Sunflower beetle
Wheat, barley, oats, lentils	Cutworms, grasshoppers
Rangeland, pastures, roadside, fence row	Grasshoppers
Flax	Cutworms, beet webworm, grasshoppers
Red clover (seed production only)	Lesser clover leaf weevil (suppression only)

Application:

Decis may be applied by air or ground equipment to all crops with the exception of alfalfa, red clover and corn, which require ground application only. Apply when insects exceed economic threshold numbers with sufficient water for good coverage. Use higher rates for severe infestations, on dense foliage or when a number of insect growth stages are present.

Alfalfa (seed production) - Use higher rates if alfalfa weevil present.

Tank Mixes:

When in a tank-mix the spray mixture must be constantly agitated throughout application. Do not allow the spray mixture to stand in the spray tank for more than 4 hours after mixing.

Decis may be tank mixed with the following herbicides: *Pardner*, *Buctril M*, *Banvel*, *MCPA*, *2,4-D*, *Puma¹²⁰ Super*. Tank mixes with *Puma¹²⁰ Super* or *Puma¹²⁰ Super plus Buctril M* are for use in spring and durum wheat only.

Bayer CropScience also supports the following mixes that are not on the *Decis* label. Apply mixes according to the most restrictive use limitations for either product:

Herbicides – Glyphosate, *Odyssey* and *Solo*

Fungicides – *Headline*, *Lance*, *Tilt*

When a tank mix is used the labels of the tank mix partners are to be consulted.

How it Works:

Decis is a non-systemic, synthetic pyrethroid which works by contact and ingestion.

Effects of Weather:

DO NOT spray under a strong temperature inversion, or when temperature exceeds 25°C as this will result in a reduction in control. Best control will be achieved when *Decis* is applied during cooler periods of the day. DO NOT apply within 1 hour of rain.

Restrictions:

Alfalfa seed production – DO NOT apply more than once per year.

Canola – May be applied by ground or air. DO NOT apply more than once per season.

Corn – DO NOT apply more than 3 times per year

Potato – (Ground) DO NOT apply more than 3 times per year. (Aerial) DO NOT apply more than 2 times per year. May be used only once per season on high organic (muck) soils.

Red clover – DO NOT apply by air. DO NOT make more than 2 applications per year. DO NOT use treated crop for feed or forage. Restricted entry interval – 12 hours

Wheat, barley, oats, flax, lentil – (Ground) DO NOT apply more than 3 times per year. (Aerial) DO NOT apply more than 2 times per year.

Storage: DO NOT store below freezing. DO NOT store near feed, food, seeds or fertilizer. Keep away from heat, sparks and open flames. If stored for 1 year or longer, shake well before using.

Others: Ground application - Observe a 16 yard (15 m) buffer zone from environmentally sensitive areas (for example, wetlands, sloughs, rivers, houses, farm buildings).

Aerial application - Leave a 110 yard (100m) buffer zone. DO NOT apply *Decis* by air when the wind speed exceeds 8 kph. In soils with high organic content (muck soils), *Decis* 5 EC should be applied only once during each crop year, prior to August 1, and at rates of no more than 80 mL/acre.

Precautions:

Decis is of high mammalian toxicity and is a severe eye and skin irritant. Avoid contacting or breathing spray mist.

This product contains petroleum distillates which are moderately to highly toxic to aquatic organisms and fish. *Decis* is toxic to bees and other beneficial insects. Avoid spraying when bees are foraging. Wear protective clothing, including goggles and respirator, when handling or spraying. DO NOT contaminate or store near feed or foodstuffs. Wash thoroughly after using *Decis*.

Hazard Rating:



Danger - Poison

For an explanation of the symbols used here see page 11.

Delegate

Insecticide Group - 5
(Refer to page 456)

Company:

Dow AgroSciences (PCP# 28778)

Formulations:

25% spinetoram formulated as wettable granules.

Container size - 840 g

Insects Controlled and Registered Crops:

CROP	RATE (g per acre)	INSECT
Potato*	65 to 97	Colorado potato beetle (time for egg hatch or small larvae)
	65	European corn borer (time to coincide with peak egg hatch)
Wheat, barley, oats, rye**	40 to 81	Armyworm (when economic thresholds dictate)

* Maximum 3 applications per year with a minimum retreatment interval of 7 days

** Maximum 3 applications per year with a minimum retreatment interval of 5 days

Application:

Apply by ground only in sufficient water volume to cover the entire plant using a combination of nozzles and pressure designed to deliver thorough, even coverage

with ASABE fine classification droplets. DO NOT apply through irrigation systems.

How it Works:

Delegate is derived from the fermentation of the bacterium *Saccharopolyspora spinosa*, which is then chemically modified to create the active ingredient. Spinetoram affects the insect nervous system. It does not interact with the known binding sites of other classes of insecticides. It works through ingestion or contact with the target insects. Target insects cease feeding within a few minutes, although death may take a few days.

Tank Mixes:

DO NOT mix this product with any other pesticide or fertilizer.

Restrictions:

Re-entry: DO NOT enter treated field for 12 hours.

Preharvest: DO NOT harvest within 21 days of application for wheat (spring and durum) or within 7 days for potato.

Grazing: No restrictions indicated.

Aerial Application: DO NOT apply by air.

Storage: Store in a cool, dry place. Keep from freezing.

Precautions: *Delegate* is toxic to bees, small wild mammals, aquatic organisms and non-target terrestrial plants, and may be toxic to certain beneficial insects exposed to direct treatment, drift or residues on flowering crops or weeds. DO NOT apply this product to flowering crops or weeds if bees are visiting the treatment area. Minimize spray drift to reduce harmful effects on bees, and other beneficial insects or small mammals in habitats next to the application site such as hedgerows and woodlands.

Buffer Zones:

Application method	Buffer Zones (metres†) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground *	10	5	1

See page 29 for an explanation of the different habitats.

* Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

† Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Tank Cleaning:

Refer to page 15.

Hazard Rating:

No specific hazard rating specified.

For an explanation of the symbols used here see page 11.

Dibrom

Insecticide Group – 1B
(Refer to page 456)

Company:

United Agri Products (PCP#7442)

Formulations:

864 g/L naled formulated as an emulsifiable concentrate.

Container size - 4 x 3.78 L jugs per case and 2 x 9.46 L jugs per case

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa, clover, vetch	Aphids, loopers, leafhoppers, Lygus bugs
Beans	Alfalfa looper, aphids
Potatoes	Flea beetles, Colorado potato beetles, leafhoppers
Rangeland, field areas and pastures.	Grasshoppers

Application:

Apply *Dibrom* by ground or air. Use designated amounts in full volumes of water. For ground application use 40-120 L of water per acre. For aerial use 4.4-8.8 L of water per acre unless otherwise stated.

How it Works:

Dibrom is an organophosphate insecticide. It acts as a contact and stomach poison.

Effects of Weather:

DO NOT apply *Dibrom* when air temperature is greater than 32°C.

Restrictions:

Environment: DO NOT contaminate any body of water, waterway or water source. *Dibrom* is moderately to highly toxic to birds, aquatic animals and other wildlife.

Re-entry interval: DO NOT enter or allow worker re-entry into treated area for 48 hours following application.

Precautions:

Concentrate may cause skin damage. DO NOT get on skin, eyes or clothing. Use waterproof gloves and face shield or goggles when handling concentrate. Harmful if swallowed. Avoid breathing spray mist.

Avoid contamination of feed, foodstuffs and drinking water.

Highly toxic to bees. Avoid application during periods of bee activity.

Hazard Rating:

Danger – Poison

For an explanation of the symbols used here see page 11.

Dimethoate

Insecticide Group – 1B

(Refer to page 456)

Company:

IPCO (*Cygon 480 EC* – PCP#9807)

Cheminova (*Cygon 480-Ag* – PCP#25651)

United Agri Products (*Lagon 480E* – PCP#9382)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Formulations:

Cygon/Lagon - 480 g/L dimethoate formulated as an emulsifiable concentrate. Container size - 10 L.

Insects Controlled and Registered Crops:

CROP	<i>Cygon 480-Ag</i> INSECT	<i>Lagon 480E</i> INSECT	<i>Cygon 480 EC</i> INSECT
Peas		Aphids	Aphids
Potatoes (ground application only)	Aphids, leafhoppers	Aphids, leafhoppers, Lygus bugs	Aphids leafhoppers, Lygus bugs
Alfalfa *(rates vary for seed and forage production)	Aphids, leafhoppers, Lygus bugs*, plant bugs*, alfalfa blotch leaf miner, grasshoppers, reduction of alfalfa weevil larvae	Aphids, leafhoppers, alfalfa blotch leafminers, grasshoppers, reduction of alfalfa weevil larvae, Lygus bugs*, plant bugs*	Aphids, blotch leafminer, grasshoppers, leafhoppers, Lygus bugs*, plant bugs*, sweet clover weevil, reduction of alfalfa weevil larvae
Canaryseed	Aphids	Aphids	Aphids
Canola/ rapeseed	Aphids, leafhoppers, grasshoppers	Aphids, leafhoppers, grasshoppers	Aphids, leafhoppers, grasshoppers
Forage crops	Lygus bugs, plant bugs, grasshoppers	Grasshoppers, aphids (suppression only of Russian wheat aphid)	Aphids, grasshoppers, leafhoppers, Lygus bugs, plant bugs, reduction of alfalfa weevil larvae
Sweet clover, red clover, alsike clover	Sweet clover weevil	Sweet clover weevil	Sweet clover weevil
Pastures, waste areas	Grasshoppers	Grasshoppers	Grasshoppers
Wheat	Wheat midge, thrips, aphids (suppression only of Russian wheat aphid)	Thrips, grasshoppers, wheat midge, Russian wheat aphid	Wheat midge, aphids, thrips, grasshoppers
Barley, oats	Thrips, aphids (suppression only of Russian wheat aphid)	Thrips, grasshoppers,	Aphids, thrips, grasshoppers
Flax	Aphids	Aphids	Aphids

CROP	<i>Cygon 480-Ag</i> INSECT	<i>Lagon 480E</i> INSECT	<i>Cygon 480 EC</i> INSECT
Rye			Grasshoppers
Soybeans		Aphids, leafhoppers, Lygus bugs, spider mites	Aphids, leafhoppers, Lygus bugs, spider mites

Application:

May be applied by air or ground equipment (unless otherwise specified above). Apply when insects exceed economic threshold numbers and use sufficient water for good coverage. Use higher rates for adult insects, heavy infestations or dense canopy.

How it Works:

Dimethoate is a broad spectrum, systemic (within the plant) and contact, organophosphate insecticide and acaricide.

Restrictions:

Grazing: Remove cattle prior to spraying. Read label carefully to determine livestock re-entry period.

Storage: Store at temperatures between 4°C and 30°C and in areas away from feed and food.

Others: DO NOT treat when bees are foraging. DO NOT apply to alfalfa in bloom. Read label carefully to determine maximum number of applications per season. Wait at least 10 days before placing leafcutter bees in treated fields.

Precautions:

Dimethoate is of high acute mammalian toxicity and is highly toxic to birds, bees and other animals. Wear a respirator, goggles, rubber gloves, rubber boots and coveralls when handling concentrate. Avoid contact with skin and eyes. DO NOT inhale spray mist.

Hazard Rating:



Warning Poison – *Lagon, Cygon 480 AG*



Danger Poison – *Cygon 480 LC*

For an explanation of the symbols used here see page 11.

Dipel 2X DF

Insecticide Group – 11
(Refer to page 456)

Company:

Valent BioSciences (PCP#26508)

Formulation:

Bacillus thuringiensis var. *Kurstaki* strain ABTS-351 fermentation solids, spores and insecticidal toxins - 57.0%

Potency: 32,000 Cabbage Looper Units (CLU) per mg (32 billion CLU per Kg)

Insects Controlled and Registered Crops:

CROP	INSECT
Sunflower	Sunflower moth
Timothy	Essex (European) skipper
Corn	European corn borer larvae
Potato	Cabbage loopers

Application:

Treat when larvae are young (early instars) before the crop is damaged. A spreader sticker such as *Triton B1956* should be used to give thorough foliage coverage.

How it Works:

Dipel is a biological stomach insecticide resulting in the larvae ceasing to eat in a few hours, with death usually occurring within 1-3 days.

Restrictions:

Storage: Store at temperatures between 0° and 25°C (cooler temperatures preferable).

Others: DO NOT allow dilute spray to stand in tank for more than 12 hours. Use product within 24 months of date of manufacture if stored at cool temperatures. Final spray solution for *Dipel* should have a pH of 5-7.

Precautions:

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. In case of contact with eyes or skin, immediately flush eyes or skin with plenty of water.

Hazard Rating:

Warning contains the allergen soy



Caution – eye irritant, skin irritant, potential sensitizer

Eco Bran

Insecticide Group – 1A

(Refer to page 456)

Company:

Peacock Industries (PCP#25815)

Formulation:

Wheat bran infused with carbaryl insecticide.

Container size - 20 kg bag; 1kg bottle

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa, beans, clover, corn, oats, rye, wheat, barley, canola, pastures, rangelands, forage grasses, field borders, headlands, rights-of-way, roadsides, wastelands	Grasshoppers

Preharvest Intervals and Livestock Re-entry Periods:

Crop	Preharvest Interval/ Livestock re-entry period
Corn	1
Alfalfa, clover	2
Beans	5
Oats, rye, wheat	14
Barley	28
Canola	Treat only seedlings
Field borders, headlands, rights-of-way, roadsides, wastelands	0
Entry of beef cattle or other livestock to pastures, rangelands or forage grasses	1
Entry of dairy cattle to pastures or rangelands, harvest of forage crops	2

Application:

For ground application only. DO NOT apply by air.

Broadcast application may be made with spreaders, hand applicators, or by hand. Use gloves and wash thoroughly following application. More information on application and applicators can be found at: <http://www.grasshoppercontrol.com>

Restrictions:

DO NOT apply within 50m of sloughs, ponds, streams, dugouts or open water. Apply when winds are between 3-8 kph and do not favour drift.

Presence of product on flowering crops such as alfalfa and clover will not harm foraging honey or leafcutter bees.

May be used in pastures while beef cattle are grazing.

Precautions:

Harmful if inhaled or swallowed. Avoid breathing dust or vapour from bait. Use only in well-ventilated areas. May cause eye irritation. Avoid contact with eyes and skin. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Keep away from heat, sparks and open flame.

Endosulfan

Insecticide Group – 2A

(Refer to page 456)

Company:

Bayer CropScience (*Thiodan 4EC* – PCP#15747)
United Agri Products (*Thionex 50WP* – PCP#15333;
Thionex EC – PCP#23453)

Different trade names refer to different companies. Note that products may have different label recommendations. Check your label for more information.

Last date of sale for Endosulfan products is December 31, 2015.

Last date of use for Endosulfan products is December 31, 2016.

Formulation:

Thiodan / Thionex - 400 g/L endosulfan formulated as an emulsifiable concentrate. Container size - 10 L

Thionex 50WP - 50% endosulfan formulated as a wettable powder. Container size - 1 kg.

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Aphid, Colorado potato beetle, flea beetle, leaf-hopper

Application:

Ground application only. Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage.

How it Works:

Endosulfan is an organochlorine insecticide that works as a contact and stomach poison.

Tank Mixes:

Compatible with most commonly used insecticides and fungicides, except Bordeaux mixture and hydrated lime.

Restrictions:

Grazing: DO NOT feed treated crop residue to livestock.

Storage: DO NOT store below freezing. If stored one year or longer, shake well before using.

Others: On sunflowers, DO NOT make more than 1 application per season.

Precautions:

Endosulfan is of high, acute mammalian toxicity and is toxic to bees. Highly toxic to fish. Hazardous if swallowed, inhaled or absorbed through the skin. Wear synthetic rubber gloves and approved respirator in prolonged spray-mixing and loading operations. DO NOT apply or allow spray drift to areas occupied by unprotected persons and animals or onto streams, lakes or ponds.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Entrust 80 W

Insecticide Group – 5

(Refer to page 456)

Company:

Dow AgroSciences (PCP#27825)

Formulation:

80% spinosad

Container size - 4 x 113.4 g packets

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle larvae and European corn borer larvae

Application:

Apply *Entrust 80 W* as a foliar spray by ground only. Apply when scouting indicates the target pest species have reached economic threshold levels. For Colorado potato beetle larvae, target eggs at hatch or small larvae. For control of European corn borer, time the application to coincide with peak egg hatch. Use higher application rate for higher pest pressure or when extended egg hatch is anticipated. If pest populations persist, a repeat application 7 to 10 days after the initial application may be necessary.

How it works:

Entrust 80 W is in the spinosine class of insecticides. It is a contact and stomach insecticide. It is derived from the fermentation of *Saccharopolyspora spinosa*.

Effects of weather:

This product has the potential for run-off. Do not spray immediately after a rainfall or if rain is forecast within 48 hours after application.

Restrictions:

Storage: Avoid freezing. DO NOT store or ship with food, feeds, drugs or clothing.

Others: DO NOT make more than 2 applications per season (maximum of 60 g/acre).

Precautions:

Entrust 80 W is highly toxic to honey bees exposed to direct spray. Avoid spraying on crops in bloom or other vegetation when bees are actively foraging. This product is toxic to other beneficial parasites and predators. DO NOT contaminate aquatic habitats such as ponds, lakes, rivers, streams and wetlands during application or when rinsing equipment and containers.

Buffer Zones: A buffer zone of 2 metres (early season) or 1 metre (late season) is required between downwind edge of spray boom and sensitive aquatic habitats.

Avoid contact with eyes, skin, and clothing.

DO NOT enter or allow worker entry into treated areas for a period of 12 hours after application.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Fulfill

Insecticide Group – 9B
(Refer to page 456)

Company:

Syngenta Canada (PCP # 27274)

Formulation:

50% pymetrozine formulated as a wettable granule

Container size – 6 X 780 g

Insects Controlled and Registered Crop:

CROP	INSECT
Potatoes	Aphids including: green peach, potato, foxglove, buckthorn

Application:

May be applied by ground or air. Apply *Fulfill* to plant foliage. Thorough spray coverage is essential for best performance. Apply *Fulfill* with sufficient water (minimum of 40 L/acre) to ensure good coverage of all plant surfaces. Higher water volumes will generally result in better coverage, especially under adverse conditions (hot, dry), where a dense plant canopy exists and/or aphid infestations are high. One additional application may be needed to control persistent aphid populations. Allow a minimum of 7 days between applications.

DO NOT apply *Fulfill* insecticide through chemigation.

DO NOT use in nurseries or in plant propagation houses, or on any plants grown for use as transplants.

The use of a non-ionic adjuvant is recommended to improve the performance of *Fulfill* insecticide under drought stress conditions.

How it Works:

Fulfill is a systemic insecticide and works primarily by ingestion and but has some contact activity. Affected aphids stop feeding shortly after exposure, but may remain on the plant foliage until they die, which is usually within 2-4 days. *Fulfill* insecticide has residual activity in the plant and will control aphids that move onto the plant after spraying.

Fulfill has shown no phytotoxicity on the varieties of potato tested when applied at the label rates.

Effects of Weather:

Fulfill insecticide exhibits movement through the leaf surface into plant tissue and is rainfast as soon as the spray solution has dried.

Restrictions:

Storage: Store in a cool, dry, place away from food, drinks, and animal feeding stuffs. Keep in the original container tightly closed.

Others: DO NOT apply by air. DO NOT exceed 2 applications (152 g product/acre) per crop per season. DO NOT apply directly to aquatic systems, permanent water bodies or areas where surface water is present. DO NOT contaminate water when cleaning equipment or disposing of equipment wash water.

A re-cropping restriction of 30 days is required for all crops.

Precautions:

May cause skin sensitization reactions. Applicators and other handlers must wear personal protective equipment including, long-sleeved shirt, long pants, waterproof gloves and shoes plus socks. DO NOT enter or allow entry into treated areas for 12 hours. DO NOT use, pour, spill, or store near heat or open flame.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Imidacloprid

Insecticide Group – 4A
(Refer to page 456)

Company:

Bayer CropScience (*Admire* 240 – PCP# 24094;
Admire SPT – PCP# 27702)
Chemnova (*Grapple* – PCP# 28726 / *Grapple₂* – PCP# 29048)
MANA Canada (*Alias* 240 SC – PCP# 28475)

Formulation:

240 g/L imidacloprid formulated as a suspension concentrate.

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, aphids, potato leafhopper, potato flea beetle

Application:

Soil application: (*Admire* 240 / *Alias* 240 SC / *Grapple* / *Grapple₂*) Apply as a narrow band in-furrow. For best results, direct spray on the seed pieces in the furrow. Scout potato fields frequently, especially during warmer part of growing season. If pest populations exceed economic thresholds apply a recommended foliar insecticide with a different mode of action than imidacloprid.

Seed piece treatment: (*Admire SPT* / *Admire* 240 / *Alias* 240 SC / *Grapple* / *Grapple₂*) Refer to Imidacloprid in the seed treatments product pages.

Foliar application: (*Admire* 240 / *Alias* 240 SC / *Grapple* / *Grapple₂*) Apply only if insect populations exceed recommended economic thresholds. For optimum control, good coverage of the foliage is needed. A maximum of 2 foliar applications may be made per crop per season. Scout fields and retreat if needed. For aphids, two applications at least seven days apart may be required to achieve control. DO NOT make a foliar application following a soil or seed treatment of the product in the same crop. Allow at least 7 days after the last application and before harvesting the crop.

How it Works:

Imidacloprid is a chloronicotinyl, systemic (within the plant) insecticide that works by contact or ingestion. Control period may vary due to climate and soil conditions

Restrictions:

DO NOT apply more than once per season as a soil application.
DO NOT follow a soil application with a foliar application.

Re-cropping: Acceptable plant-back intervals for:

Cereal grains (wheat, barley, oats) - minimum 30 days

Peas and beans - 9 months

All other food and feed crops - 12 months

Green manure and other cover crops can be grown without plant-back intervals but cannot be grazed or harvested for food or feed.

DO NOT apply in fields where imidacloprid has been used during the previous season.

DO NOT apply through any irrigation system.

Precautions:

DO NOT re-enter treated areas for 24 hours after foliar application.

Avoid application when heavy rain is forecast.

DO NOT apply product or plant treated seed pieces within 15 metres of well-heads or aquatic systems, including marshes, ponds, ditches, reservoirs, streams, lakes, etc.

DO NOT mix, load or clean spray equipment within 30 metres of well-heads or freshwater habitats.

For application with air-blast equipment, DO NOT apply within 40 metres of well-heads or aquatic systems.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and / or where the water table is shallow.

Storage: DO NOT store in or around the home. Store unused product in a cool, ventilated, dry, locked area and avoid cross-contamination with other pesticides, fertilizers, food and feed.


DO NOT use treated seed pieces for food, feed or fodder.

Imidacloprid is highly toxic to bees exposed to direct treatment or residues on flowering crops or weeds.

Highly toxic to aquatic invertebrates and birds.

If this product is to be applied to a product destined for export to the United States, contact 1-866-375-4648 or www.croplife.ca.

Hazard Rating:

 Caution – Poison

For an explanation of the symbols used here see page 11.

Imidan

Insecticide Group – 1B
(Refer to page 456)

Company:

Gowan Canada (PCP#23006)

Formulation:

50% phosmet formulated as a wettable powder in water soluble sachets.

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa	Alfalfa weevil, alfalfa blotch leafminer
Potato	Colorado potato beetle, potato flea beetle, potato leafhopper, potato aphid

Application:

Apply by ground only.

Imidan 50-WP instapak is packaged in water soluble sachets that are to be dropped into the spray tank unopened. DO NOT use in low-volume, gear-type spray equipment.

How it Works:

Imidan is an organophosphate insecticide.

Restrictions:

Storage: Keep sachets dry and DO NOT allow sachets to contact any moist surface prior to adding to spray tank. Keep water soluble sachets in the protective container and store in a cool, dry place. DO NOT store at temperatures above 40°C.

Precautions:

Harmful if swallowed, inhaled or absorbed through the skin. Wear protective clothing, including rubber gloves and goggles, during mixing, loading and spraying.

Imidan is toxic to fish and bees. Keep away from any body of water.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Lambda-cyhalothrin

Insecticide Group – 3A
(Refer to page 456)

Company:

Syngenta Canada. (*Matador* – PCP# 24984)
MANA Canada (*Silencer 120 EC* – PCP# 29052)

Formulations:

120g/L lambda-cyhalothrin formulated as an emulsifiable concentrate Container size - 4 x 3.78 L.

Insects Controlled and Registered Crops:

CROP	INSECT
Potatoes	Colorado potato beetle, European corn borer, Lygus bugs, potato flea beetle, potato leafhopper, tuber flea beetle, armyworm
Canola, mustard	Crucifer flea beetle, grasshoppers, Lygus bug, cabbage seedpod weevil (adults), cabbage looper, diamond-back moth larvae, imported cabbageworm, bertha armyworm, swede midge
Sunflower	Sunflower beetle
Wheat, barley, oats	Grasshoppers, armyworm
Alfalfa, unimproved pasture	Grasshoppers
Summerfallow (Matador only)	Grasshoppers
Flax	Grasshoppers
Alfalfa Matador – Ground or Air Silencer – Ground only	Alfalfa weevil, Lygus bugs, pea aphid, potato leafhopper
Corn	European corn borer, corn earworm, cutworms, fall armyworm, armyworm
Beans	Cutworms, corn borer, potato leafhopper, Lygus bugs

CROP	INSECT
Chickpeas	Grasshoppers, potato leafhopper, cutworms
Fababeans (broad beans)	Lygus bugs, potato leafhopper, pea aphid
Lentils	Cutworms, grasshoppers, Lygus bugs, pea aphids, potato leafhopper
Peas	Cutworms, grasshoppers, pea aphids, pea leaf weevil
Soybeans	Cutworms, grasshoppers, Lygus bugs, soybean aphids
Timothy	Grasshoppers

Application:

Aerial:

Matador and Silencer: Canola, mustard, sunflower, flax, alfalfa, unimproved pasture - DO NOT make more than 1 application at the 33.2 ml/acre rate per year.

Corn, wheat, barley, oats, potatoes, soybean, dry edible bean, pea, chickpea, lentil, favabeans - DO NOT make more than 2 applications at the 33.2 ml/ac rate per year.

Matador: Summerfallow - DO NOT make more than 1 application at the 33.2 ml/acre rate per year.

Ground:

Canola, mustard, sunflower, flax, alfalfa, unimproved pasture, summerfallow (Matador only), corn, wheat, barley, oats - DO NOT make more than 3 applications per year at the 33.2 ml/acre rate.

Potatoes - DO NOT make more than 3 applications per year at the 33.2 ml/acre rate. DO NOT make more than 2 applications per year if using the 50 ml/acre rate. DO NOT exceed 100 ml/acre of lambda-cyhalothrin per year.

Beans, chickpeas, favabeans, lentils, peas, soybeans - DO NOT make more than 3 applications per year. DO NOT graze or harvest treated forage straw or hay for livestock feed.

Timing:

For potato insects, timing of application should be based on the presence of vulnerable pest developmental stages and significant populations as determined by local monitoring.

For sunflower beetles, use the high rate to control adults.

For flea beetles, to prevent migration of over-wintering adults throughout the field, spray a 15 m strip around the field at the first sign of flea beetle feeding.

For grasshoppers, apply the low rate when grasshoppers are up to the 3rd nymphal stage (up to 1 cm in length) or when insect numbers are low. Apply the high rate when insects are larger, up to but not including, winged adults or when insect numbers are high.

For corn borer control apply Matador before the larva bores into the plant stalk or pods

How it Works:

Lambda-cyhalothrin is a synthetic pyrethroid insecticide. It is a fast acting stomach and contact insecticide effective against a broad spectrum of foliar pests. It has no fumigant or systemic activity.

Effects of Weather:

For best results, apply Lambda-cyhalothrin during the early morning before temperatures rise, and during the evening, past the heat of the day.

Tank Mixes:

Herbicides: (Ground only)

Horizon

Tralkoxydim ^Δ

Fungicides: (Tank mixes on legumes (bean, chickpea, lentil,

pea, soybean) may be applied by ground only) Refer to label for other crops.

Propiconazole ^Δ

Allegro in dry bean

Quadris (Matador only)

Quilt (Matador only)

Headline (Silencer only) on dry field pea- to control insects and diseases listed on the label of each product. Read carefully and follow all use directions and use precautions on both the *Silencer* 120 EC and *Headline* EC Fungicide labels. Failure to follow the rates of use and timing of application as recommended for each product will result in unsatisfactory control of target pest.

Touchdown Total and Truxion

^Δ Manufacturers may only support specific mixes. Contact the manufacturer for more information.

Restrictions:

Grazing: DO NOT graze or feed lactating dairy animals on treated green cereal forage or treated pasture. DO NOT apply within 14 days of livestock foraging.

Alfalfa seed from treated crops is not to be used for production of "alfalfa sprouts" for human consumption.

Storage: Store above 0 degrees C. Storage below 0 degrees C will not impair the effectiveness of Lambda-cyhalothrin. However, following such storage, agitate well before use.

Others: Allow a 7-day interval between applications. DO NOT apply within 15m of productive fisheries, water or waterfowl habitat.

Buffer Zones:

Application method	Crop	Buffer Zones (metres [†]) Required for the Protection of:	
		Aquatic Habitats of Depths	
		Less than 1 m	Greater than 1 m
Ground	All field crops	15	15
Fixed wing airplane	Potatoes, oilseed crops, cereal crops, alfalfa, unimproved pasture, summerfallow	100	20
	corn	225	20
	Legume vegetables	600	25

See page 29 for an explanation of the different habitats.

- Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.
- For tank mixes, consult the labels of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Precautions:

Lambda-cyhalothrin has potential for skin and eye irritation. Avoid splashing in eyes or on skin, particularly the face. If hands are contaminated, wash with soap and water before touching other areas of skin. This product is very toxic to bees. Avoid spraying when bees are foraging. Spray deposits should be dry before bees commence foraging in treated crops.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Lannate

Insecticide Group – 1A
(Refer to page 456)

Company:

E.I. duPont Canada (PCP#10868)

Formulations:

90% methomyl formulated as a water soluble powder.
Container size - 24 x 225 gram water soluble bags.

Insects Controlled and Registered Crops:

CROP	INSECT
Canola	Alfalfa looper, bertha armyworm, clover cutworm, beet webworm
Flax	Bertha armyworm
Peas	Alfalfa looper, pea aphid
Wheat, oats, barley	Armyworm, thrips
Potato	Aphids, leafhoppers, flea beetles, variegated cutworm

Application:

May be applied to canola, flax and cereals by air or ground equipment. Ground applications only to peas and potatoes. Apply when insects exceed threshold levels using sufficient water for good coverage.

Suggested water volumes for ground application:

Potatoes - 100 to 340 L per acre

Canola and flax - 40 L per acre

Wheat, oats, barley, peas - 40 to 140 L per acre

When applied by air, pilot should not assist in mixing and loading operations. Apply a minimum of 9 L of water per acre) for aerial application.

Use higher rates for mature insects, dense canopy or when infestations are heavy.

Apply at 5 to 7 day intervals as necessary.

Early morning or late evening sprays are recommended.

How it Works:

Lannate is a carbamate insecticide that works by contact and ingestion and has some systemic action. Rapidly degraded in green, growing plants; short term residual.

Restrictions:

Storage: DO NOT store below 0°C. Store in original container away from other pesticides, fertilizer, food or feed.

Others: DO NOT handle water soluble bags with bare hands. Sprays or drift must not contact workers, other persons or animals. The area being treated must be vacated by unprotected persons.

Precautions:

Lannate is of high acute mammalian toxicity. May be fatal if swallowed, inhaled or absorbed through the eyes. DO NOT breathe dust or spray mist. DO NOT get in eyes, on skin or on clothing.

Toxic to fish, birds and other wildlife. Keep out of any body of water. DO NOT apply where run-off is likely to occur. DO NOT allow to drift from treated areas. Highly toxic to bees exposed to direct application. DO NOT apply to areas being visited by bees. Time applications to coincide with minimum bee activity.

When mixing, loading or applying *Lannate*, wear protective clothing, goggles and an approved respirator. Wear clean clothes daily. Wash thoroughly after handling or applying.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Malathion

Insecticide Group – 1B
(Refer to page 456)

Company:

United Agri Products (Malathion 85E – PCP#8372)
IPCO (Malathion 500 – PCP#5821)

Different companies produce malathion. Note differences in label registrations, formulations and recommendations. Check your label for more information.

Formulations:

Malathion 500 – 500 g/L malathion formulated as an emulsifiable concentrate

Malathion 85E – 85% malathion formulated as an emulsifiable concentrate.

Insects Controlled and Registered Crops:

CROP OR STRUCTURE	INSECT
Alfalfa	Grasshopper, aphid, lygus bug, alfalfa weevil larvae, leafhoppers, alfalfa blotch leafminer, spider mites, spittle bugs
Clover (85E only)	Aphids, grasshoppers, leafhoppers, spider mites
Canola, mustard	Flea beetle, diamondback moth, grasshopper
Wheat, barley, oats, rye	Grasshopper, aphid, armyworm, cereal leaf beetle
Potatoes	Colorado potato beetle, leafhopper, aphid, spider mites
Canaryseed (for seed) (85E only)	Aphids
Sweet clover	Sweet clover weevil
Flax, lentils, hay, pasture	Grasshopper
Corn (grain, forage)	Earworms, European corn borers
Beans, peas	Aphids, leafhoppers, spider mites
Empty bin spray (grain bins, grain elevators, grain box cars, flour mills)	Confused flour beetles, flat grain beetles, granary weevils, grain mites, Indian meal moths, lesser grain borers, red flour beetle, rice weevils, rusty grain beetles, saw-toothed grain beetles

Application:

Malathion may be applied by air or ground equipment. Apply when insect numbers exceed economic threshold levels using sufficient water for good coverage. Use higher rates for heavy infestations, dense canopy or mature stages of insects.

How it Works:

Malathion is a non-systemic, contact, organophosphate insecticide and acaricide of brief to moderate persistence. Generally non-phytotoxic.

Effects of Weather:

For best results apply when daytime temperatures are above 20°C.

Restrictions:

Grazing: When spraying forages and pastures, cattle should be removed and returned after spraying.

Storage: DO NOT store near food or feed. Store in a cool dry place but not below -10°C. Protect from heat.

Others: Maximum of 2 applications per season. DO NOT apply to any plant in bloom. Apply to crops when bees are absent from field. Avoid contact with automobile paint and wash immediately if exposure occurs.

Precautions:

Malathion has a low acute mammalian toxicity. Highly toxic to bees and fish. Wear protective clothing to reduce skin and eye exposure.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Movento 240 SC

Insecticide Group – 23

(Refer to page 456)

Company:

Bayer Crop Science (PCP#28953)

Formulations:

Spinetoram formulated as a suspension concentrate - 240 g/L

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Aphids
Beans, chickpea, lentil, peas, soybean	Aphids

Application:

Ground application only in potatoes and soybeans. Ground or air application for beans, chickpea, lentil and peas. Apply in adequate water for uniform coverage, a minimum of 120 L/ac. If needed repeat application with a minimum of 7 to 10 day interval. DO NOT exceed a maximum of 292 mL / acre per season.

For best results apply when insect populations begin to build and before a damaging population becomes established. Select the appropriate rate depending on the development stage of the insect and level of infestation.

How it Works:

Movento is a systemic, tetracarboxylic acid insecticide. Following

application to plant foliage *Movento* moves through phloem and xylem to all plant tissues including new shoot, leaf and root growth. Mode of action is primarily by ingestion by immature insect life stages.

Restrictions:

DO NOT apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, creeks, marshes, streams, reservoirs and wetlands). DO NOT apply during periods of dead calm. Avoid application when winds are gusty. DO NOT apply droplets smaller than American Society of Agricultural Engineers (ASABE) fine classification. Boom height must be 60 cm or less above ground.

Re-Entry: DO NOT enter or allow worker entry into treated areas for a period of 12 hours.

Re-cropping: A plant back interval of 30 days is required for all crops not on the label.

Precautions:

Movento is toxic to bees through direct contamination of pollen and nectar. DO NOT apply this product during crop flowering period or when flowering weeds are present in the field.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Nolo Bait

Company:

M&R Durango, Inc. (PCP# 29197)

Formulations:

Wheat bran coated with spores of the protozoan *Nosema locustae*.

Minimum of 2.2×10^6 spores of *Nosema (Paranosema) locustae* Canning per gram.

Insects Controlled and Registered Crops:

CROP	INSECT
Crop and Rangeland	Grasshoppers

Rates:

Apply at a minimum rate of 0.45 kg per acre.

Consumption of a higher number of spores per grasshopper will increase product efficacy and decrease the amount of time required to kill grasshoppers. Where greater efficacy or faster population reduction is required, this may be achieved through multiple applications or a higher application rate to increase the amount of bait available to each grasshopper.

Application:

For best results, apply when most grasshoppers are in the 3rd instar (12 to 19 mm long).

Apply by hand, seed spreader, turbine spreader or airplane. Concentrate the application in areas of heaviest grasshopper infestation.

Insecticide Group – Biological Insecticide (Refer to page 456)

How it Works:

Nolo Bait must be consumed by the target insect to be effective. It infects the fat bodies of most species of grasshoppers and some crickets. Infection and sickness of the grasshopper begins upon ingestion of the bait by the grasshopper. As the *Nosema locustae* population increases inside the grasshopper it becomes lethargic, reduces its feeding and has lowered reproductive capacity. Grasshopper death will begin in 3 to 6 weeks. The pathogen may remain in the grasshopper population for several years following treatment.

Restrictions:

Preharvest interval: 0

Storage: Store product in original container in a cool, dry location (preferably at or below 20°C). Use within 13 weeks from the date of manufacture.

Precautions:

May cause sensitization. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Hazard Rating:



Caution – Potential sensitizer

Wheat allergen

For an explanation of the symbols used here see page 11.

Oberon

Company:

Bayer CropScience Inc. (PCP# 28905)

Formulations:

240 g/L spiromesifen formulated as a suspension concentrate
Container size - 2 L jug

Insects Controlled and Registered Crops:

CROP	INSECT
Alfalfa (seed production only)	Two-spotted spider mite
Corn	Banks grass mite, two-spotted spider mite

Application:

May be applied by ground or air.

Apply as soon as mite populations reach threshold levels. Repeat application if pest populations recover and reach economic thresholds. A minimum interval of 7 days between applications is required.

Thorough coverage of all plant parts is important for optimum performance. Use sufficient water volumes for thorough coverage - i.e. minimum of 40 to 80 litres of water per acre.

Avoid application when heavy rain is forecast.

How it Works:

Spiromesifen is in the Tetric acid class of insecticides and works by contact, inhibiting lipid biosynthesis in the insect. *Oberon* has strong adhesion to the leaf surface, and also some translaminar activity providing residual control through contact or ingestion. *Oberon* has activity on all mite developmental stages. Immature mite stages tend to be more susceptible to *Oberon* than adults.

Restrictions:

Alfalfa - DO NOT exceed 3 applications per season. Keep a minimum interval of 7 days between applications. DO NOT exceed a maximum of 1200 ml per acre of *Oberon* per season. Corn - DO NOT exceed 2 applications per season. DO NOT exceed 240 mL per acre per 14 day interval. DO NOT exceed 480 mL per acre per season.

DO NOT enter or allow entry into treated areas for a period of 12 hours after application.

Insecticide Group - 23

(Refer to page 456)

Oberon is toxic to aquatic organisms and beneficial insects such as pollinators. DO NOT apply this product directly to freshwater habitats such as lakes, rivers, sloughs, ponds, creeks, marshes, streams, reservoirs, ditches and wetlands.

Buffer Zones:

Application method	Buffer Zones (metres [†]) Required for the Protection of:		
	Aquatic Habitats of Depths		Terrestrial habitat
	Less than 1 m	Greater than 1 m	
Ground	10	3	2
Fixed wing airplane	800	100	85

See page 29 for an explanation of the different habitats.

- Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

DO NOT mix, load or clean equipment within 30 metres of wellheads or aquatic systems.

Rotational plant-back intervals for:

Field corn - immediate plant back
Wheat, barley and alfalfa - 30 days
All other crops - 12 months

Precautions:

Apply *Oberon* insecticide during pre-bloom, early bloom or late bloom when pollinators are not present in the field. DO NOT apply *Oberon* when pollinators are actively foraging in the treatment area.

Storage: Store in a cool, dry place in such a manner to prevent cross contamination with other pesticides, fertilizers, food and feed.

DO NOT store below freezing.

Hazard Rating:



Caution - Poison

Eye irritant

For an explanation of the symbols used here see page 11.

Orthene

Insecticide Group – 1B

(Refer to page 456)

Company:

Distributed by United Agri Products (PCP#14225)

Formulation:

75% acephate as a water soluble powder. Container size-case of 12 x 1.5 kg

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Green peach aphid, potato aphid, potato flea beetle, potato leafhopper, tarnished plant bug

Preharvest Interval

DO NOT apply within 21 days of harvest.

Application:

Apply with conventional ground equipment only. DO NOT apply by air.

Apply only when insects exceed economic thresholds.

Use higher rate only for heavy infestations.

How it Works:

Acephate is an organophosphate systemic insecticide that works through contact and as a stomach poison.

Effects of Weather:

DO NOT apply if rainfall is expected within 48 hours after application. Treatment areas should not be irrigated for at least 48 hours after application.

Restrictions:

Storage: Store in cool, dry place, in the original container away from food or feed. Protect from excessive heat.

Environment: *Orthene* has the potential to leach through soils to ground water. It is recommended that *Orthene* is not used on coarse textured soils or in areas where the water table may be high.

DO NOT feed foliage to livestock or allow animals to graze on treated areas.

DO NOT make more than 4 applications per season.

Others: *Orthene* is not registered in the United States. Therefore *Orthene* should not be used on any produce destined for markets in the United States.

Precautions:

Orthene is toxic to birds and mammals.

Orthene is toxic to aquatic invertebrates. DO NOT apply where runoff or drift is likely to occur.

Orthene is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. DO NOT apply this product or allow it to drift to blooming crops or weeds if bees are foraging in the treatment area. Beekeepers should be warned to protect bees from treated areas for one week after treatment.

First Aid:

If swallowed, induce vomiting and obtain medical attention or call a poison control centre immediately. In case of contact with skin, wash with soap and water. If in eyes, flush with water. See a physician if eye irritation persists. Atropine is an antidote.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Permethrin

Insecticide Group – 3A
(Refer to page 456)

Company:

FMC of Canada (*Pounce* – PCP#16688)
(Distributed by United Agri Products Canada Inc.)
United Phosphorous (*Perm-UP* – PCP#28877)
Amvac Chemical Corporation (*Ambush* – PCP#14882)

Formulations:

Pounce, *Perm-UP* - 384 g/L permethrin formulated as an emulsifiable concentrate. Container size – 1 L (12 x 1L).
Ambush - 500 g/L permethrin formulated as an emulsifiable concentrate. Container size - 1 L, 5 L

Insects Controlled and Registered Crops:

CROP	INSECT
Cereals, corn, flax, lentil, pea, potato, sunflowers	Cutworms
Canola, rapeseed	Cutworms, crucifer flea beetle
Potato	Colorado potato beetle, potato flea beetle, potato leafhopper, tarnished plant bug, variegated cutworm, European corn borer

Application:

Permethrin may be applied by ground or air. Apply when insects exceed economic threshold numbers and use sufficient water for good coverage. Use higher rates for heavy infestations, adult insects and dense foliage. For cutworm control application should be made under warm, moist conditions in the evening or at night. Use high rates if larvae are near maturity or soil conditions are dry. Do not disturb soil surface for five days after treatment.

How it Works:

Permethrin is a synthetic pyrethroid insecticide. It is a stomach and contact insecticide with no systemic or fumigant effects.

Restrictions:

Grazing: Cover crops or crops treated with permethrin should not be used as a green feed for animals.

Storage: Store above 0°C.

Others: Observe a 16 yard (15 m) setback distance for ground and 110 yard (100 m) setback distance by air near water bodies or other sensitive areas.

Precautions:

Permethrin is of low acute mammalian toxicity. Permethrin is very toxic to bees and fish. DO NOT contaminate ponds, lakes or streams while filling sprayer or spraying. Avoid spraying when bees are foraging.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Rimon 10 EC

Insecticide Group – 15
(Refer to page 456)

Company:

Chemtura (PCP#28881)

MANA Canada (PCP#28515)

Formulations:

10% novaluron formulated as an emulsifiable concentrate

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, European corn borer
Alfalfa (MANA formulation only)	Lygus bug nymphs

Application:

For ground application only.

A minimum spray volume of 40 litres per acre should be used with ground sprayer equipment. Higher water volumes will provide better coverage and product performance.

Use hollow cone, disc-core hollow cone or twin jet nozzles suitable for Insecticide spraying. Drop nozzles may be required to obtain uniform coverage against certain insect pests that develop down in the canopy. Use higher application rates and spray volumes for higher insect pressure.

Colorado potato beetle - Application should be made when the majority of the population is at egg hatch to the second instar of larval development.

European corn borer - Scout for European corn borer to monitor egg-laying and egg hatch to determine timing of application. The first application should be made just prior to egg hatch.

Re-application on a 10 to 14 day interval will be required to protect new growth or if monitoring indicates that it is necessary to keep pest populations below economic thresholds.

How it Works:

Rimon 10 EC is an insect growth regulator that must be absorbed by eggs or ingested by insect larvae to be fully effective. The primary mode of action is by disrupting cuticle formation and deposition occurring when insects

change from one developmental stage to another, resulting in death at molting. Due to this mode of action *Rimon 10 EC* does not have any effect on adult stages of insects that have completed larval development.

Restrictions:

DO NOT make more than 2 applications per year per crop per season.

DO NOT apply more than 656 ml of *Rimon 10 EC* per acre per season.

DO NOT apply within 14 days of harvest (Preharvest interval).

Precautions:

DO NOT re-enter treated areas for a period of 12 hours after application.

Rimon 10 EC is toxic to aquatic organisms. DO NOT apply *Rimon 10 EC* directly to water or to areas where surface water is present. An untreated buffer zone between the last spray swath and the edge of aquatic systems (such as rivers, streams, lakes, and other water bodies) must be established. Refer to label for specific buffer zone requirements.

Rimon 10 EC is toxic to immature insects. Minimize spray drift in habitats next to the application site (e.g. hedgerows and wood-lands) to reduce harmful effects on beneficial insects.

Rimon 10 EC may be toxic to bee colonies exposed to direct treatment, drift or residues on flowering crops or weeds.

Avoid applying this product to flowering crops or weeds if bees are visiting the treatment area.

Storage: To prevent contamination, store this product away from food or feed.

If this product is to be applied to a product destined for export to the United States, information on acceptable residue levels are available at www.croplife.ca

Hazard Rating:

⚠ Warning - May cause substantial but temporary eye injury. Harmful if absorbed through skin. DO NOT get on eyes or clothing.

Keep out of Reach of Children.

For an explanation of the symbols used here see page 11.

Sevin XLR Plus

Insecticide Group – 1A
(Refer to page 456)

Company:

Tessenderlo Kerley, Inc. (PCP#19531)

Formulation:

480 g/L carbaryl formulated as a liquid suspension

Insects Controlled and Registered Crops:

CROP	INSECT
Canola	Flea beetle
Alfalfa, clover	Grasshoppers, blister beetle, leafhoppers, alfalfa caterpillar, armyworm
Barley, oats, rye,	True armyworm, grasshoppers
Beans	Leafhoppers, lygus bugs, climbing cutworms
Ditchbanks, field borders, forage grasses, headlands, pastures, rangelands, rights-of-way, wastelands	Grasshoppers
Corn	European corn borer, corn earworm, fall armyworm, grasshoppers
Potato	Colorado potato beetle, flea beetle, leafhopper, European corn borer, climbing cutworms

Application:

May be applied by air or ground equipment. Apply when insect numbers exceed economic threshold levels and use sufficient water for good coverage. Use lower rates on young plants and early stages of insects and higher rates on mature plants and advanced stages of insects, or mature insects.

Sevin XLR Plus applications are more resistant to wash-off when applied as a concentrated suspension. To ensure wash-off resistance is retained, apply dilutions of 1 part *Sevin XLR Plus* to no greater than 39 parts water. For example, if applying 1 L/ac of *Sevin XLR Plus* DO NOT exceed 40 L/ac total application volume. Applications should be made to dry foliage to maximize wash-off resistance.

How it Works:

Sevin XLR Plus is a carbamate insecticide that works by contact (approx. 20%) and ingestion (approx. 80%). Moderate to rapid in speed of action with short to moderate residual activity (2 to 4 weeks).

Restrictions:

Storage: DO NOT store in areas where temperatures frequently exceed 38°C. Store in original container in a cool dry area out of reach of children and animals and away from food and feed.

Grazing: Remove cattle from area when spraying. Crops may be grazed or harvested for feed: Dairy animals - 48 hours after treatment; Meat animals - 24 hours after treatment.

Precautions:

Sevin XLR Plus is of moderate acute mammalian toxicity. This product is highly toxic to honeybees exposed to direct treatment on blooming crops or weeds. Harmful if inhaled or swallowed. Avoid contact with skin and eyes. Wear long-sleeve work clothing and change to clean clothing daily. Wash hands and face after handling. Avoid contamination of food, feed, water supplies, streams and ponds.

Hazard Rating:



Warning – Poison

For an explanation of the symbols used here see page 11.

Sluggo Professional

Molluscicide – no group

Company:

Engage Agro Corporation (PCP#30025)

Formulation:

0.76 % ferric phosphate in a granular formulation

Container size - 5, 25 kg bags

Pest Controlled and Registered Crops:

CROP	INSECT
Field crops	Slugs and snails

Rates:

Apply bait evenly at a rate of 10.1 to 20.2 kg / acre (2.5 to 5 g per square metre).

Application:

Apply in the evening as slugs and snails travel and feed mostly at night or early morning. DO NOT place in piles. For best results the ground should be moist but with little or no standing water.

For broadcast application, standard broadcast spreaders may be used. For row application, standard granular spreaders may be used.

At seeding and later stages, apply the bait between rows and around the perimeter of the field. Treating around the perimeter of crop areas may intercept slugs or snails migrating from daytime refuge sites.

Apply at the higher rate within the recommended rate range if the infestation is severe, if the area is heavily watered or after long periods of heavy rain.

Re-apply as the bait is consumed or at least every two weeks if slugs and snails continue to be a problem.

How it Works:

Sluggo must be consumed by the slugs or snails to be effective. After ingesting the bait, slugs and snails stop feeding providing immediate protection to plants. Affected slugs and snails die within 3 to 6 days.

Restrictions:

Avoid direct application to ponds streams and lakes. This product may be toxic to fish and other aquatic organisms.

Precautions:

Avoid contact with eyes. May cause eye irritation.

Wear chemical resistant gloves during mixing and loading activities and when applying by hand.

Hazard Rating:



Warning – contains the allergen wheat

For an explanation of the symbols used here see page 11.

Success 480 SC

Insecticide Group – 5

(Refer to page 456)

Company:

Dow AgroSciences (PCP#26835)

Formulation:

480 g/L spinosad formulated as a suspension concentrate
Container size - 1L jug

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle larvae and European corn borer larvae

Application:

Apply *Success 480 SC* as a foliar spray by ground only. DO NOT apply by air. Apply when scouting indicates the target pest species have reached economic threshold levels. For Colorado potato beetle, target eggs at hatch or small larvae. For control of European corn borer, time the application to coincide with peak egg hatch. Use higher application rate for higher pest pressure or when extended egg hatch is anticipated. If pest populations persist, a repeat application 7 to 10 days after the initial application may be necessary.

How it Works:

Success 480 SC is in the spinosyn class of insecticides. It is a contact and stomach insecticide. It is derived from the fermentation of *Saccharopolyspora spinosa*.

Effects of Weather:

This product has the potential for run-off. DO NOT spray immediately after a rainfall or if rain is forecast within 48 hours after application.

Restrictions:

Storage: Avoid freezing. DO NOT store or ship with food, feeds, drugs or clothing.

Others:

Potatoes - DO NOT apply more than a maximum seasonal rate of 100 ml/acre.

DO NOT apply within 7 days of harvest.

DO NOT enter or allow worker entry into treated areas for a period of 4 hours after application.

Precautions:

Success 480 SC is highly toxic to honey bees exposed to direct spray. Avoid spraying on crops in bloom or other vegetation when bees are actively foraging. This product is toxic to other beneficial parasites and predators. DO NOT contaminate aquatic habitats such as ponds, lakes, rivers, streams and wetlands during application or when rinsing equipment and containers.

Buffer Zones: A buffer zone of 2 metres (early season) or 1 metre (late season) is required between downwind edge of spray boom and sensitive aquatic habitats. Avoid contact with eyes, skin, and clothing.

Hazard Rating:



Caution – Poison

For an explanation of the symbols used here see page 11.

Thimet 15G

Insecticide Group – 1B

(Refer to page 456)

Company:

Amvac Chemical Corporation (PCP#10532)

Note:

Last date of sale of Thimet 15-G by retailers and distributors - 1 May 2015

Last date of use of Thimet 15-G by growers and users - 1 August 2015

Formulations:

15% phorate formulated as a granular.
Container size- 20 kg bag.

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Reduction of wireworm damage.

Application:

Ground application at seeding time: Distribute granules evenly in furrow or granules may be banded on each side of the row at planting time. Use low rate for sandy or light soils and high rate for silt or heavy soils. For use only in hoppers equipped for Lock 'N Load containers to maintain a closed system.

How it Works:

Phorate is an organophosphate insecticide that works as a systemic poison, with effective initial residual activity on soil and foliar insects.

Restrictions:

Storage: DO NOT store in or around the home. Store away from food or feed. Store open bags in labeled sealed drums or heavy plastic bags.

Others: DO NOT use in muck soils. DO NOT apply later than at planting time. Will provide reduction of wireworm damage.

Precautions:

Thimet is of high acute mammalian toxicity and is highly toxic to fish, birds and other animals. DO NOT allow product to contact eyes and skin. Poisonous by skin contact, inhalation or swallowing. Repeated inhalation or skin contact with Thimet 15G, other organophosphorus or carbamate insecticides may, without symptoms, progressively increase susceptibility to poisoning. Wear freshly-laundered, long-sleeved work clothing daily. Use rubber gloves when transferring from package to equipment. Sleeve cuffs should be worn over gloves to prevent granules from falling into the gloves. Rubber gloves should be washed with soap and water after each use. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Vydate L

Insecticide Group – 1A
(Refer to page 456)

Company:

E.I. duPont Canada (PCP#17995)

Formulation:

240 g/L oxamyl

Container size - 10 L jugs

Insects Controlled and Registered Crops:

CROP	INSECT
Potato	Colorado potato beetle, flea beetles, aphids, potato leafhopper, Lygus bugs

Application:

Apply *Vydate* as a foliar spray by ground only. Apply sufficient volume of spray solution to thoroughly wet foliage (minimum water volume 120 to 360 L per acre). Make application when insects appear at economic levels. Repeat weekly as required. Use low rate for light infestations. Use higher rate for severe infestations or if aphids are the primary pest species.

Avoid application when heavy rain is forecast.

How it Works:

Vydate is a carbamate insecticide that works by contact and ingestion.

Vydate will not control Colorado potato beetles resistant to other carbamate insecticides.

Restrictions:

DO NOT apply *Vydate* by air.

DO NOT apply within 7 days of harvest.

DO NOT apply directly to water. DO NOT apply where run-off is likely to occur. DO NOT apply when weather conditions favour drift to untreated areas and when wind speed exceeds 10 km/hr.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval of 3 days.

DO NOT apply more than 2 times per season.

Storage: DO NOT freeze. DO NOT store or transport with food, feeds, drugs or clothing.

Buffer zones:

Application method	Crop	Buffer Zones (metres [†]) Required for the Protection of:		
		Freshwater habitat of depths:		
		Less than 1 m	1 – 3 m	Greater than 3 m
Ground	Potato	2	1	0

See page 29 for an explanation of the different habitats.

- Buffer zones can be reduced by 70% when using shrouds and by 30% when using cones mounted less than 12 inches from the crop canopy.
- For tank mixes, consult the labels of the tank mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

[†] Distance measured as metres from the downwind edge of the spray boom to sensitive habitat.

Precautions:

Vydate is toxic to fish, birds and other wildlife.

Vydate is toxic to honey bees exposed to direct spray and should not be used when bees are active in the treatment area. Avoid contact with eyes, skin, and clothing.

Hazard Rating:



Danger – Poison

For an explanation of the symbols used here see page 11.

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes

Notes